



EXPLORING RYTARY

STARTING RYTARY

MYRYTARY PROGRAM

THE RYTARY COMMUNITY

Learn about the RYTARY Co-Pay Savings Card



Still got passion. Still got pride. Still got purpose.

Still got it.

Meet Denise. She's living with Parkinson's disease, but it doesn't define her. That's because she takes RYTARY, which means she has less "off" time, and more time to do what she loves.

WATCH HER RYTARY STORY

INDICATION

RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Call your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects.

SEE MORE +

Start saving with the RYTARY Co-Pay Savings Card

If you have commercial insurance, you may be eligible to:

Pay no more than \$25
for your first RYTARY prescription*

Pay as little as \$0
if you're written a second or third prescription of RYTARY within that same 30-day period†



GET STARTED

Please see eligibility criteria below.

INDICATION


RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Call your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects.


SEE MORE +

Get inspired by Stories of Parkinson's (SPARK) Ambassadors




Nancy—Still got grace
On RYTARY since 2018

"My doctor upped my starting dose,



Daran—Still got smiles
On RYTARY since 2015

"Because I have less 'off' time with



Denise—Still got passion
On RYTARY since 2016

"My doctor explained that my

INDICATION

RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

SEE MORE +

...upped the dose. It felt like my toes were dancing when I hadn't asked them to dance. So, we went back to my original dose but upped the frequency to 4 times a day rather than 3 times. Now I have less 'off' time, which makes me and the family happy."

...over the night I found with my Parkinson's diagnosis."

...may have been a bit slower, but to make sure I kept him informed on what was happening. So I tried it again. I stuck with it, and he was right. My side effects improved, and I was happy with my results with RYTARY."

Individual results may vary.

Get connected with others

Sign up for an online or local event in your area, and:

- Get informed about ways to help manage Parkinson's disease from a healthcare provider
- Get inspired by SPARK Ambassadors
- Get to know others in the Parkinson's community



INDICATION

RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

SEE MORE +

Get RYTARY — a Parkinson's disease medication that can help the real you shine through

Taking RYTARY may result in:

2X reduction in "off" time compared with Immediate-Release (IR) CD/LD

INDICATION
RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION
Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

[SEE MORE +](#)

[FIND EVENTS](#)



Get information to help you on your journey



Sign up to receive ongoing communications, including a free RYTARY Information Kit

[SIGN UP](#)

INDICATION
RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson's disease, Parkinson's disease caused by infection or inflammation of the brain, or Parkinson's disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION
Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

[SEE MORE +](#)

Get RYTARY — a Parkinson’s disease medication that can help the real you shine through

Taking RYTARY may result in:

2X reduction in “off” time compared with Immediate-Release (IR) CD/LD

In a clinical study, people with advanced Parkinson’s disease taking RYTARY experienced a 13.1% reduction in “off” time during waking hours (compared with only 6.2% for people taking IR CD/LD).[†]

INDICATION

RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson’s disease, Parkinson’s disease caused by infection or inflammation of the brain, or Parkinson’s disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

SEE MORE +

- RYTARY at study start and study end[‡]: 36.9% to 23.8%
- IR CD/LD at study start and study end[‡]: 36.0% to 29.8%

LEARN MORE

“You are *not* your disease.
You are a unique individual who happens to have Parkinson’s.
So let your story be heard.
Be your own health advocate.”

— Denise, on RYTARY since 2016

*Maximum benefit of \$100.

[†]This card is not valid for prescriptions submitted for reimbursement to Medicare, Medicaid, other federal or state programs (including any state pharmaceutical assistance programs), or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drug. This card is good for one RYTARY prescription of the type the prescription is filed by the pharmacist and dispensed to the patient. Offer good only in the US.

INDICATION

RYTARY is a prescription medication that contains a combination of carbidopa and levodopa for the treatment of Parkinson’s disease, Parkinson’s disease caused by infection or inflammation of the brain, or Parkinson’s disease like symptoms that may result from carbon monoxide or manganese poisoning.

IMPORTANT SAFETY INFORMATION

Do not take RYTARY with antidepressant medications known as nonselective monoamine oxidase (MAO) inhibitors because taking these two drugs within two weeks of each other can result in high blood pressure. Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness. Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle

SEE MORE +

Taking RYTARY may result in falling asleep while engaged in normal activities, even without warning and as late as one year after starting to take RYTARY. Other sedating medicines and alcohol taken together with RYTARY may have additional sedative effects. Tell your healthcare provider if you have any kind of sleep disorder or are experiencing drowsiness or sleepiness.

Some side effects of taking RYTARY including sleepiness and dizziness may affect your ability to drive or operate machinery. Do not drive a car, operate a machine, or do anything that requires you to be alert until you know how RYTARY affects you. Talk to your healthcare provider before you lower the dose or stop taking RYTARY, as this may result in serious side effects. Call your healthcare provider immediately if you develop withdrawal symptoms such as fever, confusion, or severe muscle stiffness.

Make sure to tell your healthcare provider if you have any heart conditions, especially if you have had a heart attack and also have irregular heartbeats. Some people with a history of or risk factors for heart disease have experienced heart problems while taking RYTARY.

Some patients taking RYTARY can experience hallucinations (unreal visions, sounds, or sensations) or abnormal thoughts and behaviors (such as excessive suspicion, believing things that are not real, confusion, agitation, aggressive behavior, and disorganized thinking). If you have hallucinations or abnormal thoughts or behaviors, talk to your healthcare provider.

Some patients taking certain medicines to treat Parkinson's disease have intense urges to gamble, increased sexual urges, other intense urges, and the inability to control those urges. If you or your family members notice that you are developing unusual urges or behaviors, talk to your healthcare provider.

Tell your healthcare provider if abnormal involuntary movements appear or get worse during treatment with RYTARY.

Tell your healthcare provider if you have ever had a peptic ulcer, because RYTARY may increase your chances of having bleeding in your stomach.

Tell your healthcare provider if you have glaucoma, because RYTARY may increase the pressure in your eyes.

Parkinson's disease patients are at an increased risk of developing melanoma, a form of skin cancer. See your healthcare provider for regular skin examinations when taking RYTARY.

The most common side effects that may occur with RYTARY include nausea, dizziness, headache, sleeplessness, abnormal dreams, dry mouth, abnormal involuntary movements, anxiety, constipation, vomiting, and low blood pressure upon rising. Rise slowly after sitting or lying down for a prolonged period.

Following use in the marketplace, some patients taking RYTARY have experienced suicidal thoughts or have attempted suicide. A causal relationship has not been established. Tell your healthcare provider if you have thoughts of suicide or have attempted suicide.

Notify your healthcare provider if you become pregnant or intend to become pregnant during therapy or if you intend to breast-feed or are breast-feeding an infant.

Adverse events following unintentional overdose with this medication have been reported. If you accidentally take more than your prescribed dose, talk to your healthcare provider right away.

Make sure you tell your healthcare provider about all of the prescription and non-prescription medications you take, including supplements, and especially those for Parkinson's disease, heart disease, blood pressure, abnormal thoughts, tuberculosis, and sleep problems, and supplements containing iron. Do not take other carbidopa/levodopa preparations with RYTARY without consulting your healthcare provider.

Be sure to take your medicine as instructed. You may take RYTARY with or without food; however, taking RYTARY with food may decrease or delay its effect. For this reason, consider taking the first dose of the day about 1 to 2 hours before eating. Swallow RYTARY whole; do not chew, divide, or crush. If you have difficulty swallowing the capsule, twist apart both halves and sprinkle the entire contents of both halves of the capsule on a small amount of applesauce (1 to 2 tablespoons). Consume the mixture immediately. Do not store the drug/food mixture for future use.

Note: The above information for patients being treated with RYTARY is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects. Tell your healthcare provider if you have any side effects while taking RYTARY. He or she can make adjustments that may reduce these effects.

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Specialty, a division of Amneal Pharmaceuticals LLC at 1-877-835-5472 or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please read the [Full Prescribing Information](#). For more information talk to your healthcare provider.

The individuals appearing in this piece were sponsored by Amneal Pharmaceuticals. Your experience and results may vary.



RYTARY® and MYRYTARY® are registered trademarks of Impax Laboratories, LLC.
© 2019 Amneal Pharmaceuticals LLC.
Distributed by Amneal Specialty, a division of Amneal Pharmaceuticals LLC. All rights reserved.
PP-RYT-US-0093 11/2019



- Contact Us
- Legal Notices
- Amneal Specialty
- Privacy Policy
- Full Prescribing Information
- Site Map

This is intended for residents of the United States only. Any product discussed herein may have different product labeling in different countries. Please read the [Full Prescribing Information](#) and discuss it with your doctor or healthcare professional.