



US 20200248174A1

(19) **United States**

(12) **Patent Application Publication**
Nishida et al.

(10) **Pub. No.: US 2020/0248174 A1**

(43) **Pub. Date: Aug. 6, 2020**

(54) **GENOMIC SEQUENCE MODIFICATION
METHOD FOR SPECIFICALLY
CONVERTING NUCLEIC ACID BASES OF
TARGETED DNA SEQUENCE, AND
MOLECULAR COMPLEX FOR USE IN
SAME**

Publication Classification

(51) **Int. Cl.**
C12N 15/10 (2006.01)
C12N 9/22 (2006.01)
C12N 9/78 (2006.01)
C12N 15/11 (2006.01)

(52) **U.S. Cl.**
 CPC *C12N 15/1024* (2013.01); *C12N 9/22*
 (2013.01); *C12N 15/102* (2013.01); *C12N*
9/78 (2013.01); *C12N 2310/3513* (2013.01);
C12Y 305/04005 (2013.01); *C07K 2319/00*
 (2013.01); *C07K 2319/81* (2013.01); *C12N*
15/11 (2013.01)

(71) Applicant: **NATIONAL UNIVERSITY
CORPORATION KOBE
UNIVERSITY, Kobe-shi (JP)**

(72) Inventors: **Keiji Nishida, Kobe-shi, Hyogo (JP);
Akihiko Kondo, Kobe-shi, Hyogo (JP);
Satomi Kojima, Kobe-shi, Hyogo (JP)**

(21) Appl. No.: **16/838,960**

(22) Filed: **Apr. 2, 2020**

Related U.S. Application Data

(63) Continuation of application No. 15/124,021, filed on
Nov. 9, 2016, now Pat. No. 10,655,123, filed as
application No. PCT/JP2015/056436 on Mar. 4, 2015.

Foreign Application Priority Data

Mar. 5, 2014 (JP) 2014-043348
Sep. 30, 2014 (JP) 2014-201859

(57) **ABSTRACT**

The invention provides a method of modifying a targeted site of a double stranded DNA, including a step of contacting a complex wherein a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in a selected double stranded DNA and a nucleic acid base converting enzyme are linked, with the double stranded DNA, to convert one or more nucleotides in the targeted site to other one or more nucleotides or delete one or more nucleotides, or insert one or more nucleotides into the targeted site, without cleaving at least one strand of the double stranded DNA in the targeted site.

Specification includes a Sequence Listing.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

Change of mutation introduction pattern

target sequence AAATGGGAGGATACGTTCTGTATGGAGCAAGGATAGGT (SEQ ID NO:46)
 TTACCGCTGCTATGGAGAGATACCTCCAGCGTATCGA (SEQ ID NO:47)

187

target 1





Fig. 7




Fig. 8




Fig. 10A




Fig. 10B




Fig. 11A




Fig. 11B

galK TCAATGGGCTAACTACGTTTC-3' (SEQ ID NO:61)
 1 TCAATGGGCTAACTACGTTTC (SEQ ID NO:61)
 2 TTAATGGGCTAACTACGTTTC (SEQ ID NO:62)
 3 TTAATGGGCTAACTACGTTTC (SEQ ID NO:62)

Fig. 11C

xpoB GCTGTCTCAGTTTATGGACC-3' (SEQ ID NO:63)
 [CGACAGAGTCAAATACCTGG-5'] (SEQ ID NO:64)

none | 1 GCTGTCTCAGTTTATGGACC (SEQ ID NO:63)
 2 GCTGTCTCAGTTTATGGACC (SEQ ID NO:63)

Rif25 | 1 GCTGTCTCAGTTTATGAACC (SEQ ID NO:66)
 2 GCTGTCTCAGTTTATGAACC (SEQ ID NO:66)

Rif50 | 1 GCTGTCTCAGTTTATAAACC (SEQ ID NO:67)
 2 GCTGTCTCAGTTTATGAACC (SEQ ID NO:66)
 [CGACAGAGTCAAATACTTGG-5'] (SEQ ID NO:68)




Fig. 12A




Fig. 12B

- TCGCTTGAACATCCAGCGAAACAGGC CCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 69)
- target 24bp
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 70)
- TCGCTTGAACATCCAGCGAAACAGGC CaCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 71)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 72)
- TCGCTTGAACATCCAGCGAAACAGGC CaCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 73)
- TCGCTTGAACATCCAGCGAAACAGGC CCaCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 74)
- target 22bp
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 75)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 76)
- TCGCTTGAACATCCAGCGAAACAGGC CCaCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 77)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 78)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 79)
- target 20bp
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 80)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 81)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 82)
- TCGCTTGAACATCCAGCGAAACAGGC CCaCCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 83)
- target 18bp
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 84)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 85)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 86)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 87)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 88)
- TCGCTTGAACATCCAGCGAAACAGGC CCCCATCGAGCAGAAA CCGTGGTGGATGGC (SEQ ID NO: 89)

Fig. 13




Fig. 14

mutation introduction method (temperature sensitive plasmid)





Fig. 15



rpoB/r GCCTGCTCAGTTTATGGACC-3' (SEQ ID NO:90)

colony1 GCCTGCTCAGTTTATGAACC (SEQ ID NO:91)

colony2 GCCTGCTCAGTTTATGAACC (c→t) (SEQ ID NO:91)





Fig. 16

galK (target 8, target 12)



galK 8 ACTACACGCAATTCAGGGCGC -3' (SEQ ID NO:92)
 colony1 ACTTACACGCAATTCAGGGCGC (SEQ ID NO:93)
 colony2 ACTTACACGCAATTCAGGGCGC (c→t) (SEQ ID NO:93)

galK 12 TCAATGGGCTAACGACGTTGG -3' (SEQ ID NO:94)
 colony1 TTATGGGCTAACGACGTTGG (SEQ ID NO:95)
 colony2 TTATGGGCTAACGACGTTGG (c→t) (SEQ ID NO:95)

**GENOMIC SEQUENCE MODIFICATION
METHOD FOR SPECIFICALLY
CONVERTING NUCLEIC ACID BASES OF
TARGETED DNA SEQUENCE, AND
MOLECULAR COMPLEX FOR USE IN
SAME**

CROSS-REFERENCE TO RELATED
APPLICATIONS

[0001] This patent application is a continuation of U.S. patent application Ser. No. 15/124,021, filed Nov. 9, 2016; which is the U.S. national phase of International Patent Application No. PCT/JP2015/056436, filed Mar. 4, 2015; which claims the benefit of Japanese Patent Application No. 2014-043348, filed on Mar. 5, 2014, and Japanese Patent Application No. 2014-201859, filed on Sep. 30, 2014, which are incorporated by reference in their entireties herein.

INCORPORATION-BY-REFERENCE OF
MATERIAL ELECTRONICALLY SUBMITTED

[0002] Incorporated by reference in its entirety herein is a computer-readable nucleotide/amino acid sequence listing submitted concurrently herewith and identified as follows: 96.9 KB ASCII (Text) file named "150161_401C1_SEQ_LISTING.txt" created Mar. 30, 2020.

TECHNICAL FIELD

[0003] The present invention relates to a modification method of a genome sequence, which enables modification of a nucleic acid base in a particular region of a genome, without cleaving double-stranded DNA (with no cleavage or single strand cleavage), and without inserting a foreign DNA fragment, and a complex of a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme used therefor.

BACKGROUND ART

[0004] In recent years, genome editing is attracting attention as a technique for modifying the target gene and genome region of interest in various species. Conventionally, as a method of genome editing, a method utilizing an artificial nuclease comprising a combination of a molecule having a sequence-independent DNA cleavage ability and a molecule having a sequence recognition ability has been proposed (non-patent document 1).

[0005] For example, a method of performing recombination at a target gene locus in DNA in a plant cell or insect cell as a host, by using a zinc finger nuclease (ZFN) wherein a zinc finger DNA binding domain and a non-specific DNA cleavage domain are linked (patent document 1); a method of cleaving or modifying a target gene in a particular nucleotide sequence or a site adjacent thereto by using TALEN wherein a transcription activator-like (TAL) effector, which is a DNA binding module that the plant pathogenic bacteria *Xanthomonas* has, and a DNA endonuclease are linked (patent document 2); a method utilizing CRISPR-Cas9 system wherein DNA sequence CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats), that functions in an acquired immune system possessed by eubacterium and archaeobacterium, and nuclease Cas (CRISPR-associated) protein family having an important function along with CRISPR are combined (patent document 3) and the like have been reported. Furthermore, a method of

cleaving a target gene in the vicinity of a particular sequence, by using artificial nuclease wherein a PPR protein configured to recognize a particular nucleotide sequence by a series of PPR motifs each consisting of 35 amino acids and recognizing one nucleic acid base, and nuclease are linked (patent document 4) has also been reported.

DOCUMENT LIST

Patent Documents

- [0006]** patent document 1: JP-B-4968498
- [0007]** patent document 2: National Publication of International Patent Application No. 2013-513389
- [0008]** patent document 3: National Publication of International Patent Application No. 2010-519929
- [0009]** patent document 4: JP-A-2013-128413
- [0010]** non-patent document
- [0011]** non-patent document 1: Kelvin M Esvelt, Harris H Wang (2013) Genome-scale engineering for systems and synthetic biology, *Molecular Systems Biology* 9: 641

SUMMARY OF THE INVENTION

Problems to be Solved by the Invention

[0012] The genome editing techniques heretofore been proposed basically presuppose double-stranded DNA breaks (DSB). However, since they involve unexpected genome modifications, side effects such as strong cytotoxicity, chromosomal rearrangement and the like occur, and they have common problems of impaired reliability in gene therapy, extremely small number of surviving cells by nucleotide modification, and difficulty in genetic modification itself in primate ovum and unicellular microorganisms.

[0013] Therefore, an object of the present invention is to provide a novel method of genome editing for modifying a nucleic acid base of a particular sequence of a gene without DSB or insertion of foreign DNA fragment, i.e., by non-cleavage of a double stranded DNA or single strand cleavage, and a complex of a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme therefor.

Means of Solving the Problems

[0014] The present inventors have conducted intensive studies in an attempt to solve the above-mentioned problems and taken note of adopting base conversion by a conversion reaction of DNA base, without accompanying DSB. The base conversion reaction by a deamination reaction of DNA base is already known; however, targeting any site by recognizing a particular sequence of DNA, and specifically modifying the targeted DNA by base conversion of DNA bases has not been realized yet.

[0015] Therefore, deaminase, that catalyzes a deamination reaction, was used as an enzyme for such conversion of nucleic acid bases, and linked to a molecule having a DNA sequence recognition ability, thereby a genome sequence was modified by nucleic acid base conversion in a region containing a particular DNA sequence.

[0016] Specifically, CRISPR-Cas system (CRISPR-mutant Cas) was used. That is, a DNA encoding an RNA molecule, wherein genome specific CRISPR-RNA:crRNA (gRNA) containing a sequence complementary to a target sequence of a gene to be modified is linked to an RNA for recruiting Cas protein (trans-activating crRNA: tracrRNA)

was produced. On the other hand, a DNA wherein a DNA encoding a mutant Cas protein (dCas), wherein cleavage ability of one or both strands of a double stranded DNA is inactivated and a deaminase gene are linked, was produced. These DNAs were introduced into a host yeast cell which comprises a gene to be modified. As a result, mutation could be introduced randomly within the range of several hundred nucleotides of the gene of interest including the target sequence. Compared to when a double mutant Cas protein, which do not cleave both of DNA strands in the double stranded DNA, was used, the mutation introduction efficiency increased when a mutant Cas protein which cleave of either one of the strands was used. In addition, it was clarified that the area of mutation region and variety of mutation vary depending on which of the DNA double strand is cleaved. Furthermore, mutation could be introduced extremely efficiently by targeting a plurality of regions in the target gene of interest. That is, a host cell introduced with DNA was seeded in a nonselective medium, and the sequence of the target gene of interest was examined in randomly selected colonies. As a result, introduction of mutation was confirmed in almost all colonies. It was also confirmed that genome editing can be simultaneously performed at a plurality of sites by targeting certain region in two or more target genes of interest. It was further demonstrated that the method can simultaneously introduce mutation into alleles of diploid or polyploid genomes, can introduce mutation into not only eukaryotic cells but also prokaryotic cells such as *Escherichia coli*, and is widely applicable irrespective of species. It was also found that editing of essential gene, which showed low efficiency heretofore, can be efficiently performed by transiently performing a nucleic acid base conversion reaction at a desired stage.

[0017] The present inventors have conducted further studies based on these findings and completed the present invention.

[0018] Accordingly, the present invention is as described below.

[0019] [1] A method of modifying a targeted site of a double stranded DNA, comprising a step of contacting a complex wherein a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in a selected double stranded DNA and a nucleic acid base converting enzyme are linked, with said double stranded DNA, to convert one or more nucleotides in the targeted site to other one or more nucleotides or delete one or more nucleotides, or insert one or more nucleotides into said targeted site, without cleaving at least one strand of said double stranded DNA in the targeted site.

[0020] [2] The method of [1], wherein the nucleic acid sequence-recognizing module is selected from the group consisting of a CRISPR-Cas system wherein at least one DNA cleavage ability of Cas is inactivated, a zinc finger motif, a TAL effector and a PPR motif.

[0021] [3] The method of [1], wherein the nucleic acid sequence-recognizing module is a CRISPR-Cas system wherein at least one DNA cleavage ability of Cas is inactivated.

[0022] [4] The method of any of [1]-[3], which uses two or more kinds of nucleic acid sequence-recognizing modules each specifically binding to a different target nucleotide sequence.

[0023] [5] The method of [4], wherein the different target nucleotide sequence is present in a different gene.

[0024] [6] The method of any of [1]-[5], wherein the nucleic acid base converting enzyme is deaminase.

[0025] [7] The method of the above-mentioned [6], wherein the deaminase is AID (AICDA).

[0026] [8] The method of any of [1]- [7], wherein the double stranded DNA is contacted with the complex by introducing a nucleic acid encoding the complex into a cell having the double stranded DNA.

[0027] [9] The method of [8], wherein the cell is a prokaryotic cell.

[0028] [10] The method of [8], wherein the aforementioned cell is a eukaryotic cell.

[0029] [11] The method of [8], wherein the cell is a cell of a microorganism.

[0030] [12] The method of [8], wherein the cell is a plant cell.

[0031] [13] The method of [8], wherein the cell is an insect cell.

[0032] [14] The method of [8], wherein the cell is an animal cell.

[0033] [15] The method of [8], wherein the aforementioned cell is a cell of a vertebrate.

[0034] [16] The method of [8], wherein the cell is a mammalian cell.

[0035] [17] The method of any of [9]- [16], wherein the cell is a polyploid cell, and a site in any targeted allele on a homologous chromosome is modified.

[0036] [18] The method of any of [8]- [17], comprising a step of introducing an expression vector comprising a nucleic acid encoding the complex in a form permitting control of an expression period into the cell, and a step of inducing expression of the nucleic acid for a period necessary for stabilizing the modification of the targeted site in the double stranded DNA.

[0037] [19] The method of the above-mentioned [18], wherein the target nucleotide sequence in the double stranded DNA is present in a gene essential for the cell.

[0038] [20] A nucleic acid-modifying enzyme complex wherein a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in a selected double stranded DNA and a nucleic acid base converting enzyme are linked, which converts one or more nucleotides in the targeted site to other one or more nucleotides or deletes one or more nucleotides, or inserts one or more nucleotides into said targeted site, without cleaving at least one strand of said double stranded DNA in the targeted site.

[0039] [21] A nucleic acid encoding the nucleic acid-modifying enzyme complex of [20].

Effect of the Invention

[0040] According to the genome editing technique of the present invention, since it is not associated with insertion of a foreign DNA or double-stranded DNA breaks, the technique is superior in safety. The technique has some possibility of providing a solution in cases where conventional methods were considered as a gene recombination, and thus biologically or legally controversial. It is also theoretically possible to set a wide range of mutation introduction from a pin point of one base to several hundred bases, and the technique can also be applied to local evolution induction by

introduction of random mutation into a particular limited region, which has been almost impossible heretofore.

BRIEF DESCRIPTION OF THE DRAWINGS

[0041] FIG. 1 is a schematic illustration showing a mechanism of the genetic modification method of the present invention using the CRISPR-Cas system.

[0042] FIG. 2 shows the results of verification, by using a budding yeast, of the effect of the genetic modification method of the present invention comprising a combination of a CRISPR-Cas system and PmCDA1 deaminase from *Petromyza marinus*.

[0043] FIG. 3 shows changes in the number of surviving cells after expression induction when a CRISPR-Cas9 system using a D10A mutant of Cas9 having a nickase activity and a deaminase, PmCDA1, are used in combination (nCas9 D10A-PmCDA1), and when conventional Cas9 having a DNA double strand cleavage ability is used.

[0044] FIG. 4 shows the results when a plurality of expression constructs are constructed such that human AID deaminase and dCas9 are linked via SH3 domain and a binding ligand thereof, wherein the express constructs are introduced into a budding yeast together with two kinds of gRNA (targeting sequences of target 4 and target 5).

[0045] FIG. 5 shows that the mutation introduction efficiency is increased by the use of Cas9 that cleaves either DNA single strand.

[0046] FIG. 6 shows that in the case where a double stranded DNA is not cleaved, the area of mutation introduction region and frequency thereof change depending on which one of the single strands is cleaved.

[0047] FIG. 7 shows that extremely high mutation introduction efficiency can be realized by targeting two regions in proximity.

[0048] FIG. 8 shows that the genetic modification method of the present invention does not require selection by marker. It was found that mutation was introduced into all colonies sequenced.

[0049] FIG. 9 shows that a plurality of sites in a genome can be simultaneously edited by the genetic modification method of the present invention. The upper panel shows the nucleotide sequence and amino acid sequence of the target site of each gene, and an arrow on the nucleotide sequence shows the target nucleotide sequence. The number at the arrow end or arrow head indicates the position of the target nucleotide sequence terminus on ORF. The lower panel shows the results of sequencing of the target site in each 5 clones of red (R) and white (W) colonies. In the sequences, the nucleotides indicated with outline characters show occurrence of base conversion. As for responsiveness to canavanine (Can^R), R shows resistance, and S shows sensitivity.

[0050] FIG. 10 shows that a mutation can be simultaneously introduced into both alleles on the homologous chromosome of diploid genome by the genetic modification method of the present invention. FIG. 10A shows homologous mutation introduction efficiency of *Adel* gene (upper panel) and *can1* gene respectively. FIG. 10B shows that homologous mutation was actually introduced into red colony (lower panel). Also, occurrence of heterologous mutation in white colony was shown (upper panel).

[0051] FIG. 11 shows that genome editing of *Escherichia coli*, a prokaryotic cell, is possible by the genetic modification method of the present invention. FIG. 11A is a sche-

matic illustration showing the plasmid used. FIG. 11B shows that a mutation (CAA→TAA) could be efficiently introduced by targeting a region in the *galK* gene. FIG. 11C shows the results of sequence analysis of each two clones of the respective colonies in a nonselective medium (none), a medium containing 25 μg/ml rifampicin (Rif25) or a medium containing 50 μg/ml rifampicin (Rif50). Introduction of a mutation imparting rifampicin resistance was confirmed (upper panel). The appearance frequency of rifampicin resistance strain was estimated to be about 10% (lower panel).

[0052] FIG. 12 shows control of the edited base sites by the length of guide RNA. FIG. 12A is a conceptual Figure of editing base site when the length of the target nucleotide sequence is 20 bases or 24 bases. FIG. 12B shows the results of editing by targeting *gsiA* gene and changing the length of the target nucleotide sequence. The mutated sites are shown with bold letters, “T” and “A” show introduction of complete mutation (C→T or G→A) into the clone, “t” shows that not less than 50% of mutation (C→T) is introduced into the clone (incomplete cloning), and “c” shows that the introduction efficiency of the mutation (C→T) into the clone is less than 50%.

[0053] FIG. 13 is a schematic illustration showing a temperature sensitive plasmid for mutation introduction, which was used in Example 11.

[0054] FIG. 14 shows the protocol of mutation introduction in Example 11.

[0055] FIG. 15 shows the results of introduction of mutation into the *rpoB* gene in Example 11.

[0056] FIG. 16 shows the results of introduction of mutation into the *galK* gene in Example 11.

DESCRIPTION OF EMBODIMENTS

[0057] The present invention provides a method of modifying a targeted site of a double stranded DNA by converting the target nucleotide sequence and nucleotides in the vicinity thereof in the double stranded DNA to other nucleotides, without cleaving at least one strand of the double stranded DNA to be modified. The method characteristically comprises a step of contacting a complex wherein a nucleic acid sequence-recognizing module that specifically binds to the target nucleotide sequence in the double stranded DNA and a nucleic acid base converting enzyme are linked, with the double stranded DNA to convert the targeted site, i.e., the target nucleotide sequence and nucleotides in the vicinity thereof, to other nucleotides.

[0058] In the present invention, the “modification” of a double stranded DNA means that a nucleotide (e.g., dC) on a DNA strand is converted to another nucleotide (e.g., dT, dA or dG), or deleted, or a nucleotide or a nucleotide sequence is inserted between certain nucleotides on the DNA strand. While the double stranded DNA to be modified is not particularly limited, it is preferably a genomic DNA. The “targeted site” of a double stranded DNA means the entire or partial “target nucleotide sequence”, which a nucleic acid sequence-recognizing module specifically recognizes and binds to, or the vicinity of the target nucleotide sequence (one or both of 5' upstream and 3' downstream), and the length thereof can be appropriately adjusted between 1 base and several hundred bases according to the object.

[0059] In the present invention, the “nucleic acid sequence-recognizing module” means a molecule or molecule complex having an ability to specifically recognize

and bind to a particular nucleotide sequence (i.e., target nucleotide sequence) on a DNA strand. Binding of the nucleic acid sequence-recognizing module to a target nucleotide sequence enables a nucleic acid base converting enzyme linked to the module to specifically act on a targeted site of a double stranded DNA.

[0060] In the present invention, the “nucleic acid base converting enzyme” means an enzyme capable of converting a target nucleotide to another nucleotide by catalyzing a reaction for converting a substituent on a purine or pyrimidine ring on a DNA base to another group or atom, without cleaving the DNA strand.

[0061] In the present invention, the “nucleic acid-modifying enzyme complex” means a molecular complex comprising a complex of the above-mentioned nucleic acid sequence-recognizing module linked with a nucleic acid base converting enzyme, wherein the complex has nucleic acid base converting enzyme activity and is imparted with a particular nucleotide sequence recognition ability. The “complex” used herein encompasses not only one composed of a plurality of molecules, but also a single molecule having a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme such as a fusion protein.

[0062] The nucleic acid base converting enzyme used in the present invention is not particularly limited as long as it can catalyze the above-mentioned reaction, and examples thereof include deaminase belonging to the nucleic acid/nucleotide deaminase superfamily, which catalyzes a deamination reaction that converts an amino group to a carbonyl group. Preferable examples thereof include cytidine deaminase capable of converting cytosine or 5-methylcytosine to uracil or thymine, respectively, adenosine deaminase capable of converting adenine to hypoxanthine, guanosine deaminase capable of converting guanine to xanthine and the like. As cytidine deaminase, more preferred is activation-induced cytidine deaminase (hereinafter also referred to as AID), which is an enzyme that introduces a mutation into an immunoglobulin gene in the acquired immunity of vertebrate or the like.

[0063] While the origin of nucleic acid base converting enzyme is not particularly limited, for example, PmCDA1 (*Petromyzon marinus* cytosine deaminase 1) from *Petromyzon marinus*, or AID (Activation-induced cytidine deaminase; AICDA) from mammal (e.g., human, swine, bovine, horse, monkey etc) can be used. The base sequence and amino acid sequence of CDS of PmCDA1 are shown in SEQ ID NOs: 1 and 2, respectively, and the base sequence and amino acid sequence of CDS of human AID are shown in SEQ ID NOs: 3 and 4, respectively.

[0064] A target nucleotide sequence in a double stranded DNA to be recognized by the nucleic acid sequence-recognizing module in the nucleic acid-modifying enzyme complex of the present invention is not particularly limited as long as the module specifically binds to any sequence in the double stranded DNA. The length of the target nucleotide sequence only needs to be sufficient for specific binding of the nucleic acid sequence-recognizing module. For example, when mutation is introduced into a particular site in the genomic DNA of a mammal, it is not less than 12 nucleotides, preferably not less than 15 nucleotides, more preferably not less than 17 nucleotides, according to the genome size thereof. While the upper limit of the length is not particularly limited, it is preferably not more than 25 nucleotides, more preferably not more than 22 nucleotides.

[0065] As the nucleic acid sequence-recognizing module in the nucleic acid-modifying enzyme complex of the present invention, CRISPR-Cas system wherein at least one DNA cleavage ability of Cas is inactivated (CRISPR-mutant Cas), zinc finger motif, TAL effector and PPR motif and the like, as well as a fragment containing a DNA binding domain of a protein that specifically binds to DNA such as restriction enzyme, transcription factor, RNA polymerase or the like, and not having a DNA double strand cleavage ability and the like can be used, but the module is not limited thereto. Preferably, the modules include CRISPR-mutant Cas, zinc finger motif, TAL effector, PPR motif and the like.

[0066] A zinc finger motif is constructed by linking 3-6 different Cys2His2 type zinc finger units (1 finger recognizes about 3 bases), and can recognize a target nucleotide sequence of 9-18 bases. A zinc finger motif can be produced by a known method such as Modular assembly method (*Nat Biotechnol* (2002) 20: 135-141), OPEN method (*Mol Cell* (2008) 31: 294-301), CoDA method (*Nat Methods* (2011) 8: 67-69), *Escherichia coli* one-hybrid method (*Nat Biotechnol* (2008) 26:695-701) and the like. The above-mentioned patent document 1 can be referred to as for the detail of the zinc finger motif production.

[0067] A TAL effector has a module repeat structure with about 34 amino acids as a unit, and the 12th and 13th amino acid residues (called RVD) of one module determine the binding stability and base specificity. Since each module is highly independent, TAL effector specific to a target nucleotide sequence can be produced by simply linking the modules. For TAL effector, production methods utilizing an open resource (REAL method (*Curr Protoc Mol Biol* (2012) Chapter 12: Unit 12.15), FLASH method (*Nat Biotechnol* (2012) 30: 460-465), and Golden Gate method (*Nucleic Acids Res* (2011) 39: e82) etc) have been established, and a TAL effector for a target nucleotide sequence can be designed relatively easily. The above-mentioned patent document 2 can be referred to as for the detail of the production of TAL effector.

[0068] PPR motif is constructed such that a particular nucleotide sequence is recognized by a series of PPR motifs each consisting of 35 amino acids and recognizing one nucleic acid base, and recognizes a target base only by 1, 4 and ii(-2) amino acids of each motif. Motif configuration has no dependency, and is free of interference of motifs on both sides. Therefore, similar to TAL effector, a PPR protein specific to the target nucleotide sequence can be produced by simply linking PPR motifs. The above-mentioned patent document 4 can be referred to as for the detail of the production of PPR motif.

[0069] When a fragment of a restriction enzyme, transcription factor, RNA polymerase or the like is used, since the DNA binding domains of these proteins are well known, a fragment containing said domain and not having a DNA double strand cleavage ability can be easily designed and constructed.

[0070] Any of the above-mentioned nucleic acid sequence-recognizing module can be provided as a fusion protein with the above-mentioned nucleic acid base converting enzyme, or a protein binding domain such as SH3 domain, PDZ domain, GK domain, GB domain and the like and a binding partner thereof may be fused with a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme, respectively, and provided as a protein complex via an interaction of the domain and a binding

partner thereof. Alternatively, a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme may be each fused with intein, and they can be linked by ligation after protein synthesis.

[0071] The nucleic acid-modifying enzyme complex of the present invention containing a complex (including fusion protein), wherein a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme are linked, may be contacted with a double stranded DNA as an enzyme reaction in a cell-free system. In view of the main object of the present invention, it is desirable to perform the contact by introducing a nucleic acid encoding the complex into a cell having the double stranded DNA of interest (e.g., genomic DNA).

[0072] Therefore, the nucleic acid sequence-recognizing module and the nucleic acid base converting enzyme are preferably prepared as a nucleic acid encoding a fusion protein thereof, or as nucleic acids encoding each of them in a form capable of forming a complex in a host cell after translation into a protein by utilizing a binding domain, intein or the like. The nucleic acid here may be a DNA or an RNA. When it is a DNA, it is preferably a double stranded DNA, and provided in the form of an expression vector disposed under regulation of a functional promoter in a host cell. When it is an RNA, it is preferably a single stranded RNA.

[0073] Since the complex of the present invention wherein a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme are linked, is not associated with double-stranded DNA breaks (DSB), genome editing with low toxicity is possible, and the genetic modification method of the present invention can be applied to a wide range of biological materials. Therefore, the cells into which nucleic acid encoding nucleic acid sequence-recognizing module and/or nucleic acid base converting enzyme is introduced can encompass cells of any species, from cells of microorganisms, such as bacterium, such as *Escherichia coli* and the like which are prokaryotes, such as yeast and the like which are lower eukaryotes, to cells of higher eukaryotes such as insect, plant and the like, and cells of vertebrates including mammals such as human and the like.

[0074] A DNA encoding a nucleic acid sequence-recognizing module such as zinc finger motif, TAL effector, PPR motif and the like can be obtained by any method mentioned above for each module. A DNA encoding a sequence-recognizing module of restriction enzyme, transcription factor, RNA polymerase and the like can be cloned by, for example, synthesizing an oligoDNA primer covering a region encoding a desired part of the protein (part containing DNA binding domain) based on the cDNA sequence information thereof, and amplifying by the RT-PCR method using, the total RNA or mRNA fraction prepared from the protein-producing cells as a template.

[0075] A DNA encoding a nucleic acid base converting enzyme can also be cloned similarly by synthesizing an oligoDNA primer based on the cDNA sequence information thereof, and amplifying by the RT-PCR method using, the total RNA or mRNA fraction prepared from the enzyme-producing cells as a template. For example, a DNA encoding PmCDA1 of *Petromyzon marinus* can be cloned by designing suitable primers for the upstream and downstream of CDS based on the cDNA sequence (accession No. EF094822) registered in the NCBI database, and cloning from mRNA *Petromyzon marinus* by the RT-PCR method. A

DNA encoding human AID can be cloned by designing suitable primers for the upstream and downstream of CDS based on the cDNA sequence (accession No. AB040431) registered in the NCBI database, and cloning from, for example, mRNA from human lymph node by the RT-PCR method.

[0076] The cloned DNA may be directly, or after digestion with a restriction enzyme when desired, or after addition of a suitable linker and/or a nuclear localization signal (each organelle transfer signal when the target double stranded DNA of interest is mitochondria or chloroplast DNA), ligated with a DNA encoding a nucleic acid sequence-recognizing module to prepare a DNA encoding a fusion protein. Alternatively, a DNA encoding a nucleic acid sequence-recognizing module, and a DNA encoding a nucleic acid base converting enzyme may be each fused with a DNA encoding a binding domain or a binding partner thereof, or both DNAs may be fused with a DNA encoding a separation intein, whereby the nucleic acid sequence-recognizing conversion module and the nucleic acid base converting enzyme are translated in a host cell to form a complex. In these cases, a linker and/or a nuclear localization signal can be linked to a suitable position of one of or both DNAs when desired.

[0077] A DNA encoding a nucleic acid sequence-recognizing module and a DNA encoding a nucleic acid base converting enzyme can be obtained by chemically synthesizing the DNA strand, or by linking partly overlapping synthesized oligoDNA short strands by utilizing the PCR method and the Gibson Assembly method to construct a DNA encoding the full length thereof. The advantage of constructing a full-length DNA by chemical synthesis or a combination of PCR method or Gibson Assembly method is that the codon used can be designed in CDS full-length according to the host into which the DNA is introduced. In the expression of a heterologous DNA, the protein expression level is expected to increase by converting the DNA sequence thereof to a codon which is highly frequently used in the host organism. As the data of codon use frequency in host used, for example, the genetic code use frequency database (www.kazusa.or.jp/codon/index.html) disclosed in the home page of Kazusa DNA Research Institute can be used, or documents showing the codon use frequency in each host may be referred to. By reference to the obtained data and the DNA sequence to be introduced, codons showing low use frequency in the host from those used for the DNA sequence may be converted to a codon coding the same amino acid and showing high use frequency.

[0078] An expression vector containing a DNA encoding a nucleic acid sequence-recognizing module and/or a nucleic acid base converting enzyme can be produced, for example, by linking the DNA to the downstream of a promoter in a suitable expression vector.

[0079] As the expression vector, plasmids from *Escherichia coli* (e.g., pBR322, pBR325, pUC12, pUC13); plasmids from *Bacillus subtilis* (e.g., pUB110, pTP5, pC194); plasmids from yeast (e.g., pSH19, pSH15); insect cell expression plasmids (e.g., pFast-Bac); animal cell expression plasmids (e.g., pA1-11, pXT1, pRc/CMV, pRc/RSV, pCDNAI/Neo); bacteriophages such as λ phage and the like; insect virus vectors such as baculovirus and the like (e.g., BmNPV, AcNPV); animal virus vectors such as retrovirus, vaccinia virus, adenovirus and the like, are used.

[0080] As the promoter, any promoter appropriate for a host used for gene expression can be used. In a conventional method involving DSB, since the survival rate of the host cell sometimes decreases markedly due to the toxicity, it is desirable to increase the number of cells by the start of the induction by using an inductive promoter. However, since sufficient cell proliferation can also be achieved by expressing the nucleic acid-modifying enzyme complex of the present invention, a constitutive promoter can also be used without limitation.

[0081] For example, when the host is an animal cell, SRA promoter, SV40 promoter, LTR promoter, CMV (cytomegalovirus) promoter, RSV (Rous sarcoma virus) promoter, MoMuLV (Moloney mouse leukemia virus) LTR, HSV-TK (simple herpes virus thymidine kinase) promoter and the like are used. Of these, CMV promoter, SRA promoter and the like are preferable.

[0082] When the host is *Escherichia coli*, trp promoter, lac promoter, recA promoter, λP_L promoter, Ipp promoter, T7 promoter and the like are preferable.

[0083] When the host is genus *Bacillus*, SPO1 promoter, SPO2 promoter, penP promoter and the like are preferable.

[0084] When the host is a yeast, Gal1/10 promoter, PHO5 promoter, PGK promoter, GAP promoter, ADH promoter and the like are preferable.

[0085] When the host is an insect cell, polyhedrin promoter, P10 promoter and the like are preferable.

[0086] When the host is a plant cell, CaMV35S promoter, CaMV19S promoter, NOS promoter and the like are preferable.

[0087] As the expression vector, besides those mentioned above, one containing enhancer, splicing signal, terminator, polyA addition signal, a selection marker such as drug resistance gene, auxotrophic complementary gene and the like, replication origin and the like on demand can be used.

[0088] An RNA encoding a nucleic acid sequence-recognizing module and/or a nucleic acid base converting enzyme can be prepared by, for example, transcription to mRNA in an in vitro transcription system known per se by using a vector encoding DNA encoding the above-mentioned nucleic acid sequence-recognizing module and/or a nucleic acid base converting enzyme as a template.

[0089] A complex of a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme can be intracellularly expressed by introducing an expression vector containing a DNA encoding a nucleic acid sequence-recognizing module and/or a nucleic acid base converting enzyme into a host cell, and culturing the host cell.

[0090] As the host, genus *Escherichia*, genus *Bacillus*, yeast, insect cell, insect, animal cell and the like are used.

[0091] As the genus *Escherichia*, *Escherichia coli* K12-DH1 [*Proc. Natl. Acad. Sci. USA*, 60, 160 (1968)], *Escherichia coli* JM103 [*Nucleic Acids Research*, 9, 309 (1981)], *Escherichia coli* JA221 [*Journal of Molecular Biology*, 120, 517 (1978)], *Escherichia coli* HB101 [*Journal of Molecular Biology*, 41, 459 (1969)], *Escherichia coli* C600 [*Genetics*, 39, 440 (1954)] and the like are used.

[0092] As the genus *Bacillus*, *Bacillus subtilis* M1114 [*Gene*, 24, 255 (1983)], *Bacillus subtilis* 207-21 [*Journal of Biochemistry*, 95, 87 (1984)] and the like are used.

[0093] As the yeast, *Saccharomyces cerevisiae* AH22, AH22R, NA87-11A, DKD-5D, 20B-12, *Schizosaccharomyces pombe* NCYC1913, NCYC2036, *Pichia pastoris* KM71 and the like are used.

[0094] As the insect cell when the virus is AcNPV, cells of established line from cabbage armyworm larva (*Spodoptera frugiperda* cell; Sf cell), MG1 cells from the mid-intestine of *Trichoplusia ni*, High Five™ cells from an egg of *Trichoplusia ni*, cells from *Mamestra brassicae*, cells from *Estigmene acrea* and the like are used. When the virus is BmNPV, cells of established line from *Bombyx mori* (*Bombyx mori* N cell; BmN cell) and the like are used as insect cells. As the Sf cell, for example, Sf9 cell (ATCC CRL1711), Sf21 cell [all above, *In Vivo*, 13, 213-217 (1977)] and the like are used.

[0095] As the insect, for example, larva of *Bombyx mori*, *Drosophila*, cricket and the like are used [*Nature*, 315, 592 (1985)].

[0096] As the animal cell, cell lines such as monkey COS-7 cell, monkey Vero cell, Chinese hamster ovary (CHO) cell, dhfr gene-deficient CHO cell, mouse L cell, mouse AtT-20 cell, mouse myeloma cell, rat GH3 cell, human FL cell and the like, pluripotent stem cells such as iPS cell, ES cell and the like of human and other mammals, and primary cultured cells prepared from various tissues are used. Furthermore, zebrafish embryo, *Xenopus* oocyte and the like can also be used.

[0097] As the plant cell, suspend cultured cells, callus, protoplast, leaf segment, root segment and the like prepared from various plants (e.g., grain such as rice, wheat, corn and the like, product crops such as tomato, cucumber, egg plant and the like, garden plants such as carnation, *Eustoma russellianum* and the like, experiment plants such as tobacco, *Arabidopsis thaliana* and the like) are used.

[0098] All the above-mentioned host cells may be haploid (monoploid), or polyploid (e.g., diploid, triploid, tetraploid and the like). In the conventional mutation introduction methods, mutation is, in principle, introduced into only one homologous chromosome to produce a heterologous genotype. Therefore, the desired feature is not expressed unless it is a dominant mutation, and making it homologous inconveniently requires labor and time. In contrast, according to the present invention, since mutations can be introduced into all alleles on the homologous chromosome in the genome, desired feature can be expressed in a single generation even in the case of recessive mutation (FIG. 10), which is extremely useful since the problem of the conventional method can be solved.

[0099] An expression vector can be introduced by a known method (e.g., lysozyme method, competent method, PEG method, CaCl₂ coprecipitation method, electroporation method, the microinjection method, the particle gun method, lipofection method, *Agrobacterium* method and the like) according to the kind of the host.

[0100] *Escherichia coli* can be transformed according to the methods described in, for example, *Proc. Natl. Acad. Sci. USA*, 69, 2110 (1972), *Gene*, 17, 107 (1982) and the like.

[0101] A vector can be introduced into the genus *Bacillus* according to the methods described in, for example, *Molecular & General Genetics*, 168, 111 (1979) and the like.

[0102] A vector can be introduced into a yeast according to the methods described in, for example, *Methods in Enzymology*, 194, 182-187 (1991), *Proc. Natl. Acad. Sci. USA*, 75, 1929 (1978) and the like.

[0103] A vector can be introduced into an insect cell and an insect according to the methods described in, for example, *Bio/Technology*, 6, 47-55 (1988) and the like.

[0104] A vector can be introduced into an animal cell according to the methods described in, for example, *Cell Engineering* additional volume 8, New Cell Engineering Experiment Protocol, 263-267 (1995) (published by Shujunsha), and *Virology*, 52, 456 (1973).

[0105] A cell introduced with a vector can be cultured according to a known method according to the kind of the host.

[0106] For example, when *Escherichia coli* or genus *Bacillus* is cultured, a liquid medium is preferable as a medium used for the culture. The medium preferably contains a carbon source, nitrogen source, inorganic substance and the like necessary for the growth of the transformant. Examples of the carbon source include glucose, dextrin, soluble starch, sucrose and the like; examples of the nitrogen source include inorganic or organic substances such as ammonium salts, nitrate salts, corn steep liquor, peptone, casein, meat extract, soybean cake, potato extract and the like; and examples of the inorganic substance include calcium chloride, sodium dihydrogen phosphate, magnesium chloride and the like. The medium may contain yeast extract, vitamins, growth promoting factor and the like. The pH of the medium is preferably about 5-about 8.

[0107] As a medium for culturing *Escherichia coli*, for example, M9 medium containing glucose, casamino acid [*Journal of Experiments in Molecular Genetics*, 431-433, Cold Spring Harbor Laboratory, New York 1972] is preferable. Where necessary, for example, agents such as 3 β -indolylacrylic acid may be added to the medium to ensure an efficient function of a promoter. *Escherichia coli* is cultured at generally about 15-about 43° C. Where necessary, aeration and stirring may be performed.

[0108] The genus *Bacillus* is cultured at generally about 30-about 40° C. Where necessary, aeration and stirring may be performed.

[0109] Examples of the medium for culturing yeast include Burkholder minimum medium [*Proc. Natl. Acad. Sci. USA*, 77, 4505 (1980)], SD medium containing 0.5% casamino acid [*Proc. Natl. Acad. Sci. USA*, 81, 5330 (1984)] and the like. The pH of the medium is preferably about 5-about 8. The culture is performed at generally about 20° C.-about 35° C. Where necessary, aeration and stirring may be performed.

[0110] As a medium for culturing an insect cell or insect, for example, Grace's Insect Medium [*Nature*, 195, 788 (1962)] containing an additive such as inactivated 10% bovine serum and the like as appropriate and the like are used. The pH of the medium is preferably about 6.2-about 6.4. The culture is performed at generally about 27° C. Where necessary, aeration and stirring may be performed.

[0111] As a medium for culturing an animal cell, for example, minimum essential medium (MEM) containing about 5-about 20% of fetal bovine serum [*Science*, 122, 501 (1952)], Dulbecco's modified Eagle medium (DMEM) [*Virology*, 8, 396 (1959)], RPMI 1640 medium [*The Journal of the American Medical Association*, 199, 519 (1967)], 199 medium [*Proceeding of the Society for the Biological Medicine*, 73, 1 (1950)] and the like are used. The pH of the medium is preferably about 6-about 8. The culture is performed at generally about 30° C.-about 40° C. Where necessary, aeration and stirring may be performed.

[0112] As a medium for culturing a plant cell, for example, MS medium, LS medium, B5 medium and the like are used. The pH of the medium is preferably about 5-about 8. The

culture is performed at generally about 20° C.-about 30° C. Where necessary, aeration and stirring may be performed.

[0113] As mentioned above, a complex of a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme, i.e., nucleic acid-modifying enzyme complex, can be expressed intracellularly.

[0114] An RNA encoding a nucleic acid sequence-recognizing module and/or a nucleic acid base converting enzyme can be introduced into a host cell by microinjection method, lipofection method and the like. RNA introduction can be performed once or multiple times (e.g., 2-5 times) at suitable intervals.

[0115] When a complex of a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme is expressed by an expression vector or RNA molecule introduced into the cell, the nucleic acid sequence-recognizing module specifically recognizes and binds to a target nucleotide sequence in the double stranded DNA (e.g., genomic DNA) of interest and, due to the action of the nucleic acid base converting enzyme linked to the nucleic acid sequence-recognizing module, base conversion occurs in the sense strand or antisense strand of the targeted site (whole or partial target nucleotide sequence or appropriately adjusted within several hundred bases including the vicinity thereof) and a mismatch occurs in the double stranded DNA (e.g., when cytidine deaminase such as PmCDA1, AID and the like is used as a nucleic acid base converting enzyme, cytosine on the sense strand or antisense strand at the targeted site is converted to uracil to cause U:G or G:U mismatch). When the mismatch is not correctly repaired, and when repaired such that a base of the opposite strand forms a pair with a base of the converted strand (T-A or A-T in the above-mentioned example), or when another nucleotide is further substituted (e.g., U→A, G) or when one to several dozen bases are deleted or inserted during repair, various mutations are introduced.

[0116] As for zinc finger motif, production of many actually functional zinc finger motifs is not easy, since production efficiency of a zinc finger that specifically binds to a target nucleotide sequence is not high and selection of a zinc finger having high binding specificity is not easy. While TAL effector and PPR motif have a high degree of freedom of target nucleic acid sequence recognition as compared to zinc finger motif, a problem remains in the efficiency since a large protein needs to be designed and constructed every time according to the target nucleotide sequence.

[0117] In contrast, since the CRISPR-Cas system recognizes the sequence of double stranded DNA of interest by a guide RNA complementary to the target nucleotide sequence, any sequence can be targeted by simply synthesizing an oligoDNA capable of specifically forming a hybrid with the target nucleotide sequence.

[0118] Therefore, in a more preferable embodiment of the present invention, a CRISPR-Cas system wherein at least one DNA cleavage ability of Cas is inactivated (CRISPR-mutant Cas) is used as a nucleic acid sequence-recognizing module.

[0119] FIG. 1 is a schematic illustration showing the double stranded DNA modification method of the present invention using CRISPR-mutant Cas as a nucleic acid sequence-recognizing module.

[0120] The nucleic acid sequence-recognizing module of the present invention using CRISPR-mutant Cas is provided as a complex of an RNA molecule consisting of a guide

RNA complementary to the target nucleotide sequence and tracrRNA necessary for recruiting mutant Cas protein, and a mutant Cas protein.

[0121] The Cas protein used in the present invention is not particularly limited as long as it belongs to the CRISPR system, and is preferably Cas9. Examples of Cas9 include, but are not limited to, Cas9 (SpCas9 from *Streptococcus pyogenes*, Cas9 (StCas9) from *Streptococcus thermophilus* and the like, preferably SpCas9. As a mutant Cas used in the present invention, either a Cas having cleavage ability of both strands of the double stranded DNA is inactivated, or a Cas having nickase activity wherein only one of the cleavage ability of only one of the strands is inactivated, can be used. For example, in the case of SpCas9, a D1 OA mutant wherein the 10th Asp residue is converted to an Ala residue and lacking cleavage ability of a strand opposite to the strand forming a complementary strand with a guide RNA, or H840A mutant wherein the 840th His residue is converted to an Ala residue and lacking cleavage ability of strand complementary to guide RNA, or a double mutant thereof can be used, and another mutant Cas can be used similarly.

[0122] A nucleic acid base converting enzyme is provided as a complex with mutant Cas by a method similar to the linking scheme with the above-mentioned zinc finger and the like. Alternatively, a nucleic acid base converting enzyme and mutant Cas can also be linked by utilizing RNA scaffold with RNA aptamers MS2F6, PP7 and the like and binding proteins thereto. Guide RNA forms a complementary strand with the target nucleotide sequence, mutant Cas is recruited by the attached tracrRNA and mutant Cas recognizes DNA cleavage site recognition sequence PAM (protospacer adjacent motif) (when SpCas9 is used, PAM is 3 bases of NGG (N is any base), and, theoretically, can target any position on the genome). One or both DNAs cannot be cleaved, and, due to the action of the nucleic acid base converting enzyme linked to the mutant Cas, base conversion occurs in the targeted site (appropriately adjusted within several hundred bases including whole or partial target nucleotide sequence) and a mismatch occurs in the double stranded DNA. When the mismatch is not correctly repaired, and when repaired such that a base of the opposite strand forms a pair with a base of the converted strand, or when another nucleotide is further converted or when one to several dozen bases are deleted or inserted during repair, various mutations are introduced (see, e.g., FIG. 2).

[0123] Even when CRISPR-mutant Cas is used as a nucleic acid sequence-recognizing module, a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme are introduced, desirably in the form of a nucleic acid encoding same, into a cell having a double stranded DNA of interest, similar to when zinc finger and the like are used as a nucleic acid sequence-recognizing module.

[0124] A DNA encoding Cas can be cloned by a method similar to the above-mentioned method for a DNA encoding a nucleic acid base converting enzyme, from a cell producing the enzyme. A mutant Cas can be obtained by introducing a mutation to convert an amino acid residue of the part important for the DNA cleavage activity (e.g., 10th Asp residue and 840th His residue for Cas9, though not limited thereto) to another amino acid, into a DNA encoding cloned Cas, by a site specific mutation induction method known per se.

[0125] Alternatively, a DNA encoding mutant Cas can also be constructed as a DNA having codon usage suitable

for expression in a host cell to be used, by a method similar to those mentioned above for a DNA encoding a nucleic acid sequence-recognizing module and a DNA encoding a nucleic acid base converting enzyme, and in a combination with chemical synthesis or PCR method or Gibson Assembly method. For example, CDS sequence and amino acid sequence optimized for the expression of SpCas9 in eukaryotic cells are shown in SEQ ID NOs: 5 and 6. In the sequence shown in SEQ ID NO: 5, when "A" is converted to "C" in base No. 29, a DNA encoding a D1 OA mutant can be obtained, and when "CA" is converted to "GC" in base Nos. 2518-2519, a DNA encoding an H840A mutant can be obtained.

[0126] A DNA encoding a mutant Cas and a DNA encoding a nucleic acid base converting enzyme may be linked to allow for expression as a fusion protein, or designed to be separately expressed using a binding domain, intein or the like, and form a complex in a host cell via protein-protein interaction or protein ligation.

[0127] The obtained DNA encoding a mutant Cas and/or a nucleic acid base converting enzyme can be inserted into the downstream of a promoter of an expression vector similar to the one mentioned above, according to the host.

[0128] On the other hand, a DNA encoding guide RNA and tracrRNA can be obtained by designing an oligoDNA sequence linking guide RNA sequence complementary to the target nucleotide sequence and known tracrRNA sequence (e.g., gtttagagctagaatagcaagttaaaataaggctagtc-cggtatcaactgaaaaagtggcaccgagtcggtggtgctttt; SEQ ID NO: 7) and chemically synthesizing using a DNA/RNA synthesizer.

[0129] While the length of the guide RNA sequence is not particularly limited as long as it can specifically bind to a target nucleotide sequence, for example, it is 15-30 nucleotides, preferably 18-24 nucleotides.

[0130] While a DNA encoding guide RNA and tracrRNA can also be inserted into an expression vector similar to the one mentioned above, according to the host. As the promoter, pol III promoter (e.g., SNR6, SNR52, SCR1, RPR1, U6, H1 promoter etc.) and terminator (e.g., T_e sequence) are preferably used.

[0131] An RNA encoding mutant Cas and/or a nucleic acid base converting enzyme can be prepared by, for example, transcription to mRNA in an in vitro transcription system known per se by using a vector encoding the above-mentioned mutant Cas and/or DNA encoding a nucleic acid base converting enzyme as a template.

[0132] Guide RNA-tracrRNA can be obtained by designing an oligoDNA sequence in which a sequence complementary to the target nucleotide sequence and known tracrRNA sequence are linked, and chemically synthesizing using a DNA/RNA synthesizer.

[0133] A DNA or RNA encoding mutant Cas and/or a nucleic acid base converting enzyme, guide RNA-tracrRNA or a DNA encoding same can be introduced into a host cell by a method similar to the above, according to the host.

[0134] Since conventional artificial nuclease accompanies Double-stranded DNA breaks (DSB), inhibition of growth and cell death assumedly caused by disordered cleavage (off-target cleavage) of chromosome may occur by targeting a sequence in the genome. The effect thereof is particularly fatal for many microorganisms and prokaryotes, and prevents applicability. In the present invention, mutation is introduced not by DNA cleavage but by a conversion

reaction of the substituent on the DNA base (particularly deamination reaction), and therefore, drastic reduction of toxicity can be realized. In fact, as shown in the comparison tests using a budding yeast as a host in the below-mentioned Examples, when Cas9 having a conventional type of DSB activity is used, the number of surviving cells decreases by induction of expression, whereas it was confirmed that the cells continued to grow and the number of surviving cells increased by the technique of the present invention using a combination of mutant Cas and a nucleic acid base converting enzyme in combination (FIG. 3).

[0135] The modification of the double stranded DNA in the present invention does not preclude occurrence of cleavage of the double stranded DNA in a site other than the targeted site (appropriately adjusted within several hundred bases including whole or partial target nucleotide sequence). However, one of the greatest advantages of the present invention is avoidance of toxicity by off-target cleavage, which is generally applicable to any species. In one preferable embodiment, therefore, the modification of the double stranded DNA in the present invention is not associated with cleavage of DNA strand not only in a targeted site of a selected double stranded DNA but in other sites.

[0136] As shown in the below-mentioned Examples, when Cas having a nickase activity capable of cleaving only one of the strands of the double stranded DNA is used as a mutant Cas (FIG. 5), the mutation introduction efficiency increases as compared to when mutant Cas which is incapable of cleaving both strands is used. Therefore, for example, besides a nucleic acid sequence-recognizing module and a nucleic acid base converting enzyme, linking a protein having a nickase activity, thereby cleaving only a DNA single strand in the vicinity of the target nucleotide sequence, the mutation introduction efficiency can be improved while avoiding the strong toxicity of DSB.

[0137] Furthermore, a comparison of the effects of mutant Cas having two kinds of nickase activity of cleaving different strand reveals that using one of the mutant Cas results in mutated sites accumulating near the center of the target nucleotide sequence, and using another mutant Cas results in various mutations which are randomly introduced into region of several hundred bases from the target nucleotide sequence (FIG. 6). Therefore, by selecting a strand to be cleaved by the nickase, a mutation can be introduced into a particular nucleotide or nucleotide region at a pinpoint, or various mutations can be randomly introduced into a comparatively wide range, which can be properly adopted according to the object. For example, when the former technique is applied to genetically diseased iPS cell, a cell transplantation therapeutic agent with a lower risk of rejection can be produced by repairing mutation of the pathogenic gene in an iPS cell produced from the patients' own cell, and differentiating the cell into the somatic cell of interest.

[0138] Example 7 and the subsequent Examples mentioned below show that a mutation can be introduced into a particular nucleotide almost at a pinpoint. For pinpoint introduction of a mutation into a desired nucleotide, the target nucleotide sequence should be set to show certain regularity of the positional relationship between a nucleotide desired to be introduced with a mutation and the target nucleotide sequence. CRISPR-Cas system is used as a nucleic acid sequence-recognizing module and AID is used as a nucleic acid base converting enzyme, the target nucleotide

sequence can be designed such that C (or G in the opposite strand) into which a mutation is desired to be introduced is at 2-5 nucleotides from the 5'-end of the target nucleotide sequence. As mentioned above, the length of the guide RNA sequence can be appropriately determined to fall between 15-30 nucleotides, preferably 18-24 nucleotides. Since the guide RNA sequence is a sequence complementary to the target nucleotide sequence, the length of the target nucleotide sequence changes when the length of the guide RNA sequence is changed; however, the regularity that a mutation is likely to be introduced into C or G at 2-5 nucleotides from the 5'-end irrespective of the length of the nucleotide, is maintained (FIG. 12). Therefore, by appropriately determining the length of the target nucleotide sequence (guide RNA as a complementary strand thereof), the site of a base into which a mutation can be introduced can be shifted. As a result, restriction by DNA cleavage site recognition sequence PAM (NGG) can also be removed, and the degree of freedom of mutation introduction becomes higher.

[0139] As shown in the below-mentioned Examples, when sequence-recognizing modules are produced corresponding to a plurality of target nucleotide sequences in proximity, and simultaneously used, the mutation introduction efficiency drastically increases relative to when a single nucleotide sequence is used as a target (FIG. 7). As the effect thereof, similar mutation induction is realized even when both target nucleotide sequences partly overlap or when the both are apart by about 600 bp. It can occur when both target nucleotide sequences are in the same direction (target nucleotide sequences are present on the same strand) (FIG. 7), and when they are opposed (target nucleotide sequences are present on each strand of double stranded DNA) (FIG. 4).

[0140] As shown in the below-mentioned Examples, the genome sequence modification method of the present invention can introduce mutation into almost all cells in which the nucleic acid-modifying enzyme complex of the present invention has been expressed, by selecting a suitable target nucleotide sequence (FIG. 8). Thus, insertion and selection of a selection marker gene, which are essential in the conventional genome editing, are not necessary. This dramatically facilitates and simplifies gene manipulation and extends the applicability to crop breeding and the like since a recombinant organism with foreign DNA is not produced.

[0141] Since the genome sequence modification method of the present invention shows extremely high mutation introduction efficiency, and does not require selection by markers, a plurality of DNA regions at completely different positions can be modified as targets (FIG. 9). Therefore, in one preferable embodiment of the present invention, two or more kinds of nucleic acid sequence-recognizing modules that specifically bind to different target nucleotide sequences (which may be present in one target gene of interest, or two or more different target genes of interest, which may be present on the same chromosome or different chromosomes) can be used. In this case, each one of these nucleic acid sequence-recognizing modules and nucleic acid base converting enzyme form a nucleic acid-modifying enzyme complex. Here, a common nucleic acid base converting enzyme can be used. For example, when CRISPR-Cas system is used as a nucleic acid sequence-recognizing module, a common complex of a Cas protein and a nucleic acid base converting enzyme (including fusion protein) is

used, and two or more kinds of chimeric RNAs of tracrRNA and each of two or more guide RNAs that respectively form a complementary strand with a different target nucleotide sequences are produced and used as guide RNA-tracrRNAs. On the other hand, when zinc finger motif, TAL effector and the like are used as nucleic acid sequence-recognizing modules, for example, a nucleic acid base converting enzyme can be fused with a nucleic acid sequence-recognizing module that specifically binds to a different target nucleotide.

[0142] To express the nucleic acid-modifying enzyme complex of the present invention in a host cell, as mentioned above, an expression vector containing a DNA encoding the nucleic acid-modifying enzyme complex, or an RNA encoding the nucleic acid-modifying enzyme complex is introduced into a host cell. For efficient introduction of mutation, it is desirable to maintain an expression of nucleic acid-modifying enzyme complex at a given level or above for not less than a given period. From such aspect, introduction of an expression vector autonomously replicatable in a host cell (plasmid etc.) is reliable. However, since the plasmid etc. are foreign DNAs, they are preferably removed rapidly after successful introduction of mutation. Therefore, although it varies depending on the kind of host cell and the like, for example, the introduced plasmid is desirably removed from the host cell after a lapse of 6 hr-2 days from the introduction of an expression vector by using various plasmid removal methods which are well known in the art.

[0143] Alternatively, as long as sufficient expression of a nucleic acid-modifying enzyme complex for the introduction of mutation is achieved, it is also preferable to introduce mutation into the target double stranded DNA of interest by transient expression by using an expression vector without autonomous replicatability in a host cell (e.g., vector lacking replication origin that functions in a host cell and/or gene encoding protein necessary for replication etc.) or RNA.

[0144] Expression of target gene is suppressed while the nucleic acid-modifying enzyme complex of the present invention is expressed in a host cell to perform a nucleic acid base conversion reaction. Therefore, it was difficult to directly edit a gene essential for the survival of the host cell as a target gene (result in side effects such as growth inhibition of host, unstable mutation introduction efficiency, mutation of site different from target and the like). In the present invention, direct editing of an essential gene has been successfully and efficiently realized by causing a nucleic acid base conversion reaction at a desired stage, and transiently expressing the nucleic acid-modifying enzyme complex of the present invention in a host cell for a period necessary for stabilizing the modification of the targeted site. While the period necessary for a nucleic acid base conversion reaction and stabilizing the modification of the targeted site varies depending on the kind of the host cell, culture conditions and the like, host cells of 2-20 generations are generally considered to be necessary. For example, when the host cell is a yeast or bacterium (e.g., *Escherichia coli*), expression of a nucleic acid-modifying enzyme complex needs to be induced for 5-10 generations. Those of ordinary skill in the art can appropriately determine a preferable expression induction period based on the doubling time of the host cell under culture conditions used. For example, when a budding yeast is subjected to liquid culture in a 0.02% galactose inducer medium, the expression induction period is, for example, 20-40 hr. The expression induction

period of the nucleic acid encoding the nucleic acid-modifying enzyme complex of the present invention may be extended beyond the above-mentioned "period necessary for establishing the modification of the targeted site" to the extent not causing side effects to the host cell.

[0145] As a means for transiently expressing the nucleic acid-modifying enzyme complex of the present invention at a desired stage for a desired period, a method comprising producing a construct (expression vector) containing a nucleic acid encoding the nucleic acid-modifying enzyme complex (a DNA encoding a guide RNA-tracrRNA and a DNA encoding a mutant Cas and nucleic acid base substitution enzyme in the case of CRISPR-Cas system), in a manner that the expression period can be controlled, and introducing the construct into a host cell can be used. The "manner that the expression period can be controlled" is specifically, for example, a nucleic acid encoding the nucleic acid-modifying enzyme complex of the present invention placed under regulation of an inducible regulatory region. While the "inducible regulatory region" is not particularly limited, it is, for example, an operon of a temperature sensitive (ts) mutation repressor and an operator regulated thereby in microorganism cells of bacterium (e.g., *Escherichia coli*), yeast and the like. Examples of the ts mutation repressor include, but are not limited to, ts mutation of *cl* repressor from λ phage. In the case of λ phage *cl* repressor (ts), it is linked to an operator to suppress expression of gene in the downstream at not more than 30° C. (e.g., 28° C.). At a high temperature of not less than 37° C. (e.g., 42° C.), it is dissociated from the operator to allow for induction of gene expression (FIGS. 13 and 14). Therefore, the period when the expression of the target gene is suppressed can be minimized by culturing a host cell introduced with a nucleic acid encoding nucleic acid-modifying enzyme complex generally at not more than 30° C., raising the temperature to not less than 37° C. at an appropriate stage, performing culture for a given period to carry out a nucleic acid base conversion reaction and, after introduction of mutation into the target gene, rapidly lowering the temperature to not more than 30° C. Thus, even when an essential gene for the host cell is targeted, it can be efficiently edited while suppressing the side effects (FIG. 15).

[0146] When temperature sensitive mutation is utilized, for example, a temperature sensitive mutant of a protein necessary for autonomous replication of a vector is included in a vector containing a DNA encoding the nucleic acid-modifying enzyme complex of the present invention. As a result, autonomous replication becomes impossible rapidly after expression of the nucleic acid-modifying enzyme complex, and the vector naturally falls off during the cell division. Examples of the temperature sensitive mutant protein include, but are not limited to, a temperature sensitive mutant of Rep101 ori necessary for the replication of pSC101 ori. Rep101 ori (ts) acts on pSC101 ori to enable autonomous replication of plasmid at not more than 30° C. (e.g., 28° C.), but loses function at not less than 37° C. (e.g., 42° C.), and plasmid cannot replicate autonomously. Therefore, a combined use with *cl* repressor (ts) of the above-mentioned A phage simultaneously enables transient expression of the nucleic acid-modifying enzyme complex of the present invention, and removal of the plasmid.

[0147] On the other hand, when a higher eukaryotic cell such as animal cell, insect cell, plant cell and the like is used as a host cell, a DNA encoding the nucleic acid-modifying

enzyme complex of the present invention is introduced into a host cell under regulation of an inducible promoter (e.g., metallothionein promoter (induced by heavy metal ion), heat shock protein promoter (induced by heat shock), Tet-ON/Tet-OFF system promoter (induced by addition or removal of tetracycline or a derivative thereof), steroid-responsive promoter (induced by steroid hormone or a derivative thereof) etc.), the induction substance is added to the medium (or removed from the medium) at an appropriate stage to induce expression of the nucleic acid-modifying enzyme complex, culture is performed for a given period to carry out a nucleic acid base conversion reaction and, introduction of mutation into the target gene, transient expression of the nucleic acid-modifying enzyme complex can be realized.

[0148] In Prokaryotic cells such as *Escherichia coli* and the like, inducible promoters can also be used. Examples of such inducible promoters include, but are not limited to, lac promoter (induced by IPTG), cspA promoter (induced by cold shock), araBAD promoter (induced by arabinose) and the like.

[0149] Alternatively, the above-mentioned inducible promoters can also be utilized as a vector removal mechanism when higher eukaryotic cells such as animal cell, insect cell, plant cell and the like are used as a host cell. That is, a vector is loaded with a replication origin that can function in a host cell, and a nucleic acid encoding a protein necessary for replication thereof (e.g., SV40 ori and large T antigen, oriP and EBNA-1 etc. for animal cells), and the expression of the nucleic acid encoding the protein is regulated by the above-mentioned inducible promoter. As a result, while the vector is autonomously replicatable in the presence of an induction substance, when the induction substance is removed, autonomous replication does not occur, and the vector naturally falls off during cell division (conversely, autonomous replication becomes impossible by the addition of tetracycline and doxycycline in the case of Tet-OFF system vector).

[0150] The present invention is explained in the following by referring to Examples, which are not to be construed as limitative.

EXAMPLE

[0151] In the below-mentioned Examples 1-6, experiments were performed as follows.

<Cell Line, Culture, Transformation, and Expression Induction>

[0152] Budding yeast *Saccharomyces cerevisiae* BY4741 strain (requiring leucine and uracil) was cultured in a standard YPDA medium or SD medium with a Dropout composition satisfying the auxotrophicity. The culture performed was standing culture in an agar plate or shaking culture in a liquid medium between 25° C. and 30° C. Transformation was performed by a lithium acetate method, and selection was made in SD medium matching appropriate auxotrophicity. For expression induction by galactose, after preculture overnight in an appropriate SD medium, culture in SR medium overnight with carbon source changed from 2% glucose to 2% raffinose, and further culture in SGal medium for 3 hr to about two nights with carbon source changed to 0.2-2% galactose were conducted for expression induction.

[0153] For the measurement of the number of surviving cells and Can1 mutation rate, a cell suspension was appropriately diluted, and applied on SD plate medium and SD-Arg+60 mg/l Canavanine plate medium or SD+300 mg/l Canavanine plate medium, and the number of colonies that emerge 3 days later was counted as the number of surviving cells. Using the number of surviving colonies in SD plate as the total number of cells, and the number of surviving colonies in Canavanine plate as the number of resistant mutant strain, the mutation rate was calculated and evaluated. The site of mutation introduction was identified by amplifying DNA fragments containing the target gene region of each strain by a colony PCR method, performing DNA sequencing, and performing an alignment analysis based on the sequence of *Saccharomyces* Genome Database (www.yeastgenome.org).

<Nucleic Acid Operation>

[0154] DNA was processed or constructed by any of PCR method, restriction enzyme treatment, ligation, Gibson Assembly method, and artificial chemical synthesis. For plasmid, as the yeast/*Escherichia coli* shuttle vector, pRS315 for leucine selection and pRS426 for uracil selection were used as the backbone. Plasmid was amplified by *Escherichia coli* line XL-10 gold or DH5 α , and introduced into yeast by the lithium acetate method.

<Construct>

[0155] For inducible expression, budding yeast pGal1/10 (SEQ ID NO: 8), which is a bidirectional promoter inducible by galactose, was used. At the downstream of a promoter, a nuclear localization signal (ccc aag aag aag agg aag ggt; SEQ ID NO: 9(PKPKKRV; encoding SEQ ID NO: 10)) was added to Cas9 gene ORF from *Streptococcus pyogenes* having a codon optimized for eukaryon expression (SEQ ID NO: 5) and ORF (SEQ ID NO: 1 or 3) of deaminase gene (PmCDA1 from *Petromyza marinus* or hAID from human) was ligated via a linker sequence, and expressed as a fusion protein. As a linker sequence, GS linker (repeat of ggt gga gga ggt tct; SEQ ID NO: 11 (encoding GGGGS; SEQ ID NO: 12)), Flag® tag (gac tat aag gac cagcag gga gac tac aag gat cat gat att gat tac aaa gac gat gac gat aag; SEQ ID NO: 13 (encoding DYKDHDGDYKDHDIDYKDDDDK; SEQ ID NO: 14)), Strep-tag® (tgg agc cac ccg cag ttc gaa aaa; SEQ ID NO: 15 (encoding WSHQPQFEK; SEQ ID NO: 16)), and other domains are selected and used in combination. Here, particularly, 2xGS, SH3 domain (SEQ ID NO: 17 and 18), and Flag® tag were ligated and used. As a terminator, ADH1 terminator from budding yeast (SEQ ID NO: 19) and Top2 terminator (SEQ ID NO: 20) were ligated. In the domain integration method, Cas9 gene ORF was ligated to SH3 domain via 2xGS linker to give one protein, deaminase was added with SH3 ligand sequence (SEQ ID NOs: 21 and 22) as another protein, and they were ligated to Gal1/10 promoter on both sides. And they were simultaneously expressed. These were incorporated into pRS315 plasmid.

[0156] In Cas9, mutation to convert the 10th aspartic acid to alanine (D10A, corresponding DNA sequence mutation a29c) and mutation to convert the 840th histidine to alanine (H840A, corresponding DNA sequence mutation ca2518gc) were introduced to remove cleavage ability of either side of DNA strand.

[0157] gRNA as a chimeric structure with tracrRNA (from *Streptococcus pyogenes*; SEQ ID NO: 7) was disposed between SNR52 promoter (SEQ ID NO: 23) and Sup4 terminator (SEQ ID NO: 24), and incorporated into pRS426 plasmid. As gRNA target base sequence, CAN1 gene ORF, 187-206 (gatacgttctctatggagga; SEQ ID NO: 25) (target 1), 786-805 (ttggagaaaccaggtgcct; SEQ ID NO: 26) (target 3), 793-812 (aacccaggtgcctgggtcc; SEQ ID NO: 27) (target 4), 563-582 (ttggccaagtcattcaatt; SEQ ID NO: 28) (target 2) and complementary strand sequence of 767-786 (ataacg-gaatccaactgggc; SEQ ID NO: 29) (target 5r) were used. For simultaneous expression of a plurality of targets, using a sequence from a promoter to a terminator as one set, and a plurality of the sets were incorporated into the same plasmid. They were introduced into cells along with Cas9-deaminase expression plasmid, intracellularly expressed, and a complex of gRNA-tracrRNA and Cas9-deaminase was formed.

Example 1: Modification of Genome Sequence by Linking DNA Sequence Recognition Ability of CRISPR-Cas to Deaminase PmCDA1

[0158] To test the effect of genome sequence modification technique of the present invention by utilizing deaminase and CRISPR-Cas nucleic acid sequence recognition ability, introduction of mutation into CAN1 gene encoding canavanine transporter, whose gene deficit results in canavanine-resistance, was attempted. As gRNA, a sequence complementary to 187-206 (target 1) of CAN1 gene ORF was used, a chimeric RNA expression vector obtained by linking thereto tracrRNA from *Streptococcus pyogenes*, and a vector expressing a protein obtained by fusing dCas9 with impaired nuclease activity by introducing mutations (D10A and H840A) into Cas9 (SpCas9) from *Streptococcus pyogenes* with PmCDA1 from *Petromyzon marinus* as deaminase were constructed, introduced into the budding yeast by the lithium acetate method, and coexpressed. The results are shown in FIG. 2. When cultured on a canavanine-containing SD plate, only the cells subjected to introduction and expression of gRNA-tracrRNA and dCas9-PmCDA1 formed colony. The resistant colony was picked up and the sequence of CAN1 gene region was determined. As a result, it was confirmed that a mutation was introduced into the target nucleotide sequence (target 1) and the vicinity thereof.

Example 2: Drastic Reduction of side Effects•Toxicity

[0159] In conventional Cas9 and other artificial nucleases (ZFN, TALEN), inhibition of growth and cell death assumedly caused by disordered cleavage of chromosome occur by targeting a sequence in the genome. The effect thereof is particularly fatal for many microorganisms and prokaryotes, and prevents applicability.

[0160] Therefore, to verify the safety and cell toxicity of the genome sequence modification technique of the present invention, a comparative test with conventional CRISPR-Cas9 was performed. Using sequences (targets 3, 4) in the CAN1 gene as gRNA target, the surviving cells were counted immediately after the start of expression induction by galactose and at 6 hr after the induction based on the colony forming ability on an SD plate. The results are shown in FIG. 3. In conventional Cas9, the growth was inhibited and cell death was induced, which decreased the number of surviving cells. In contrast, by the present technique (nCas9

D10A-PmCDA1), the cells could continue to grow, and the number of surviving cells drastically increased.

Example 3: Use of Different Linking Scheme

[0161] Whether mutation can be introduced into a targeted gene even when Cas9 and deaminase are not used as a fusion protein but when a nucleic acid-modifying enzyme complex is formed via a binding domain and a ligand thereof was examined. As Cas9, dCas9 used in Example 1 was used and human AID instead of PmCDA1 was used as deaminase. SH3 domain was fused with the former, and a binding ligand thereof was fused with the latter to produce various constructs shown in FIG. 4. In addition, sequences (target 4,5r) in the CAN1 gene were used as gRNA targets. These constructs were introduced into a budding yeast. As a result, even when dCas9 and deaminase were linked via the binding domain, mutation was efficiently introduced into the targeted site of the CAN1 gene (FIG. 4). The mutation introduction efficiency was remarkably improved by introducing a plurality of binding domains into dCas9. The main site of mutation introduction was 782nd (g782c) of ORF.

Example 4: High Efficiency and Changes in Mutation Pattern by Use of Nickase

[0162] In the same manner as in Example 1 except that D10A mutant nCas9 (D10A) that cleaves only a strand complementary to gRNA, or H840A mutant nCas9 (H840A) that cleaves only a strand opposite to a strand complementary to gRNA was used instead of dCas9, mutation was introduced into the CAN1 gene, and the sequence in the CAN1 gene region of the colony generated on a canavanine-containing SD plate was examined. It was found that the efficiency increases in the former (nCas9 (D10A)) as compared to dCas9 (FIG. 5), and the mutation gathers in the center of the target sequence (FIG. 6). Therefore, this method enables pinpoint introduction of mutation. On the other hand, it was found in the latter (nCas9 (H840A)) that a plurality of random mutations were introduced into a region of several hundred bases from the targeted nucleotide (FIG. 6) along with an increase in the efficiency as compared to dCas9 (FIG. 5).

[0163] Similar remarkable introduction of mutation could be confirmed even when the target nucleotide sequence was changed. In this genome editing system using CRISPR-Cas9 system and cytidine deaminase, it was confirmed as shown in Table 1 that cytosine present within the range of about 2-5 bp from the 5'-side of the target nucleotide sequence (20 bp) were preferentially deaminated. Therefore, by setting the target nucleotide sequence based on this regularity and further combining with nCas9 (D10A), precise genome editing of 1 nucleotide unit is possible. On the other hand, a plurality of mutations can be simultaneously inserted within the range of about several hundred bp in the vicinity of the target nucleotide sequence by using nCas9 (H840A). Furthermore, site specificity may possibly be further varied by changing the linking scheme of deaminase.

[0164] These results show that the kind of Cas9 protein can be changed properly according to the object.

TABLE 1

position of CAN1 gene ORF	sequence (SEQ ID NO:)	site of main mutation introduction
187-206 (target 1)	Gatacgttctctatggagga (25)	c191a, g226t
563-582 (target 2)	Ttgccaagtcattcaattt (28)	cc567at, c567del
786-805 (target 3) and 793-812 (target 4)	Ttgagaaacccaggtgcct (26) Aaccaggtgcctgggtcc (27)	cc795tt, cc796tt
767-786 (complementary strand) (target 5r)	Ataacggaatccaactgggc (29)	g782c

Example 5: Efficiency Increases Synergistically by Targeting a Plurality of DNA Sequences in Proximity

[0165] Efficiency drastically increased by simultaneously using a plurality of targets in proximity rather than a single target (FIG. 7). In fact, 10-20% of cells had canavanine-resistant mutation (targets 3, 4). In the Figure, gRNA1 and gRNA2 target target 3 and target 4, respectively. As deaminase, PmCDA1 was used. The effect thereof was confirmed to occur not only when the sequences partly overlapped (targets 3, 4), but also when they were apart by about 600 bp (targets 1, 3). The effect was found both when the DNA sequences were in the same direction (targets 3, 4) and opposing (targets 4, 5) (FIG. 4).

Example 6: Genetic Modification not Requiring Selection Marker

[0166] As for the cells (Targets 3, 4) in which target 3 and target 4 were targeted in Example 5, 10 colonies were randomly selected from those grown on a non-selected (canavanine-free) plate (SD plate not containing Leu and Ura) and the sequences of the CAN1 gene region were determined. As a result, mutation was introduced into the targeted site of the CAN1 gene in all examined colonies (FIG. 8). That is, editing can be expected in almost all expressed cells by selecting a suitable target sequence according to the present invention. Therefore, insertion of a marker gene and selection, which are essential for the conventional gene manipulation, are not necessary. This dramatically facilitates and simplifies gene manipulation and extends the applicability to crop breeding and the like since a recombinant organism with foreign DNA is not produced.

[0167] In the following Examples, experiment techniques shared by Examples 1-6 were performed in the same manner as above.

Example 7: Simultaneous Editing of a Plurality of Sites (Different Gene)

[0168] In a general gene manipulation method, mutation of only one site is generally achieved by one operation due to various restrictions. Thus, whether a simultaneous mutation operation of a plurality of sites is possible using the method of the present invention was tested.

[0169] Using the ORF of positions 3 to 22 of Ade1 gene of budding yeast BY4741 strain as the first target nucleotide sequence (Ade1 target 5:GTCAATTACGAAGACTGAAC; SEQ ID NO: 30), and the ORF of positions 767-786 (complementary strand) of Can1 gene as the second target nucleotide sequence (Can1 target8 (786-767; ATAACG-GAATCCAACCTGGGC; SEQ ID NO: 29), both DNAs encoding chimeric RNAs of two kinds of gRNAs each containing a nucleotide sequence complementary thereto and tracrRNA (SEQ ID NO: 7) were placed on the same plasmid (pRS426), and introduced into BY4741 strain together with plasmid nCas9 D10A-PmCDA1 containing a nucleic acid encoding a fusion protein of mutant Cas9 and PmCDA1, and expressed, and introduction of mutation into the both genes was verified. The cells were cultured in an SD drop-out medium (uracil and leucine deficient; SD-UL) as a base, which maintains plasmid. The cells were appropriately diluted, applied on SD-UL and canavanine addition medium and allowed to form a colony. After 2 days of culture at 28° C., colonies were observed, and the incidence of red colony due to Ade1 mutation, and the survival rate in a canavanine medium were respectively counted. The results are shown in Table 2.

TABLE 2

medium	incidence of red colony	survival rate in Canavanine medium (Can)	red colony and Can survival rate
SD-UL	0.54 ± 0.04		
+canavanine	0.64 ± 0.14	0.51 ± 0.15	0.31 ± 0.04

[0170] As a phenotype, the proportion of introduction of mutation into both Ade1 gene and Can1 gene was high and about 31%.

[0171] Then, a colony on an SD-UL medium was subjected to PCR amplification followed by sequencing. Regions containing ORF of each of Ade1 and Can1 were amplified, and sequence information of about 500 b sequences surrounding the target sequence was obtained. To be specific, 5 red colonies and 5 white colonies were analyzed to find conversion of the 5th C of Ade1 gene ORF to G in all red colonies and the 5th C to T in all white colonies (FIG. 9). While the mutation rate of the target is 100%, as the mutation rate in light of the object of gene destruction, the desired mutation rate is 50% since the 5th C needs to be changed to G to be a stop codon. Similarly, as for the Can1 gene, mutation was confirmed in the 782nd G of ORF in all clones (FIG. 9); however, since only the mutation to C affords canavanine-resistance, the desired mutation rate is 70%. Desired mutations in both genes were simultaneously obtained in 40% clones (4 clones out of 10 clones) by investigation, and practically high efficiency was obtained.

Example 8: Editing of Polyploid Genome

[0172] Many organisms have diploid or polyploid genome. In the conventional mutation introduction methods, mutation is, in principle, introduced into only one homologous chromosome to produce a heterologous genotype. Therefore, desired feature is not obtained unless it is a dominant mutation, and making it homologous requires labor and time. Thus, whether the technique of the present

invention can introduce mutation into all target alleles on the homologous chromosome in the genome was tested.

[0173] That is, simultaneous editing of Ade1 and Can1 genes was performed in budding yeast YPH501 strain as a diploid strain. The phenotype of these gene mutations (red colony and canavanine-resistant) is a recessive phenotype, and therefore, these phenotypes do not appear unless both mutations of homologous gene (homologous mutation) are introduced.

[0174] Using the ORF of positions 1173-1154 (complementary strand) of Ade1 gene (Ade1 target 1: GTCAATAGGATCCCTTTT; SEQ ID NO: 31) or of positions 3-22 (Ade1 target 5: GTCAATTACGAAGACTGAAC; SEQ ID NO: 30) as the first target nucleotide sequence, and the ORF of positions 767-786 (complementary strand) of Can1 gene as the second target nucleotide sequence (Can1 target8: ATAACGGAATCCAACCTGGGC; SEQ ID NO: 29), both DNAs encoding chimeric RNAs of two kinds of gRNAs each containing a nucleotide sequence complementary thereto and tracrRNA (SEQ ID NO: 7) were placed on the same plasmid (pRS426), and introduced into BY4741 strain together with plasmid nCas9 D10A-PmCDA1 containing a nucleic acid encoding a fusion protein of mutant Cas9 and PmCDA1, and expressed, and introduction of mutation into each gene was verified.

[0175] As a result of colony count, it was found that each characteristic of phenotype could be obtained at a high probability (40% - 70%) (FIG. 10A).

[0176] To confirm mutation, Ade1 target region of each of white colony and red colony was sequenced to confirm overlapping of sequence signals indicating heterologous mutation in the target site of white colony (FIG. 10B, upper panel, G and T signals overlap at ↓). Phenotype was confirmed to be absent in colony with heterologous mutation. On the other hand, homologous mutation free of overlapping signal was confirmed in red colony (FIG. 10B, lower panel, T signal at ↓).

Example 9: Genome Editing in *Escherichia coli*

[0177] In this Example, it is demonstrated that this technique effectively functions in *Escherichia coli*, which is a representative bacterium model organism. Particularly, conventional nuclease type genome editing technique is fatal for bacteria, and the application is difficult. Thus, the superiority of this technique is emphasized. In combination with yeast, which is an eukaryote model cell, it is shown that this technique is widely applicable to any species irrespective of prokaryon and eukaryon.

[0178] Amino acid mutation of DIOA and H840A were introduced (dCas9) into *Streptococcus pyogenes* Cas9 gene containing bidirectional promoter region, and a construct to be expressed as a fusion protein with PmCDA1 via a linker sequence was constructed, and chimeric gRNAs encoding a sequence complementary to each of the target nucleotide sequences was simultaneously included in a plasmid (full-length nucleotide sequence is shown in SEQ ID NO: 32, in which sequence, a sequence complementary to each of the target sequences is introduced into the site of n₂₀) (FIG. 11A).

[0179] First, the ORF of positions 426-445 (T CAA TGG GCT AAC TAC GTT C; SEQ ID NO: 33) of *Escherichia coli* galK gene was introduced as a target nucleotide sequence into a plasmid, various *Escherichia coli* strains (XL10-gold, DH5a, MG1655, BW25113) were transformed

with the plasmid by calcium method or electroporation method, SOC medium was added, recovery culture was performed overnight, plasmid carrying cells were selected from ampicillin-containing LB medium, and colony was formed. Introduction of mutation was verified by direct-sequence from colony PCR. The results are shown in FIG. 11B.

[0180] Independent colony (1-3) was selected randomly, and sequence was analyzed. As a result, the 427-position C of ORF was converted to T (clones 2, 3) at a probability of not less than 60%, and the occurrence of gene destruction generating a stop codon (TAA) was confirmed.

[0181] Then, with a complementary sequence (5'-GGTC-CATAAAGTGGAGACAGC-3'; SEQ ID NO: 34) of 1530-1549 base region of rpoB gene ORF, which is an essential gene, as a target, particular point mutation was introduced by a method similar to the above-mentioned method to try to impart rifampicin-resistant function to *Escherichia coli*. The sequences of colonies selected in a nonselective medium (none), a 25 µg/ml rifampicin (Rif25) and 50 µg/ml rifampicin (Rif50)-containing medium were analyzed. As a result, it was confirmed that conversion of the 1546-position G of ORF to A introduced amino acid mutation from Asp(GAC) to Asn(AAC), and rifampicin-resistance was imparted (FIG. 11C, upper panel). A 10-fold dilution series of the cell suspension after transformation treatment was spotted on a nonselective medium (none), a 25 µg/ml rifampicin (Rif25) and 50 µg/ml rifampicin (Rif50)-containing medium and cultured. As a result, it is estimated that rifampicin-resistant strain was obtained at about 10% frequency (FIG. 11C, lower panel).

[0182] As shown above, by this technique, a new function can be added by particular point mutation, rather than simple gene destruction. This technique is superior since essential gene is directly edited.

Example 10: Adjustment of editing base site by gRNA length

[0183] Conventionally, the gRNA length relative to a target nucleotide sequence was 20b as basic, and cytosine (or guanine in opposite strand) in a site of 2-5b from the 5'-terminus thereof (15-19b upstream of PAM sequence) is used as a mutation target. Whether expression of different gRNA length can shift the site of the base to be the target was examined (FIG. 12A).

[0184] Experimental Example performed on *Escherichia coli* is shown in FIG. 12B. A site containing many cytosines on *Escherichia coli* genome was searched for, and the experiment was performed using *gsiA* gene, which is a putative ABC-transporter. Substituted cytosine was examined while changing the length of the target to 24 bp, 22 bp, 20 bp, 18 bp to find that the 898th, 899th cytosine was substituted by thymine in the case of 20 bp (standard length). When the target site is longer than 20 bp, the 896th and 897th cytosines were also substituted, and when the target site was shorter, the 900th and 901st cytosines were also substituted. In fact, the target site could be shifted by changing the length of the gRNA.

Example 11: Development of Temperature Dependent Genome Editing Plasmid

[0185] A plasmid that induces expression of the nucleic acid-modifying enzyme complex of the present invention

under high temperature conditions was designed. While optimizing efficiency by limitatively controlling the expression state, reduction of side effects (growth inhibition of host, unstable mutation introduction efficiency, mutation of site different from target and the like) was aimed. Simultaneously, a simultaneous and easy removal of plasmid after editing was intended by combining a mechanism for ceasing the replication of plasmid at a high temperature. The detail of the experiment is shown below.

[0186] With temperature sensitive plasmid pSC101-Rep101 system (sequence of pSC101 ori is shown in SEQ ID NO: 35, and sequence of temperature sensitive Rep101 is shown in SEQ ID NO: 36) as a backbone, temperature sensitive A repressor (c1857) system was used for expression induction. For genome editing, G113E mutation imparting RecA resistance was introduced into A repressor, to ensure normal function even under SOS response (SEQ ID NO: 37). dCas9-PmCDA1 (SEQ ID NO: 38) was ligated to Right Operator (SEQ ID NO: 39), and gRNA (SEQ ID NO: 40) was ligated to the downstream of Left Operator (SEQ ID NO: 41) to regulate the expression (full-length nucleotide sequence of the constructed expression vector is shown in SEQ ID NO: 42). During culture at not more than 30° C., transcription of gRNA and expression of dCas9-PmCDA1 are suppressed, and the cells can grow normally. When cultured at not less than 37° C., transcription of gRNA and expression of dCas9-PmCDA1 are induced, and replication of plasmid is suppressed simultaneously. Therefore, a nucleic acid-modifying enzyme complex necessary for genome editing is transiently supplied, and plasmid can be removed easily after editing (FIG. 13).

[0187] Specific protocol of the base substitution is shown in FIG. 14.

[0188] The culture temperature for plasmid construction is set at around 28° C., and an *Escherichia coli* colony retaining the desired plasmid is first established. Then, the colony is directly used, or after plasmid extraction when the strain is changed, transformation with the target strain is performed again, and the obtained colony is used. Liquid culture at 28° C. is performed overnight. Thereafter, the colony is diluted with the medium, induction culture is performed at 42° C. for 1 hr to overnight, the cell suspension is appropriately diluted and spread or spotted on a plate to acquire a single colony.

[0189] As a verification experiment, point mutation introduction into essential gene rpoB was performed. When rpoB, which is one of the RNA polymerase-constituting factors, is deleted or its function is lost, the *Escherichia coli* will not survive. On the other hand, it is known that resistance to antibiotic rifampicin (Rif) is acquired when point mutation is entered at a particular site. Therefore, aiming at such introduction of point mutation, a target site is selected and assay was performed.

[0190] The results are shown in FIG. 15. In the upper left panel, the left shows an LB (chloramphenicol addition) plate, and the right shows a rifampicin-added LB (chloram-

phenicol addition) plate, and samples with or without chloramphenicol were prepared and cultured at 28° C. or 42° C. When cultured at 28° C., the rate of Rif resistance is low; however, when cultured at 42° C., rifampicin resistance was obtained with extremely high efficiency. When the colonies (non-selection) obtained on LB were sequenced by 8 colonies, the 1546th guanine (G) was substituted by adenine (A) in not less than 60% of the strain cultured at 42° C. (lower and upper left panels). It is clear that the base is also completely substituted in actual sequence spectrum (lower right panel).

[0191] Similarly, base substitution of galK, which is one of the factors involved in the galactose metabolism, was performed. Since metabolism of 2-deoxy-galactose (2DOG), which is an analogue of galactose, by galK is fatal to *Escherichia coli*, this was used as a selection method. Target site was set such that missense mutation is induced in target 8, and that stop codon is entered in target 12 (FIG. 16 lower right).

[0192] The results are shown in FIG. 16. In the upper left and lower left panels, the left shows an LB (chloramphenicol addition) plate, and the right shows a 2-DOG-added LB (chloramphenicol addition) plate, and samples with or without chloramphenicol were prepared and cultured at 28° C. or 42° C. In target 8, colony was produced only slightly on a 2-DOG addition plate (upper left panel), 3 colonies on LB (red frame) were sequenced to determine that the 61st cytosine (C) was substituted by thymine (T) in all colonies (upper right). This mutation is assumed to be insufficient to lose function of galK. On the other hand, in target 12, colony was obtained on 2-DOG addition plate by culture at 28° C. and 42° C. (lower left panel). 3 colonies on LB were sequenced to determine that the 271st cytosine was substituted by thymine in all colonies (lower right). It was shown that mutation can be also introduced stably and highly efficiently in such different targets.

[0193] The contents disclosed in any publication cited herein, including patents and patent applications, are hereby incorporated in their entireties by reference, to the extent that they have been disclosed herein.

[0194] This application is based on patent application Nos. 2014-43348 and 2014-201859 filed in Japan (filing dates: Mar. 5, 2014 and Sep. 30, 2014, respectively), the contents of which are incorporated in full herein.

INDUSTRIAL APPLICABILITY

[0195] The present invention makes it possible to safely introduce site specific mutation into any species without insertion of a foreign DNA or double-stranded DNA breaks. It is also possible to set a wide range of mutation introduction from a pin point of one base to several hundred bases, and the technique can also be applied to topical evolution induction by introduction of random mutation into a particular restricted region, which has been almost impossible heretofore, and is extremely useful.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 95

<210> SEQ ID NO 1

<211> LENGTH: 624

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Petromyzon marinus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(624)

<400> SEQUENCE: 1

atg acc gac gct gag tac gtg aga atc cat gag aag ttg gac atc tac      48
Met Thr Asp Ala Glu Tyr Val Arg Ile His Glu Lys Leu Asp Ile Tyr
1           5           10

acg ttt aag aaa cag ttt ttc aac aac aaa aaa tcc gtg tcg cat aga      96
Thr Phe Lys Lys Gln Phe Phe Asn Asn Lys Lys Ser Val Ser His Arg
           20           25           30

tgc tac gtt ctc ttt gaa tta aaa cga cgg ggt gaa cgt aga gcg tgt     144
Cys Tyr Val Leu Phe Glu Leu Lys Arg Arg Gly Glu Arg Arg Ala Cys
           35           40           45

ttt tgg ggc tat gct gtg aat aaa cca cag agc ggg aca gaa cgt ggc     192
Phe Trp Gly Tyr Ala Val Asn Lys Pro Gln Ser Gly Thr Glu Arg Gly
50           55           60

att cac gcc gaa atc ttt agc att aga aaa gtc gaa gaa tac ctg cgc     240
Ile His Ala Glu Ile Phe Ser Ile Arg Lys Val Glu Glu Tyr Leu Arg
65           70           75           80

gac aac ccc gga caa ttc acg ata aat tgg tac tca tcc tgg agt cct     288
Asp Asn Pro Gly Gln Phe Thr Ile Asn Trp Tyr Ser Ser Trp Ser Pro
           85           90           95

tgt gca gat tgc gct gaa aag atc tta gaa tgg tat aac cag gag ctg     336
Cys Ala Asp Cys Ala Glu Lys Ile Leu Glu Trp Tyr Asn Gln Glu Leu
           100          105          110

cgg ggg aac ggc cac act ttg aaa atc tgg gct tgc aaa ctc tat tac     384
Arg Gly Asn Gly His Thr Leu Lys Ile Trp Ala Cys Lys Leu Tyr Tyr
           115          120          125

gag aaa aat gcg agg aat caa att ggg ctg tgg aac ctc aga gat aac     432
Glu Lys Asn Ala Arg Asn Gln Ile Gly Leu Trp Asn Leu Arg Asp Asn
130          135          140

ggg gtt ggg ttg aat gta atg gta agt gaa cac tac caa tgt tgc agg     480
Gly Val Gly Leu Asn Val Met Val Ser Glu His Tyr Gln Cys Cys Arg
145          150          155          160

aaa ata ttc atc caa tcg tcg cac aat caa ttg aat gag aat aga tgg     528
Lys Ile Phe Ile Gln Ser Ser His Asn Gln Leu Asn Glu Asn Arg Trp
           165          170          175

ctt gag aag act ttg aag cga gct gaa aaa cga cgg agc gag ttg tcc     576
Leu Glu Lys Thr Leu Lys Arg Ala Glu Lys Arg Arg Ser Glu Leu Ser
           180          185          190

att atg att cag gta aaa ata ctc cac acc act aag agt cct gct gtt     624
Ile Met Ile Gln Val Lys Ile Leu His Thr Thr Lys Ser Pro Ala Val
           195          200          205

```

```

<210> SEQ ID NO 2
<211> LENGTH: 208
<212> TYPE: PRT
<213> ORGANISM: Petromyzon marinus

<400> SEQUENCE: 2

Met Thr Asp Ala Glu Tyr Val Arg Ile His Glu Lys Leu Asp Ile Tyr
1           5           10           15

Thr Phe Lys Lys Gln Phe Phe Asn Asn Lys Lys Ser Val Ser His Arg
           20           25           30

Cys Tyr Val Leu Phe Glu Leu Lys Arg Arg Gly Glu Arg Arg Ala Cys
           35           40           45

```

-continued

Phe Trp Gly Tyr Ala Val Asn Lys Pro Gln Ser Gly Thr Glu Arg Gly
 50 55 60

Ile His Ala Glu Ile Phe Ser Ile Arg Lys Val Glu Glu Tyr Leu Arg
 65 70 75 80

Asp Asn Pro Gly Gln Phe Thr Ile Asn Trp Tyr Ser Ser Trp Ser Pro
 85 90 95

Cys Ala Asp Cys Ala Glu Lys Ile Leu Glu Trp Tyr Asn Gln Glu Leu
 100 105 110

Arg Gly Asn Gly His Thr Leu Lys Ile Trp Ala Cys Lys Leu Tyr Tyr
 115 120 125

Glu Lys Asn Ala Arg Asn Gln Ile Gly Leu Trp Asn Leu Arg Asp Asn
 130 135 140

Gly Val Gly Leu Asn Val Met Val Ser Glu His Tyr Gln Cys Cys Arg
 145 150 155 160

Lys Ile Phe Ile Gln Ser Ser His Asn Gln Leu Asn Glu Asn Arg Trp
 165 170 175

Leu Glu Lys Thr Leu Lys Arg Ala Glu Lys Arg Arg Ser Glu Leu Ser
 180 185 190

Ile Met Ile Gln Val Lys Ile Leu His Thr Thr Lys Ser Pro Ala Val
 195 200 205

<210> SEQ ID NO 3
 <211> LENGTH: 600
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(600)

<400> SEQUENCE: 3

atg gac agc ctc ttg atg aac cgg agg aag ttt ctt tac caa ttc aaa 48
 Met Asp Ser Leu Leu Met Asn Arg Arg Lys Phe Leu Tyr Gln Phe Lys
 1 5 10 15

aat gtc cgc tgg gct aag ggt cgg cgt gag acc tac ctg tgc tac gta 96
 Asn Val Arg Trp Ala Lys Gly Arg Arg Glu Thr Tyr Leu Cys Tyr Val
 20 25 30

gtg aag agg cgt gac agt gct aca tcc ttt tca ctg gac ttt ggt tat 144
 Val Lys Arg Arg Asp Ser Ala Thr Ser Phe Ser Leu Asp Phe Gly Tyr
 35 40 45

ctt cgc aat aag aac ggc tgc cac gtg gaa ttg ctc ttc ctc cgc tac 192
 Leu Arg Asn Lys Asn Gly Cys His Val Glu Leu Leu Phe Leu Arg Tyr
 50 55 60

atc tcg gac tgg gac cta gac cct ggc cgc tgc tac cgc gtc acc tgg 240
 Ile Ser Asp Trp Asp Leu Asp Pro Gly Arg Cys Tyr Arg Val Thr Trp
 65 70 75 80

ttc acc tcc tgg agc ccc tgc tac gac tgt gcc cga cat gtg gcc gac 288
 Phe Thr Ser Trp Ser Pro Cys Tyr Asp Cys Ala Arg His Val Ala Asp
 85 90 95

ttt ctg cga ggg aac ccc tac ctc agt ctg agg atc ttc acc gcg cgc 336
 Phe Leu Arg Gly Asn Pro Tyr Leu Ser Leu Arg Ile Phe Thr Ala Arg
 100 105 110

ctc tac ttc tgt gag gac cgc aag gct gag ccc gag ggg ctg cgg cgg 384
 Leu Tyr Phe Cys Glu Asp Arg Lys Ala Glu Pro Glu Gly Leu Arg Arg
 115 120 125

ctg cac cgc gcc ggg gtg caa ata gcc atc atg acc ttc aaa gat tat 432
 Leu His Arg Ala Gly Val Gln Ile Ala Ile Met Thr Phe Lys Asp Tyr

-continued

130	135	140	
ttt tac tgc tgg aat act	ttt gta gaa aac cat	gaa aga act ttc aaa	480
Phe Tyr Cys Trp Asn Thr	Phe Val Glu Asn His	Glu Arg Thr Phe Lys	
145	150	155	160
gcc tgg gaa ggg ctg cat	gaa aat tca gtt cgt	ctc tcc aga cag ctt	528
Ala Trp Glu Gly Leu His	Glu Asn Ser Val Arg	Leu Ser Arg Gln Leu	
	165	170	175
cgg cgc atc ctt ttg ccc	ctg tat gag gtt gat	gac tta cga gac gca	576
Arg Arg Ile Leu Leu Pro	Leu Tyr Glu Val Asp	Asp Leu Arg Asp Ala	
	180	185	190
ttt cgt act ttg gga ctt	ctc gac		600
Phe Arg Thr Leu Gly Leu	Leu Asp		
	195	200	

<210> SEQ ID NO 4
 <211> LENGTH: 200
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Asp Ser Leu Leu Met	Asn Arg Arg Lys Phe	Leu Tyr Gln Phe Lys		
1	5	10	15	
Asn Val Arg Trp Ala Lys	Gly Arg Arg Glu Thr	Tyr Leu Cys Tyr Val		
	20	25	30	
Val Lys Arg Arg Asp Ser	Ala Thr Ser Phe Ser	Leu Asp Phe Gly Tyr		
	35	40	45	
Leu Arg Asn Lys Asn Gly	Cys His Val Glu Leu	Leu Phe Leu Arg Tyr		
	50	55	60	
Ile Ser Asp Trp Asp Leu	Asp Pro Gly Arg Cys	Tyr Arg Val Thr Trp		
	65	70	75	80
Phe Thr Ser Trp Ser Pro	Cys Tyr Asp Cys Ala	Arg His Val Ala Asp		
	85	90	95	
Phe Leu Arg Gly Asn Pro	Tyr Leu Ser Leu Arg	Ile Phe Thr Ala Arg		
	100	105	110	
Leu Tyr Phe Cys Glu Asp	Arg Lys Ala Glu Pro	Glu Gly Leu Arg Arg		
	115	120	125	
Leu His Arg Ala Gly Val	Gln Ile Ala Ile Met	Thr Phe Lys Asp Tyr		
	130	135	140	
Phe Tyr Cys Trp Asn Thr	Phe Val Glu Asn His	Glu Arg Thr Phe Lys		
	145	150	155	160
Ala Trp Glu Gly Leu His	Glu Asn Ser Val Arg	Leu Ser Arg Gln Leu		
	165	170	175	
Arg Arg Ile Leu Leu Pro	Leu Tyr Glu Val Asp	Asp Leu Arg Asp Ala		
	180	185	190	
Phe Arg Thr Leu Gly Leu	Leu Asp			
	195	200		

<210> SEQ ID NO 5
 <211> LENGTH: 4116
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Streptococcus pyogenes-derived Cas9 CDS
 optimized for eucaryotic expression.
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(4116)

-continued

<400> SEQUENCE: 5

atg gac aag aag tac tcc att ggg ctc gat atc ggc aca aac agc gtc	48
Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser Val	
1 5 10 15	
ggg tgg gcc gtc att acg gac gag tac aag gtg ccg agc aaa aaa ttc	96
Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys Phe	
20 25 30	
aaa gtt ctg ggc aat acc gat cgc cac agc ata aag aag aac ctc att	144
Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu Ile	
35 40 45	
ggc gcc ctc ctg ttc gac tcc ggg gag acg gcc gaa gcc acg cgg ctc	192
Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg Leu	
50 55 60	
aaa aga aca gca cgg cgc aga tat acc cgc aga aag aat cgg atc tgc	240
Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys	
65 70 75 80	
tac ctg cag gag atc ttt agt aat gag atg gct aag gtg gat gac tct	288
Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp Ser	
85 90 95	
ttc ttc cat agg ctg gag gag tcc ttt ttg gtg gag gag gat aaa aag	336
Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys Lys	
100 105 110	
cac gag cgc cac cca atc ttt ggc aat atc gtg gac gag gtg gcg tac	384
His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala Tyr	
115 120 125	
cat gaa aag tac cca acc ata tat cat ctg agg aag aag ctt gta gac	432
His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val Asp	
130 135 140	
agt act gat aag gct gac ttg cgg ttg atc tat ctc gcg ctg gcg cat	480
Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala His	
145 150 155 160	
atg atc aaa ttt cgg gga cac ttc ctc atc gag ggg gac ctg aac cca	528
Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn Pro	
165 170 175	
gac aac agc gat gtc gac aaa ctc ttt atc caa ctg gtt cag act tac	576
Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr Tyr	
180 185 190	
aat cag ctt ttc gaa gag aac ccg atc aac gca tcc gga gtt gac gcc	624
Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp Ala	
195 200 205	
aaa gca atc ctg agc gct agg ctg tcc aaa tcc cgg cgg ctc gaa aac	672
Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu Asn	
210 215 220	
ctc atc gca cag ctc cct ggg gag aag aag aac ggc ctg ttt ggt aat	720
Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly Asn	
225 230 235 240	
ctt atc gcc ctg tca ctc ggg ctg acc ccc aac ttt aaa tct aac ttc	768
Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn Phe	
245 250 255	
gac ctg gcc gaa gat gcc aag ctt caa ctg agc aaa gac acc tac gat	816
Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr Asp	
260 265 270	
gat gat ctc gac aat ctg ctg gcc cag atc ggc gac cag tac gca gac	864
Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala Asp	
275 280 285	
ctt ttt ttg gcg gca aag aac ctg tca gac gcc att ctg ctg agt gat	912

-continued

Leu	Phe	Leu	Ala	Ala	Lys	Asn	Leu	Ser	Asp	Ala	Ile	Leu	Leu	Ser	Asp	
290						295					300					
att	ctg	cga	gtg	aac	acg	gag	atc	acc	aaa	gct	ccg	ctg	agc	gct	agt	960
Ile	Leu	Arg	Val	Asn	Thr	Glu	Ile	Thr	Lys	Ala	Pro	Leu	Ser	Ala	Ser	
305					310					315					320	
atg	atc	aag	cgc	tat	gat	gag	cac	cac	caa	gac	ttg	act	ttg	ctg	aag	1008
Met	Ile	Lys	Arg	Tyr	Asp	Glu	His	His	Gln	Asp	Leu	Thr	Leu	Leu	Lys	
				325					330						335	
gcc	ctt	gtc	aga	cag	caa	ctg	cct	gag	aag	tac	aag	gaa	att	ttc	ttc	1056
Ala	Leu	Val	Arg	Gln	Gln	Leu	Pro	Glu	Lys	Tyr	Lys	Glu	Ile	Phe	Phe	
			340					345					350			
gat	cag	tct	aaa	aat	ggc	tac	gcc	gga	tac	att	gac	ggc	gga	gca	agc	1104
Asp	Gln	Ser	Lys	Asn	Gly	Tyr	Ala	Gly	Tyr	Ile	Asp	Gly	Gly	Ala	Ser	
			355				360						365			
cag	gag	gaa	ttt	tac	aaa	ttt	att	aag	ccc	atc	ttg	gaa	aaa	atg	gac	1152
Gln	Glu	Glu	Phe	Tyr	Lys	Phe	Ile	Lys	Pro	Ile	Leu	Glu	Lys	Met	Asp	
			370			375					380					
ggc	acc	gag	gag	ctg	ctg	gta	aag	ctt	aac	aga	gaa	gat	ctg	ttg	cgc	1200
Gly	Thr	Glu	Glu	Leu	Leu	Val	Lys	Leu	Asn	Arg	Glu	Asp	Leu	Leu	Arg	
385					390					395					400	
aaa	cag	cgc	act	ttc	gac	aat	gga	agc	atc	ccc	cac	cag	att	cac	ctg	1248
Lys	Gln	Arg	Thr	Phe	Asp	Asn	Gly	Ser	Ile	Pro	His	Gln	Ile	His	Leu	
				405					410						415	
ggc	gaa	ctg	cac	gct	atc	ctc	agg	cgg	caa	gag	gat	ttc	tac	ccc	ttt	1296
Gly	Glu	Leu	His	Ala	Ile	Leu	Arg	Arg	Gln	Glu	Asp	Phe	Tyr	Pro	Phe	
			420					425						430		
ttg	aaa	gat	aac	agg	gaa	aag	att	gag	aaa	atc	ctc	aca	ttt	cgg	ata	1344
Leu	Lys	Asp	Asn	Arg	Glu	Lys	Ile	Glu	Lys	Ile	Leu	Thr	Phe	Arg	Ile	
			435				440						445			
ccc	tac	tat	gta	ggc	ccc	ctc	gcc	cgg	gga	aat	tcc	aga	ttc	gcg	tgg	1392
Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg	Gly	Asn	Ser	Arg	Phe	Ala	Trp	
			450			455					460					
atg	act	cgc	aaa	tca	gaa	gag	acc	atc	act	ccc	tgg	aac	ttc	gag	gaa	1440
Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile	Thr	Pro	Trp	Asn	Phe	Glu	Glu	
465					470					475					480	
gtc	gtg	gat	aag	ggg	gcc	tct	gcc	cag	tcc	ttc	atc	gaa	agg	atg	act	1488
Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln	Ser	Phe	Ile	Glu	Arg	Met	Thr	
				485					490					495		
aac	ttt	gat	aaa	aat	ctg	cct	aac	gaa	aag	gtg	ctt	cct	aaa	cac	tct	1536
Asn	Phe	Asp	Lys	Asn	Leu	Pro	Asn	Glu	Lys	Val	Leu	Pro	Lys	His	Ser	
				500					505					510		
ctg	ctg	tac	gag	tac	ttc	aca	ggt	tat	aac	gag	ctc	acc	aag	gtc	aaa	1584
Leu	Leu	Tyr	Glu	Tyr	Phe	Thr	Val	Tyr	Asn	Glu	Leu	Thr	Lys	Val	Lys	
			515				520							525		
tac	gtc	aca	gaa	ggg	atg	aga	aag	cca	gca	ttc	ctg	tct	gga	gag	cag	1632
Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro	Ala	Phe	Leu	Ser	Gly	Glu	Gln	
			530			535					540					
aag	aaa	gct	atc	gtg	gac	ctc	ctc	ttc	aag	acg	aac	cgg	aaa	gtt	acc	1680
Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe	Lys	Thr	Asn	Arg	Lys	Val	Thr	
545					550					555					560	
gtg	aaa	cag	ctc	aaa	gaa	gac	tat	ttc	aaa	aag	att	gaa	tgt	ttc	gac	1728
Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe	Lys	Lys	Ile	Glu	Cys	Phe	Asp	
				565					570					575		
tct	ggt	gaa	atc	agc	gga	gtg	gag	gat	cgc	ttc	aac	gca	tcc	ctg	gga	1776
Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu	Gly	
				580				585						590		
acg	tat	cac	gat	ctc	ctg	aaa	atc	att	aaa	gac	aag	gac	ttc	ctg	gac	1824

-continued

Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu	Asp	
		595					600						605			
aat	gag	gag	aac	gag	gac	att	ctt	gag	gac	att	gtc	ctc	acc	ctt	acg	1872
Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu	Thr	
	610					615					620					
ttg	ttt	gaa	gat	agg	gag	atg	att	gaa	gaa	cgc	ttg	aaa	act	tac	gct	1920
Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu	Glu	Arg	Leu	Lys	Thr	Tyr	Ala	
	625				630					635					640	
cat	ctc	ttc	gac	gac	aaa	gtc	atg	aaa	cag	ctc	aag	agg	cgc	cga	tat	1968
His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys	Gln	Leu	Lys	Arg	Arg	Arg	Tyr	
				645					650					655		
aca	gga	tgg	ggg	cgg	ctg	tca	aga	aaa	ctg	atc	aat	ggg	atc	cga	gac	2016
Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys	Leu	Ile	Asn	Gly	Ile	Arg	Asp	
				660				665					670			
aag	cag	agt	gga	aag	aca	atc	ctg	gat	ttt	ctt	aag	tcc	gat	gga	ttt	2064
Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp	Phe	Leu	Lys	Ser	Asp	Gly	Phe	
		675				680							685			
gcc	aac	cgg	aac	ttc	atg	cag	ttg	atc	cat	gat	gac	tct	ctc	acc	ttt	2112
Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile	His	Asp	Asp	Ser	Leu	Thr	Phe	
	690					695					700					
aag	gag	gac	atc	cag	aaa	gca	caa	ggt	tct	ggc	cag	ggg	gac	agt	ctt	2160
Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val	Ser	Gly	Gln	Gly	Asp	Ser	Leu	
	705				710					715					720	
cac	gag	cac	atc	gct	aat	ctt	gca	ggt	agc	cca	gct	atc	aaa	aag	gga	2208
His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly	Ser	Pro	Ala	Ile	Lys	Lys	Gly	
				725					730					735		
ata	ctg	cag	acc	ggt	aag	gtc	gtg	gat	gaa	ctc	gtc	aaa	gta	atg	gga	2256
Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp	Glu	Leu	Val	Lys	Val	Met	Gly	
			740					745					750			
agg	cat	aag	ccc	gag	aat	atc	ggt	atc	gag	atg	gcc	cga	gag	aac	caa	2304
Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile	Glu	Met	Ala	Arg	Glu	Asn	Gln	
		755					760						765			
act	acc	cag	aag	gga	cag	aag	aac	agt	agg	gaa	agg	atg	aag	agg	att	2352
Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser	Arg	Glu	Arg	Met	Lys	Arg	Ile	
		770				775						780				
gaa	gag	ggt	ata	aaa	gaa	ctg	ggg	tcc	caa	atc	ctt	aag	gaa	cac	cca	2400
Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser	Gln	Ile	Leu	Lys	Glu	His	Pro	
		785			790					795					800	
ggt	gaa	aac	acc	cag	ctt	cag	aat	gag	aag	ctc	tac	ctg	tac	tac	ctg	2448
Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu	Lys	Leu	Tyr	Leu	Tyr	Tyr	Leu	
				805					810					815		
cag	aac	ggc	agg	gac	atg	tac	gtg	gat	cag	gaa	ctg	gac	atc	aat	cgg	2496
Gln	Asn	Gly	Arg	Asp	Met	Tyr	Val	Asp	Gln	Glu	Leu	Asp	Ile	Asn	Arg	
				820				825					830			
ctc	tcc	gac	tac	gac	gtg	gat	cat	atc	gtg	ccc	cag	tct	ttt	ctc	aaa	2544
Leu	Ser	Asp	Tyr	Asp	Val	Asp	His	Ile	Val	Pro	Gln	Ser	Phe	Leu	Lys	
		835					840					845				
gat	gat	tct	att	gat	aat	aaa	gtg	ttg	aca	aga	tcc	gat	aaa	aat	aga	2592
Asp	Asp	Ser	Ile	Asp	Asn	Lys	Val	Leu	Thr	Arg	Ser	Asp	Lys	Asn	Arg	
				850		855						860				
ggg	aag	agt	gat	aac	gtc	ccc	tca	gaa	gaa	ggt	gtc	aag	aaa	atg	aaa	2640
Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu	Glu	Val	Val	Lys	Lys	Met	Lys	
				865		870				875					880	
aat	tat	tgg	cgg	cag	ctg	ctg	aac	gcc	aaa	ctg	atc	aca	caa	cgg	aag	2688
Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg	Lys	
				885					890					895		
ttc	gat	aat	ctg	act	aag	gct	gaa	cga	ggt	ggc	ctg	tct	gat	ttg	gat	2736

-continued

Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg	Gly	Gly	Leu	Ser	Glu	Leu	Asp		
			900					905					910				
aaa	gcc	ggc	ttc	atc	aaa	agg	cag	ctt	ggt	gag	aca	cgc	cag	atc	acc	2784	
Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu	Val	Glu	Thr	Arg	Gln	Ile	Thr		
		915					920					925					
aag	cac	gtg	gcc	caa	att	ctc	gat	tca	cgc	atg	aac	acc	aag	tac	gat	2832	
Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser	Arg	Met	Asn	Thr	Lys	Tyr	Asp		
	930					935					940						
gaa	aat	gac	aaa	ctg	att	cga	gag	gtg	aaa	ggt	att	act	ctg	aag	tct	2880	
Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val	Lys	Val	Ile	Thr	Leu	Lys	Ser		
	945				950					955					960		
aag	ctg	gtc	tca	gat	ttc	aga	aag	gac	ttt	cag	ttt	tat	aag	gtg	aga	2928	
Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp	Phe	Gln	Phe	Tyr	Lys	Val	Arg		
				965					970					975			
gag	atc	aac	aat	tac	cac	cat	gcg	cat	gat	gcc	tac	ctg	aat	gca	gtg	2976	
Glu	Ile	Asn	Asn	Tyr	His	His	Ala	His	Asp	Ala	Tyr	Leu	Asn	Ala	Val		
				980			985							990			
gta	ggc	act	gca	ctt	atc	aaa	aaa	tat	ccc	aag	ctt	gaa	tct	gaa	ttt	3024	
Val	Gly	Thr	Ala	Leu	Ile	Lys	Lys	Tyr	Pro	Lys	Leu	Glu	Ser	Glu	Phe		
		995					1000					1005					
gtt	tac	gga	gac	tat	aaa	gtg	tac	gat	ggt	agg	aaa	atg	atc	gca		3069	
Val	Tyr	Gly	Asp	Tyr	Lys	Val	Tyr	Asp	Val	Arg	Lys	Met	Ile	Ala			
	1010					1015						1020					
aag	tct	gag	cag	gaa	ata	ggc	aag	gcc	acc	gct	aag	tac	ttc	ttt		3114	
Lys	Ser	Glu	Gln	Glu	Ile	Gly	Lys	Ala	Thr	Ala	Lys	Tyr	Phe	Phe			
	1025					1030						1035					
tac	agc	aat	att	atg	aat	ttt	ttc	aag	acc	gag	att	aca	ctg	gcc		3159	
Tyr	Ser	Asn	Ile	Met	Asn	Phe	Phe	Lys	Thr	Glu	Ile	Thr	Leu	Ala			
	1040					1045						1050					
aat	gga	gag	att	cgg	aag	cga	cca	ctt	atc	gaa	aca	aac	gga	gaa		3204	
Asn	Gly	Glu	Ile	Arg	Lys	Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	Glu			
	1055					1060						1065					
aca	gga	gaa	atc	gtg	tgg	gac	aag	ggt	agg	gat	ttc	gcg	aca	gtc		3249	
Thr	Gly	Glu	Ile	Val	Trp	Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	Val			
	1070					1075						1080					
cgg	aag	gtc	ctg	tcc	atg	ccg	cag	gtg	aac	atc	ggt	aaa	aag	acc		3294	
Arg	Lys	Val	Leu	Ser	Met	Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	Thr			
	1085					1090						1095					
gaa	gta	cag	acc	gga	ggc	ttc	tcc	aag	gaa	agt	atc	ctc	ccg	aaa		3339	
Glu	Val	Gln	Thr	Gly	Gly	Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	Lys			
	1100					1105						1110					
agg	aac	agc	gac	aag	ctg	atc	gca	cgc	aaa	aaa	gat	tgg	gac	ccc		3384	
Arg	Asn	Ser	Asp	Lys	Leu	Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	Pro			
	1115					1120						1125					
aag	aaa	tac	ggc	gga	ttc	gat	tct	cct	aca	gtc	gct	tac	agt	gta		3429	
Lys	Lys	Tyr	Gly	Gly	Phe	Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	Val			
	1130					1135						1140					
ctg	ggt	gtg	gcc	aaa	gtg	gag	aaa	ggg	aag	tct	aaa	aaa	ctc	aaa		3474	
Leu	Val	Val	Ala	Lys	Val	Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	Lys			
	1145					1150						1155					
agc	gtc	aag	gaa	ctg	ctg	ggc	atc	aca	atc	atg	gag	cga	tca	agc		3519	
Ser	Val	Lys	Glu	Leu	Leu	Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	Ser			
	1160					1165						1170					
ttc	gaa	aaa	aac	ccc	atc	gac	ttt	ctc	gag	gcg	aaa	gga	tat	aaa		3564	
Phe	Glu	Lys	Asn	Pro	Ile	Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	Lys			
	1175					1180						1185					
gag	gtc	aaa	aaa	gac	ctc	atc	att	aag	ctt	ccc	aag	tac	tct	ctc		3609	

-continued

Glu Val	Lys Lys Asp	Leu Ile	Ile Lys Leu Pro	Lys Tyr Ser Leu	
1190		1195		1200	
ttt gag	ctt gaa aac ggc	cgg aaa cga atg ctc	gct agt gcg ggc		3654
Phe Glu	Leu Glu Asn Gly	Arg Lys Arg Met Leu	Ala Ser Ala Gly		
1205		1210	1215		
gag ctg	cag aaa ggt aac	gag ctg gca ctg ccc	tct aaa tac gtt		3699
Glu Leu	Gln Lys Gly Asn	Glu Leu Ala Leu Pro	Ser Lys Tyr Val		
1220		1225	1230		
aat ttc	ttg tat ctg gcc	agc cac tat gaa aag	ctc aaa ggg tct		3744
Asn Phe	Leu Tyr Leu Ala	Ser His Tyr Glu Lys	Leu Lys Gly Ser		
1235		1240	1245		
ccc gaa	gat aat gag cag	aag cag ctg ttc gtg	gaa caa cac aaa		3789
Pro Glu	Asp Asn Glu Gln	Lys Gln Leu Phe Val	Glu Gln His Lys		
1250		1255	1260		
cac tac	ctt gat gag atc	atc gag caa ata agc	gaa ttc tcc aaa		3834
His Tyr	Leu Asp Glu Ile	Ile Glu Gln Ile Ser	Glu Phe Ser Lys		
1265		1270	1275		
aga gtg	atc ctc gcc gac	gct aac ctc gat aag	gtg ctt tct gct		3879
Arg Val	Ile Leu Ala Asp	Ala Asn Leu Asp Lys	Val Leu Ser Ala		
1280		1285	1290		
tac aat	aag cac agg gat	aag ccc atc agg gag	cag gca gaa aac		3924
Tyr Asn	Lys His Arg Asp	Lys Pro Ile Arg Glu	Gln Ala Glu Asn		
1295		1300	1305		
att atc	cac ttg ttt act	ctg acc aac ttg ggc	gcg cct gca gcc		3969
Ile Ile	His Leu Phe Thr	Leu Thr Asn Leu Gly	Ala Pro Ala Ala		
1310		1315	1320		
ttc aag	tac ttc gac acc	acc ata gac aga aag	cgg tac acc tct		4014
Phe Lys	Tyr Phe Asp Thr	Thr Ile Asp Arg Lys	Arg Tyr Thr Ser		
1325		1330	1335		
aca aag	gag gtc ctg gac	gcc aca ctg att cat	cag tca att acg		4059
Thr Lys	Glu Val Leu Asp	Ala Thr Leu Ile His	Gln Ser Ile Thr		
1340		1345	1350		
ggg ctc	tat gaa aca aga	atc gac ctc tct cag	ctc ggt gga gac		4104
Gly Leu	Tyr Glu Thr Arg	Ile Asp Leu Ser Gln	Leu Gly Gly Asp		
1355		1360	1365		
agc agg	gct gac				4116
Ser Arg	Ala Asp				
1370					

<210> SEQ ID NO 6
 <211> LENGTH: 1372
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 6

Met Asp Lys Lys Tyr Ser	Ile Gly Leu Asp Ile Gly Thr Asn Ser Val
1	5 10 15
Gly Trp Ala Val Ile Thr Asp	Glu Tyr Lys Val Pro Ser Lys Lys Phe
20	25 30
Lys Val Leu Gly Asn Thr Asp	Arg His Ser Ile Lys Lys Asn Leu Ile
35	40 45
Gly Ala Leu Leu Phe Asp Ser	Gly Glu Thr Ala Glu Ala Thr Arg Leu
50	55 60
Lys Arg Thr Ala Arg Arg	Arg Tyr Thr Arg Arg Lys Asn Arg Ile Cys
65	70 75 80

-continued

Tyr	Leu	Gln	Glu	Ile	Phe	Ser	Asn	Glu	Met	Ala	Lys	Val	Asp	Asp	Ser
				85					90					95	
Phe	Phe	His	Arg	Leu	Glu	Glu	Ser	Phe	Leu	Val	Glu	Glu	Asp	Lys	Lys
			100					105					110		
His	Glu	Arg	His	Pro	Ile	Phe	Gly	Asn	Ile	Val	Asp	Glu	Val	Ala	Tyr
			115				120					125			
His	Glu	Lys	Tyr	Pro	Thr	Ile	Tyr	His	Leu	Arg	Lys	Lys	Leu	Val	Asp
	130					135					140				
Ser	Thr	Asp	Lys	Ala	Asp	Leu	Arg	Leu	Ile	Tyr	Leu	Ala	Leu	Ala	His
	145				150					155					160
Met	Ile	Lys	Phe	Arg	Gly	His	Phe	Leu	Ile	Glu	Gly	Asp	Leu	Asn	Pro
				165						170				175	
Asp	Asn	Ser	Asp	Val	Asp	Lys	Leu	Phe	Ile	Gln	Leu	Val	Gln	Thr	Tyr
			180					185					190		
Asn	Gln	Leu	Phe	Glu	Glu	Asn	Pro	Ile	Asn	Ala	Ser	Gly	Val	Asp	Ala
		195					200					205			
Lys	Ala	Ile	Leu	Ser	Ala	Arg	Leu	Ser	Lys	Ser	Arg	Arg	Leu	Glu	Asn
	210					215					220				
Leu	Ile	Ala	Gln	Leu	Pro	Gly	Glu	Lys	Lys	Asn	Gly	Leu	Phe	Gly	Asn
	225				230					235					240
Leu	Ile	Ala	Leu	Ser	Leu	Gly	Leu	Thr	Pro	Asn	Phe	Lys	Ser	Asn	Phe
				245					250					255	
Asp	Leu	Ala	Glu	Asp	Ala	Lys	Leu	Gln	Leu	Ser	Lys	Asp	Thr	Tyr	Asp
			260					265					270		
Asp	Asp	Leu	Asp	Asn	Leu	Leu	Ala	Gln	Ile	Gly	Asp	Gln	Tyr	Ala	Asp
		275					280					285			
Leu	Phe	Leu	Ala	Ala	Lys	Asn	Leu	Ser	Asp	Ala	Ile	Leu	Leu	Ser	Asp
	290					295					300				
Ile	Leu	Arg	Val	Asn	Thr	Glu	Ile	Thr	Lys	Ala	Pro	Leu	Ser	Ala	Ser
	305				310					315					320
Met	Ile	Lys	Arg	Tyr	Asp	Glu	His	His	Gln	Asp	Leu	Thr	Leu	Leu	Lys
				325					330						335
Ala	Leu	Val	Arg	Gln	Gln	Leu	Pro	Glu	Lys	Tyr	Lys	Glu	Ile	Phe	Phe
			340					345					350		
Asp	Gln	Ser	Lys	Asn	Gly	Tyr	Ala	Gly	Tyr	Ile	Asp	Gly	Gly	Ala	Ser
		355					360					365			
Gln	Glu	Glu	Phe	Tyr	Lys	Phe	Ile	Lys	Pro	Ile	Leu	Glu	Lys	Met	Asp
	370					375					380				
Gly	Thr	Glu	Glu	Leu	Leu	Val	Lys	Leu	Asn	Arg	Glu	Asp	Leu	Leu	Arg
	385				390					395					400
Lys	Gln	Arg	Thr	Phe	Asp	Asn	Gly	Ser	Ile	Pro	His	Gln	Ile	His	Leu
				405					410					415	
Gly	Glu	Leu	His	Ala	Ile	Leu	Arg	Arg	Gln	Glu	Asp	Phe	Tyr	Pro	Phe
			420					425					430		
Leu	Lys	Asp	Asn	Arg	Glu	Lys	Ile	Glu	Lys	Ile	Leu	Thr	Phe	Arg	Ile
		435					440					445			
Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg	Gly	Asn	Ser	Arg	Phe	Ala	Trp
	450					455					460				
Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile	Thr	Pro	Trp	Asn	Phe	Glu	Glu
	465				470					475					480
Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln	Ser	Phe	Ile	Glu	Arg	Met	Thr

-continued

485				490				495							
Asn	Phe	Asp	Lys	Asn	Leu	Pro	Asn	Glu	Lys	Val	Leu	Pro	Lys	His	Ser
			500						505					510	
Leu	Leu	Tyr	Glu	Tyr	Phe	Thr	Val	Tyr	Asn	Glu	Leu	Thr	Lys	Val	Lys
		515					520						525		
Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro	Ala	Phe	Leu	Ser	Gly	Glu	Gln
	530					535					540				
Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe	Lys	Thr	Asn	Arg	Lys	Val	Thr
545					550					555					560
Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe	Lys	Lys	Ile	Glu	Cys	Phe	Asp
			565						570					575	
Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu	Gly
			580						585					590	
Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu	Asp
		595					600						605		
Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu	Thr
	610					615					620				
Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu	Glu	Arg	Leu	Lys	Thr	Tyr	Ala
625					630					635					640
His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys	Gln	Leu	Lys	Arg	Arg	Arg	Tyr
			645						650						655
Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys	Leu	Ile	Asn	Gly	Ile	Arg	Asp
			660						665				670		
Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp	Phe	Leu	Lys	Ser	Asp	Gly	Phe
		675					680						685		
Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile	His	Asp	Asp	Ser	Leu	Thr	Phe
	690					695					700				
Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val	Ser	Gly	Gln	Gly	Asp	Ser	Leu
705					710					715					720
His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly	Ser	Pro	Ala	Ile	Lys	Lys	Gly
			725						730						735
Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp	Glu	Leu	Val	Lys	Val	Met	Gly
			740						745				750		
Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile	Glu	Met	Ala	Arg	Glu	Asn	Gln
		755					760						765		
Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser	Arg	Glu	Arg	Met	Lys	Arg	Ile
	770					775					780				
Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser	Gln	Ile	Leu	Lys	Glu	His	Pro
785					790					795					800
Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu	Lys	Leu	Tyr	Leu	Tyr	Tyr	Leu
			805						810						815
Gln	Asn	Gly	Arg	Asp	Met	Tyr	Val	Asp	Gln	Glu	Leu	Asp	Ile	Asn	Arg
			820						825				830		
Leu	Ser	Asp	Tyr	Asp	Val	Asp	His	Ile	Val	Pro	Gln	Ser	Phe	Leu	Lys
		835					840						845		
Asp	Asp	Ser	Ile	Asp	Asn	Lys	Val	Leu	Thr	Arg	Ser	Asp	Lys	Asn	Arg
		850				855					860				
Gly	Lys	Ser	Asp	Asn	Val	Pro	Ser	Glu	Glu	Val	Val	Lys	Lys	Met	Lys
865					870					875					880
Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg	Lys
			885						890						895

-continued

Phe Asp Asn Leu Thr Lys Ala Glu Arg Gly Gly Leu Ser Glu Leu Asp
 900 905 910

Lys Ala Gly Phe Ile Lys Arg Gln Leu Val Glu Thr Arg Gln Ile Thr
 915 920 925

Lys His Val Ala Gln Ile Leu Asp Ser Arg Met Asn Thr Lys Tyr Asp
 930 935 940

Glu Asn Asp Lys Leu Ile Arg Glu Val Lys Val Ile Thr Leu Lys Ser
 945 950 955 960

Lys Leu Val Ser Asp Phe Arg Lys Asp Phe Gln Phe Tyr Lys Val Arg
 965 970 975

Glu Ile Asn Asn Tyr His His Ala His Asp Ala Tyr Leu Asn Ala Val
 980 985 990

Val Gly Thr Ala Leu Ile Lys Lys Tyr Pro Lys Leu Glu Ser Glu Phe
 995 1000 1005

Val Tyr Gly Asp Tyr Lys Val Tyr Asp Val Arg Lys Met Ile Ala
 1010 1015 1020

Lys Ser Glu Gln Glu Ile Gly Lys Ala Thr Ala Lys Tyr Phe Phe
 1025 1030 1035

Tyr Ser Asn Ile Met Asn Phe Phe Lys Thr Glu Ile Thr Leu Ala
 1040 1045 1050

Asn Gly Glu Ile Arg Lys Arg Pro Leu Ile Glu Thr Asn Gly Glu
 1055 1060 1065

Thr Gly Glu Ile Val Trp Asp Lys Gly Arg Asp Phe Ala Thr Val
 1070 1075 1080

Arg Lys Val Leu Ser Met Pro Gln Val Asn Ile Val Lys Lys Thr
 1085 1090 1095

Glu Val Gln Thr Gly Gly Phe Ser Lys Glu Ser Ile Leu Pro Lys
 1100 1105 1110

Arg Asn Ser Asp Lys Leu Ile Ala Arg Lys Lys Asp Trp Asp Pro
 1115 1120 1125

Lys Lys Tyr Gly Gly Phe Asp Ser Pro Thr Val Ala Tyr Ser Val
 1130 1135 1140

Leu Val Val Ala Lys Val Glu Lys Gly Lys Ser Lys Lys Leu Lys
 1145 1150 1155

Ser Val Lys Glu Leu Leu Gly Ile Thr Ile Met Glu Arg Ser Ser
 1160 1165 1170

Phe Glu Lys Asn Pro Ile Asp Phe Leu Glu Ala Lys Gly Tyr Lys
 1175 1180 1185

Glu Val Lys Lys Asp Leu Ile Ile Lys Leu Pro Lys Tyr Ser Leu
 1190 1195 1200

Phe Glu Leu Glu Asn Gly Arg Lys Arg Met Leu Ala Ser Ala Gly
 1205 1210 1215

Glu Leu Gln Lys Gly Asn Glu Leu Ala Leu Pro Ser Lys Tyr Val
 1220 1225 1230

Asn Phe Leu Tyr Leu Ala Ser His Tyr Glu Lys Leu Lys Gly Ser
 1235 1240 1245

Pro Glu Asp Asn Glu Gln Lys Gln Leu Phe Val Glu Gln His Lys
 1250 1255 1260

His Tyr Leu Asp Glu Ile Ile Glu Gln Ile Ser Glu Phe Ser Lys
 1265 1270 1275

-continued

Arg Val Ile Leu Ala Asp Ala Asn Leu Asp Lys Val Leu Ser Ala
 1280 1285 1290

Tyr Asn Lys His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu Asn
 1295 1300 1305

Ile Ile His Leu Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala Ala
 1310 1315 1320

Phe Lys Tyr Phe Asp Thr Thr Ile Asp Arg Lys Arg Tyr Thr Ser
 1325 1330 1335

Thr Lys Glu Val Leu Asp Ala Thr Leu Ile His Gln Ser Ile Thr
 1340 1345 1350

Gly Leu Tyr Glu Thr Arg Ile Asp Leu Ser Gln Leu Gly Gly Asp
 1355 1360 1365

Ser Arg Ala Asp
 1370

<210> SEQ ID NO 7
 <211> LENGTH: 83
 <212> TYPE: DNA
 <213> ORGANISM: Streptococcus pyogenes

<400> SEQUENCE: 7

gttttagagc tagaaatagc aagttaaaat aaggctagtc cgttatcaac ttgaaaaagt 60
 ggcaccgagt cggtggtgct ttt 83

<210> SEQ ID NO 8
 <211> LENGTH: 665
 <212> TYPE: DNA
 <213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 8

tttcaaaaat tcttactttt tttttggatg gacgcaaaga agtttaataa tcatattaca 60
 tggcattacc accatataca tatccatata catatocata tctaattctta cttatatggt 120
 gtggaaatgt aaagagcccc attatcttag cctaaaaaaa cttctctttt ggaactttca 180
 gtaatacgct taactgctca ttgctatatt gaagtacgga ttagaagccg ccgagcgggt 240
 gacagccctc cgaaggaaga ctctctctcg tgcgtctctg tcttcaccgg tgcggttctc 300
 gaaacgcaga tgtgcctcgc gccgcactgc tccgaacaat aaagattcta caatactagc 360
 ttttatggtt atgaagagga aaaattggca gtaacctggc ccacaaaacc ttcaaatgaa 420
 cgaatcaaat taacaacctat aggatgataa tgcgattagt tttttagcct tattttctggg 480
 gtaattaatc agcgaagcga tgatttttga tctattaaca gatataaaa tgcaaaaact 540
 gcataaccac ttttaactaat actttcaaca ttttcggttt gtattacttc ttattcaaat 600
 gtaataaaaag tatcaacaaa aaattgttaa tatacctcta tactttaacg tcaaggagaa 660
 aaaac 665

<210> SEQ ID NO 9
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Nuclear transition signal.
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(21)

-continued

<400> SEQUENCE: 9

ccc aag aag aag agg aag gtg 21
 Pro Lys Lys Lys Arg Lys Val
 1 5

<210> SEQ ID NO 10
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

Pro Lys Lys Lys Arg Lys Val
 1 5

<210> SEQ ID NO 11
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GS linker
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(15)

<400> SEQUENCE: 11

ggt gga gga ggt tct 15
 Gly Gly Gly Gly Ser
 1 5

<210> SEQ ID NO 12
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 12

Gly Gly Gly Gly Ser
 1 5

<210> SEQ ID NO 13
 <211> LENGTH: 66
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Flag tag
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(66)

<400> SEQUENCE: 13

gac tat aag gac cac gac gga gac tac aag gat cat gat att gat tac 48
 Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp Tyr
 1 5 10 15

aaa gac gat gac gat aag 66
 Lys Asp Asp Asp Asp Lys
 20

<210> SEQ ID NO 14
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 14

Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp Ile Asp Tyr
 1 5 10 15

Lys Asp Asp Asp Asp Lys
 20

<210> SEQ ID NO 15
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Strep-tag
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(24)

<400> SEQUENCE: 15

tgg agc cac ccg cag ttc gaa aaa 24
 Trp Ser His Pro Gln Phe Glu Lys
 1 5

<210> SEQ ID NO 16
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 16

Trp Ser His Pro Gln Phe Glu Lys
 1 5

<210> SEQ ID NO 17
 <211> LENGTH: 171
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: SH3 domain
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(171)

<400> SEQUENCE: 17

gca gag tat gtg cgg gcc ctc ttt gac ttt aat ggg aat gat gaa gaa 48
 Ala Glu Tyr Val Arg Ala Leu Phe Asp Phe Asn Gly Asn Asp Glu Glu
 1 5 10 15

gat ctt ccc ttt aag aaa gga gac atc ctg aga atc cgg gat aag cct 96
 Asp Leu Pro Phe Lys Lys Gly Asp Ile Leu Arg Ile Arg Asp Lys Pro
 20 25 30

gaa gag cag tgg tgg aat gca gag gac agc gaa gga aag agg ggg atg 144
 Glu Glu Gln Trp Trp Asn Ala Glu Asp Ser Glu Gly Lys Arg Gly Met
 35 40 45

att cct gtc cct tac gtg gag aag tat 171
 Ile Pro Val Pro Tyr Val Glu Lys Tyr
 50 55

<210> SEQ ID NO 18
 <211> LENGTH: 57
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 18

Ala Glu Tyr Val Arg Ala Leu Phe Asp Phe Asn Gly Asn Asp Glu Glu
 1 5 10 15
 Asp Leu Pro Phe Lys Lys Gly Asp Ile Leu Arg Ile Arg Asp Lys Pro
 20 25 30
 Glu Glu Gln Trp Trp Asn Ala Glu Asp Ser Glu Gly Lys Arg Gly Met
 35 40 45
 Ile Pro Val Pro Tyr Val Glu Lys Tyr
 50 55

<210> SEQ ID NO 19
 <211> LENGTH: 188
 <212> TYPE: DNA
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 19

gcgaatttct tatgatttat gatttttatt attaaataag ttataaaaaa aataagtgtta 60
 taaaaatttt aaagtgactc ttaggtttta aaacgaaaat tcttattcct gagtaactct 120
 ttctgttagg tcaggttgct ttctcaggta tagcatgagg tcgctcttat tgaccacacc 180
 tctaccgg 188

<210> SEQ ID NO 20
 <211> LENGTH: 417
 <212> TYPE: DNA
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 20

ataccaggca tggagcttat ctggccggtt cgagttttcg acgagtttgg agacattcct 60
 tatagatgtc cttttttttt aatgatattc gttaaagaac aaaaagtcaa agcagtttaa 120
 cctaacacct gttgttgatg ctacttgaaa caaggcttct aggcgaatac ttaaaaagg 180
 aatttcaata gcggtttata tatctgtttg cttttcaaga tattatgtaa acgcacgatg 240
 tttttcgccc aggctttatt tttttgttg ttggtgtcct ctgaagaat tttctegggc 300
 agatctttgt cggaatgtaa aaaagcgcgt aattaaactt tctattatgc tgactaaaat 360
 ggaagtgatc accaaaggct atttctgatt atataatcta gtcattactc gctcgag 417

<210> SEQ ID NO 21
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: SH3-binding ligand
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(33)

<400> SEQUENCE: 21

cct cca cct get ctg cca cct aag aga agg aga 33
 Pro Pro Pro Ala Leu Pro Pro Lys Arg Arg Arg
 1 5 10

<210> SEQ ID NO 22
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

```

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 29

ataacggaat ccaactgggc                20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 30

gtcaattacg aagactgaac                20

<210> SEQ ID NO 31
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 31

gtcaatagga tcccctttt                19

<210> SEQ ID NO 32
<211> LENGTH: 10126
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Plasmid carrying dCas9-PmCDA1 fusion protein
and chimeric RNA targeting galK gene of E.coli
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (5561)..(5580)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 32

atcgccattc gccattcagg ctgcgcaact gttgggaagg gcgatcggtg cgggcctctt    60
cgctattaacg ccagctggcg aaaggggggat gtgctgcaag gcgattaagt tgggtaacgc    120
cagggttttc ccagtcacga cgttgtaaaa cgacggccag tgaattcgag ctcggtaccc    180
ggccgcaaac aacagataaa acgaaaggcc cagtctttcg actgagcctt tcgttttatt    240
tgatgcctgt caagtaacag caggactctt agtgggtgtg agtattttta cctgaatcat    300
aatggacaac tcgctccgtc gtttttcagc tcgcttcaaa gtcttctcaa gccatctatt    360
ctcattcaat tgattgtgcg acgattggat gaatattttc ctgcaacatt ggtagtgttc    420
acttaccatt acattcaacc caaccccggt atctctgagg ttccacagcc caatttgatt    480
cctcgcattt ttctcgtaat agagtttgca agcccagatt ttcaaagtgt ggcggttccc    540
cgcagctcc  tgggtatacc attctaagat cttttcagcg caatctgcac aaggactcca    600
ggatgagtac caatttatcg tgaattgtcc ggggttgcg cgcaggattt cttcgacttt    660
tctaagteta aagatttcgg cgtgaatgcc acgttctgtc ccgctctgtg gtttattcac    720
agcatagccc caaaaacacg ctctacgttc accccgctgc ttttaattcaa agagaacgta    780
gcatctatgc gacacggatt ttttgtgtt gaaaaactgt ttcttaaacg tgtagatgtc    840
caacttctca tggattctca cgtactcagc gtcggtcac  ctgacttat  cgtcacgtc    900
tttgtaatca atatcatgat cctttagtgc tccgtcgtgg tccttatagt ctccggactc    960

```

-continued

gagcctagac ttatcgtcat cgtctttgta atcaatatca tgatccttgt agtctccgtc	1020
gtggtoctta tagtctccgg aatacttctc cacgtaaggg acaggaatca tccccctctt	1080
tccttcgctg tcctctgcat tccaccactg ctccctcaggc ttatcccgga ttctcaggat	1140
gtctcctttc ttaaagggaa gatcctcttc atcattccca ttaaagtcaa agagggctcg	1200
cacatactca gcagaacctc cacctccaga acctcctcca ccgtcacctc ctagtctgact	1260
caaatcaatg cgtgtttcat aaagaccagt gatggattga tggataagag tggcatctaa	1320
aacttccttt gtagacgtat atcgtttacg atcaattggt gtatcaaaat atttaaaagc	1380
agcgggagct ccaagattcg tcaacgtaaa taaatgaata atattttctg cttgttcacg	1440
tattggtttg tctctatggt tgttatatgc actaagaact ttatctaaat tggcatctgc	1500
taaaataaca cgcttagaaa attcactgat ttgctcaata atctcatcta aataatgctt	1560
atgctgctcc acaacaatt gtttttggtc gttatcttct ggactaccct tcaacttttc	1620
ataatgacta gctaaatata aaaaattcac atatttgctt ggcagagcca gctcatttcc	1680
tttttgtaat tctccggcac tagccagcat ccgtttacga ccgttttcta actcaaaaag	1740
actatattta ggtagttaa tgattaagtc ttttttaact tccttatatc ctttagcttc	1800
taaaaagtea atcggatttt tttcaagga acttctttcc ataattgtga tccctagtaa	1860
ctctttaacg gattttaact tcttcgattt cctcttttcc accttagcaa ccaactaggac	1920
tgaataagct accggtggac tatcaaaacc acctatattt tttggatccc agtctttttt	1980
acgagcaata agcttgcgac aatttctttt tggtaaaatt gactccttgg agaatccgcc	2040
tgtctgtact tctgttttct tgacaatatt gacttggggc atggacaata ctttgcgcac	2100
tgtggcaaaa tctcgcctt tatcccagac aatttctcca gtttcccct tagtttcgat	2160
tagagggcgt ttgcgaatct ctccatttgc aagtgttaatt tctgttttga agaagttcat	2220
gatattagag taaaagaaat attttgoggt tgctttgcct atttcttgcct cagacttagc	2280
aatcatttta cgaacatcat aaactttata atcaccatag acaaactccg attcaagttt	2340
tggatatttc ttaatcaaaag cagttccaac gacggcattt agatacgcac catgggcatg	2400
atggtaattg ttaatctcac gtactttata gaattggaaa tcttttcgga agtcagaaac	2460
taatttagat ttttaaggtaa tcactttaac ctctcgaata agtttatcat tttcatcgta	2520
tttagtattc atcgcactat ccaaaatttg tgccacatgc ttagtgattt ggcgagtttc	2580
aaccaattgg cgtttgataa aaccagcttt atcaagttca ctcaaacctc cacgttcagc	2640
tttcgttaaa ttatcaaaact tacggtgagt gattaacttg gcgtttagaa gttgtctcca	2700
atagtttttc atctttttga ctactcttc acttggaaag ttatccgatt taccacgatt	2760
tttatcagaa cgcgttaaga ccttattgtc tattgaatcg tctttaagga aactttgtgg	2820
aacaatggca tcgacatcat aatcacttaa acgattaata tctaattctt ggtccacata	2880
catgtctctt ccatttttga gataatagag atagagcttt tcatttttga attgagtatt	2940
ttcaacagga tgetctttaa gaatctgact tccctaattct ttgatacctt cttcgattcg	3000
tttcatacgc tctcgcgaat ttttctggcc cttttgagtt gtctgatttt cacgtgccat	3060
ttcaataacg atattttctg gcttatgccc ccccatctact ttgaccaatt catcaacaac	3120
ttttacagtc tgtaaaatac cttttttaat agcagggcta ccagctaaat ttgcaatatg	3180
ttcatgtaaa ctatcgcctt gtccagacac ttgtgctttt tgaatgtctt ctttaaatgt	3240

-continued

caaacatca	tcattgatca	gctgcataaa	attgctgattg	gcaaaacat	ctgatttcaa	3300
aaaatcfaat	attgttttgc	cagattgctt	atccctaata	ccattaatca	attttcgaga	3360
caaacgtccc	caaccagtat	aacggcgacg	tttaagctgt	ttcatcacct	tatcatcaaa	3420
gaggtgagca	tatgttttaa	gtctttctc	aatcatctcc	ctatcttcaa	ataaggtcaa	3480
tgtaaaaaca	atatoctcta	agatatcttc	atctttctca	ttatocaaaa	aatctttatc	3540
tttaataaatt	tttagcaaat	catggttagt	acctaataaa	gcattaaatc	tatcttcaac	3600
tcctgaaatt	tcaaacctat	caaacattc	tatttttttg	aaataatctt	cttttaattg	3660
cttaacgggt	acttttcgat	ttgttttgaa	gagtaaatca	acaatggctt	tctttctgtc	3720
acctgaaaga	aatgctgggt	ttgcattcc	ttcagtaaca	tatttgacct	ttgtcaattc	3780
gttataaaoc	gtaaaatact	cataaagcaa	actatgtttt	gtagtactt	ttcatttgg	3840
aagattttta	tcaagtttg	tcattgcttc	aataaatgat	tgagctgaag	cacctttatc	3900
gacaactctc	tcaaaattcc	atggggtaat	tgctttctca	gacttccgag	tcattccatg	3960
aaaacgacta	ttgccacgag	ccaatggacc	aacataataa	ggaattcgaa	aagtcaagat	4020
tttttcaatc	ttctcagcat	tgctttttaa	aaatggataa	aagcttctct	gtcttctcaa	4080
aatagcatgc	agctcaccca	agtgaatttg	atggggaata	gagcctgtgt	caaaggtccg	4140
ttgcttgccg	agcaaatctt	cacgatttag	ttccaccaat	aattctcag	taccatccat	4200
tttttctaaa	attggtttga	taaatttata	aaattctctc	tggttagctc	ccccatcaat	4260
ataacctgca	tatcctgttt	ttgattgatc	aaaaaagatt	tctttatact	tttctggaag	4320
ttgttgctga	actaaagctt	ttaaagagt	caagtcttga	tgatgttcat	cgtagcgttt	4380
aatcattgaa	gctgataggg	gagccttagt	tatttcagta	ttactctta	ggatattctga	4440
aagtaaaata	gcattctgata	aattcttagc	tgccaaaaac	aaatcagcat	attgatctcc	4500
aatttgccgc	aataaattat	ctaaatcctc	atcgttaagta	tcttttgaaa	gctgtaattt	4560
agcattctct	gccaaatcaa	aatttgattt	aaaattaggg	gtcaaaccca	atgacaaagc	4620
aatgagattc	ccaaataagc	catttttctc	ctcacggggg	agctgagcaa	tgagattttc	4680
taatcgtctt	gatttactca	atcgtgcaga	aagaatcgtc	ttagcatcta	ctccacttgc	4740
gttaaataggg	ttttcttcaa	ataattgatt	gtaggtttgt	accaactgga	taaatagttt	4800
gtccacatca	ctattatcag	gatttaaatc	tcctcctaatc	aaaaaatgac	cacgaaactt	4860
aatcatatgc	gctaaggcca	aatagattaa	gcgcaaatcc	gctttatcag	tagaatctac	4920
caattttttt	cgcagatgat	agatagttgg	atatttctca	tgataagcaa	cttcatctac	4980
tatatttcca	aaaataggat	gacgttcatg	cttcttctct	tcttccacca	aaaaagactc	5040
ttcaagctga	tgaagaaac	tatcatctac	tttcgccatc	tcatttgaaa	aaatctctctg	5100
tagataacaa	atacattctc	tcgcagctgt	ataccttcta	cgagctgtcc	gtttgagacg	5160
agtcgcttcc	gctgtctctc	cactgtcaaa	taaaagagcc	cctataagat	tttttttgat	5220
actgtggcgg	tctgtatttc	ccagaacctt	gaacttttta	gacggaacct	tatattctac	5280
agtgatcacc	gccatccoga	cgctatttgg	gccgatagct	aagcctattg	agtatttctt	5340
atccattttt	gcctcctaaa	atgggcccct	taaattaaat	ccataatgag	tttgatgatt	5400
tcaataatag	ttttaatgac	ctccgaaatt	agtttaatat	gctttaattt	ttctttttca	5460
aaatatctct	tcaaaaaata	ttaccaata	cttaataata	aatagattat	aacacaaat	5520

-continued

tcttttgaca agtagtttat tttgttataa ttctatagta nnnnnnnnnn nnnnnnnnnn 5580
gttttagagc tagaaatagc aagttaaaat aaggctagtc cgttatcaac ttgaaaaagt 5640
ggcaccgagt cgggtcgtttt tttgatactt ctattctact ctgactgcaa accaaaaaaa 5700
caagcgcttt caaaacgctt gttttatcat ttttagggaa attaatctct taatcctttt 5760
atcattctac atttaggcgc tgccatcttg ctaaaacctac taagctccac aggatgattt 5820
cgtaatcccg caagaggccc ggcagtaccg gcataaccaa gcctatgctt acagcatcca 5880
gggtgacggt gccgaggatg acgatgagcg cattgttaga tttcatacac ggtgcctgac 5940
tgcgttagca atttaactgt gataaactac cgcattaaag cttatcgatg ataagctgtc 6000
aaacatgaga attacaactt atatcgtatg gggctgactt caggtgctac atttgaagag 6060
ataaattgca ctgaaatcta gtcggatcct cgctcactga ctcgctgcgc tcggtcgttc 6120
ggctgcggcg agcgggatca gctcactcaa aggcggtaat acggttatcc acagaatcag 6180
gggataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 6240
agcccgctt gctggcgctt ttccatagtc tccgcccccc tgacgagcat cacaaaaatc 6300
gacgctcaag tcagaggtgg cgaaacccga caggactata aagataccag gcgtttcccc 6360
ctggaagetc cctcgtgcgc tctcctgttc cgaccctgcc gcttaccgga tacctgtccg 6420
cctttctccc ttcgggaagc gtggcgcttt ctcatagctc acgctgtagg tatctcagtt 6480
cgggttaggt cgttcctccc aagctgggct gtgtgcaaga accccccgtt cagccccgacc 6540
gctgcgcctt atccggtaac tatcgtcttg agtccaacct ggtaagacac gacttatcgc 6600
cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag 6660
agttctttaa gtgggtggct aactacggct aactagaag gacagtattt ggtatctgcg 6720
ctctgctgaa gccagttacc ttcggaaaaa gagttgtag ctcttgatcc ggcaaaaaa 6780
ccaccgctgg tagcgggtgt ttttttgtt gcaagcagca gattacgcgc agaaaaaag 6840
gatctcaaga agatcctttg atcttttcta cggggtctga cgctcagtg aacgaaaact 6900
cacgttaagg gattttggc atgagattat caaaaaggat cttcacctag atccttttaa 6960
atataaatg aagttttaa tcaatctaaa gtatatatga gtaacttgg tctgacagtt 7020
accaatgctt aatcagtgag gcacctatct cagcgatctg tctatctcgt tcatccatag 7080
ttgctgact ccccgctgtg tagataacta cgatacggga gggcttacca tctggcccca 7140
gtgctgcaat gataccgca gaaccacgct caccgctccc agatttatca gcaataaacc 7200
agccagcccg aagggccgag cgcagaagtg gtccctgcaac tttatccgcc tccatccagt 7260
ctattaattg ttgocgggaa gctagagtaa gtggttcgcc agttaatagt ttgocgaacg 7320
ttggtgcaat tgctgcaggc atcgtgggtg cacgctcgtc gtttggtagt gcttcattca 7380
gctccggttc ccaacgatca agcaggtta catgatcccc catgttctgc aaaaaagcgg 7440
ttagctcctt cggctcctcg atcgttgta gaagtaagtt ggccgagtg ttatcactca 7500
tggttatggc agcactgcat aattctctta ctgctatgcc atccgtaaga tgcttttctg 7560
tgactggtga gtactcaacc aagtcattct gagaatagtg tatgocggca ccgagttgct 7620
cttgcocggc gtcaacacgg gataataccg cgccacatag cagaacttta aaagtgetca 7680
tcattgaaa acgttctctg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca 7740
gttcgatgta acccaactcg gcaccaact gatcttcagc atcttttact ttcaccagcg 7800

-continued

tttctgggtg agcaaaaaa ggaaggcaaa atgccgcaaa aaagggata agggcgacac	7860
ggaaatgttg aatactcata ctcttccttt ttcaatatta ttgaagcatt taccaggggt	7920
attgtctcat gagcggatag atatttgaat gtatttagaa aaataaaca ataggggttc	7980
cgcgacatt tccccgaaa gtgccactg acgtcaatgc cgagcgaag cgagccgaag	8040
ggtagcattt acgtagata acccctgat atgctccgac gctttatata gaaaagaaga	8100
ttcaactagg taaaacttta atataggtg agatgataag gtttataagg aattgtttg	8160
ttctaatttt tcaactcatt tgttctaatt tcttttaaca aatgttcttt tttttttaga	8220
acagttaga tatagttaga atagtttaaa ataaggagtg agaaaaagat gaaagaaga	8280
tatggaacag tctataaagg ctctcagagg ctcatagacg aagaaagtgg agaagcata	8340
gaggtagaca agttataccg taacaaacg tctgtaact tcgtaaaggc atatatagt	8400
caattaataa gtatgtaga tatgattggc ggaaaaaac ttaaatcgt taactatc	8460
ctagataatg tccacttaag taacaataca atgatagcta caacaagaga aatagcaaaa	8520
gctacaggaa caagtctaca aacagtaata acaacactta aaatcttaga agaagaaat	8580
attataaaaa gaaaaactgg agtattaatg ttaaacctg aactactaat gagaggcgac	8640
gacaaaaac aaaaaactc cttactcgaa tttgggaact ttgagcaaga ggcaaatgaa	8700
atagattgac ctccaataa caccacgtag ttattgggag gtcaatctat gaaatcgat	8760
taagcttttt ctaattcaca taagcgtgca ggtttaagt acataaaaa tataatgaaa	8820
aaaagcatca ttatactaac gttataccaa cattatactc tcattatact aattgcttat	8880
tccaatttc tattggttg aaccaacagg cgttagtggt ttgttgagtt ggtacttca	8940
tgggattaat cccatgaaac cccaaccaa ctgcgcaag ctttggttaa cacacacgc	9000
attccaacca atagttttct cggcataaag ccatgctctg acgttaaat gcactaatgc	9060
cttaaaaaa cattaaagtc taacacacta gacttattta ctctgtaatt aagtcgttaa	9120
accgtgtgct ctacgaccaa aagtataaaa cctttaagaa cttcttttt tcttgtaaaa	9180
aaagaaacta gataaatctc tcatatcttt tattcaataa tcgcatcaga ttgcagtata	9240
aatttaacga tcaactcatca tgttcatatt taccagagct cgtgctataa ttatactaat	9300
tttataagga gaaaaaata aagagggtta taatgaacga gaaaaatata aaacacagtc	9360
aaaactttat tacttcaaaa cataatatag ataaaaaat gacaaatata agattaatg	9420
aacatgataa tatctttgaa atcggctcag gaaaaggca tttaccctt gaattagtag	9480
agaggtgtaa tttcgtaact gccattgaaa tagaccataa attatgcaa actacagaaa	9540
ataaacttgt tgatcacgat aatttccaag ttttaacaa ggatatattg cagttaaat	9600
ttcctaaaaa ccaatcctat aaaatattg gtaatatacc ttataacata agtacggata	9660
taatacgcga aattgttttt gatagtagat ctgatgagat ttatttaac gtggaatac	9720
ggtttgctaa aagattatta aatacaaac gctcattggc attattttta atggcagaag	9780
ttgatatttc tatattaagt atggttccaa gagaatattt tcatcctaaa cctaaagtga	9840
atagctcact taccagatta aatagaaaa aatcaagaat atcacacaaa gataaacaga	9900
agtataatta tttcgttatg aaatgggtta acaagaata caagaaata tttaaaaaa	9960
atcaatttaa caattcctta aaacatgcag gaattgacga tttaaacaat attagctttg	10020
aacaattctt atctcttttc aatagctata aattatttaa taagtaagtt aaggatgca	10080

-continued

taaactgcat cccttaactt gttttctgtg tacctatatt ttgtga	10126
<p><210> SEQ ID NO 33 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Escherichia coli</p>	
<400> SEQUENCE: 33	
tcaatgggct aactacgttc	20
<p><210> SEQ ID NO 34 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Escherichia coli</p>	
<400> SEQUENCE: 34	
ggtccataaa ctgagacagc	20
<p><210> SEQ ID NO 35 <211> LENGTH: 223 <212> TYPE: DNA <213> ORGANISM: Escherichia coli</p>	
<400> SEQUENCE: 35	
gagttataca cagggtctggg atctattctt tttatcttt tttattctt ctttattcta	60
taaattataa ccacttgaat ataacaaaa aaaacacaca aaggtctagc ggaatttaca	120
gagggtctag cagaatttac aagttttcca gcaaaggctc agcagaattt acagataccc	180
acaactcaaa ggaaaaggac tagtaattat cattgactag ccc	223
<p><210> SEQ ID NO 36 <211> LENGTH: 951 <212> TYPE: DNA <213> ORGANISM: Escherichia coli</p>	
<400> SEQUENCE: 36	
atgtctgaat tagttgtttt caaagcaaat gaactagcga ttagtcgcta tgacttaacg	60
gagcatgaaa ccaagctaatt tttatgctgt gtggcactac tcaaccccac gattgaaaac	120
cctacaatga aagaacggac ggtatcgctc acttataacc aatacgttca gatgatgaac	180
atcagtaggg aaaatgctta tgggtgatta gctaaagcaa ccagagagct gatgacgaga	240
actgtggaat tcaggaatcc tttgggttaa ggctttgaga tttccagtg gacaaactat	300
gccaagttct caagcgaaaa attagaatta gtttttagtg aagagatatt gccttatctt	360
ttccagttaa aaaaattcat aaaatataat ctggaacatg ttaagtcttt tgaaaaacaa	420
tactctatga ggatttatga gtggttatta aaagaactaa cacaaaagaa aactcacaag	480
gcaaatatag agattagcct tgatgaattt aagttcatgt taatgcttga aaataactac	540
catgagttta aaaggcttaa ccaatggggt ttgaaaccaa taagtaaga tttaaacact	600
tacagcaata tgaaattggt ggttgataag cgaggccgcc cgactgatac gttgattttc	660
caagttgaac tagatagaca aatggatctc gtaaccgaac ttgagaacaa ccagataaaa	720
atgaatggtg acaaaatacc aacaaccatt acatcagatt cctacctaca taacggacta	780
agaaaaacac tacacgatgc ttttaactgca aaaattcagc tcaccagttt tgaggcaaaa	840
tttttgagtg acatgcaaag taagcatgat ctcaatgggt cgttctcatg gctcagcaca	900

-continued

aaacaacgaa ccacactaga gaacatactg gctaaatcag gaaggatctg a 951

<210> SEQ ID NO 37

<211> LENGTH: 714

<212> TYPE: DNA

<213> ORGANISM: Bacteriophage lambda

<400> SEQUENCE: 37

```

tcagccaaac gtctcttcag gccactgact agcgataact ttccccacaa cggaacaact    60
ctcattgcat gggatcattg ggtactgtgg gtttagtggt tgtaaaaaa cctgaccgct    120
atccctgac agtttcttga aggtaaactc atcaccccca agtctggcta tgcagaaatc    180
acctggctca acagcctgct cagggtaaac gagaattaac attccgtcag gaaagcttgg    240
cttgagcct gttggtgogg tcatggaatt accttcaacc tcaagccaga atgcagaatc    300
actggctttt ttggttggc ttcccatct ctccgcatca cctttggtaa aggttctaa    360
cttaggtgag aacatccctg cctgaacatg agaaaaaaca gggtaactcat actcacttct    420
aagtgacggc tgcatactaa ccgcttcata catctcgtag atttctctgg cgattgaagg    480
gctaaattct tcaacgctaa ctttgagaat ttttgtaagc aatgcccgt tataagcatt    540
taatgcattg atgccattaa ataagcacc aacgcctgac tgcgccatcc ceatcttgtc    600
tgcgacagat tcctgggata agccaagttc attttcttt ttttcataaa ttgctttaag    660
gcgacgtgcg tcctcaagct gctcctgtgt taatggtttc tttttgtgc tcat      714

```

<210> SEQ ID NO 38

<211> LENGTH: 5097

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: dCas9-PmCDA1

<400> SEQUENCE: 38

```

atggataaga aatactcaat aggccttagct atcggcacia atagcgtcgg atgggcggtg    60
atcactgatg aatataaggt tccgtctaaa aagttcaagg ttctgggaaa tacagaccgc    120
cacagtatca aaaaaatct tataggggct cttttatttg acagtggaga gacagcggaa    180
gcgactcgtc tcaaacggac agctcgtaga aggtatacac gtcggaagaa tcgtatttgt    240
tatctacagg agatttttcc aaatgagatg gcgaaagtag atgatagttt ctttcatcga    300
cttgaagagt ctttttgggt ggaagaagac aagaagcatg aacgtcatcc tatttttggg    360
aatatagtag atgaagttgc ttatcatgag aaatatccaa ctatctatca tctgcaaaaa    420
aaattggtag attctactga taaagcggat ttgctcttaa tctatttggc cttagcgcac    480
atgattaagt ttcgtggtca ttttttgatt gagggagatt taaatcctga taatagtgat    540
gtggacaaac tatttatcca gttggtacaa acctacaatc aattatttga agaaaaccct    600
attaacgcaa gtggagtaga tgctaaagcg attctttctg cagcattgag taaatcaaga    660
cgattagaaa atctcattgc tcagctcccc ggtgagaaga aaaatggctt atttgggaat    720
ctcattgctt tgtcattggg tttgaccctc aattttaaat caaattttga tttggcagaa    780
gatgctaaat tacagctttc aaaagatact tacgatgatg atttagataa tttattggcg    840
caaattggag atcaatatgc tgatttggtt ttggcagcta agaatttatc agatgctatt    900
ttactttcag atatcctaag agtaaaact gaaataacta aggcctccct atcagcttca    960

```

-continued

atgattaaac	gctacgatga	acatcatcaa	gacttgactc	ttttaaagc	tttagttcga	1020
caacaacttc	cagaaaagta	taaagaaatc	ttttttgatc	aatcaaaaa	cggatatgca	1080
ggttatattg	atgggggagc	tagccaagaa	gaattttata	aatttatcaa	accaatttta	1140
gaaaaaatgg	atggtactga	ggaattattg	gtgaaactaa	atcgtgaaga	ttgctgcgc	1200
aagcaacgga	cctttgacaa	cggctctatt	ccccatcaa	ttcacttggg	tgagctgcat	1260
gctattttga	gaagacaaga	agacttttat	ccatttttaa	aagacaatcg	tgagaagatt	1320
gaaaaaatct	tgacttttcg	aattccttat	tatgttggtc	cattggcgcg	tggcaatagt	1380
cgttttgcat	ggatgactcg	gaagtctgaa	gaaacaatta	ccccatgga	ttttgaagaa	1440
gttctcgata	aagggtcttc	agctcaatca	tttattgaac	gcatgacaaa	ctttgataaa	1500
aatcttccaa	atgaaaaagt	actaccaaaa	catagtttgc	tttatgagta	tttacggtt	1560
tataacgaat	tgacaaaagt	caaatatggt	actgaaggaa	tgcgaaaacc	agcatttctt	1620
tcagggtgaa	agaagaaagc	cattgttgat	ttactcttca	aaacaaatcg	aaaagtaacc	1680
gttaagcaat	taaaagaaga	ttatttcaaa	aaaatagaat	gtttgatag	tgttgaatt	1740
tcaggagtgt	aagatagatt	taatgcttca	ttaggtacct	acatgattt	gctaaaaatt	1800
attaagata	aagatttttt	ggataatgaa	gaaaatgaag	atatcttaga	ggatattggt	1860
ttaacattga	ccttatttga	agataggag	atgattgagg	aaagacttaa	aacatagct	1920
cacctctttg	atgataaggt	gatgaaacag	cttaaacctc	gccgttatac	tggttgggga	1980
cgtttgtctc	gaaaattgat	taatggtatt	agggataaag	aatctggcaa	aacaatatta	2040
gattttttga	aatcagatgg	ttttgccaat	cgcaatttta	tgcagctgat	ccatgatgat	2100
agtttgacat	ttaaagaaga	cattcaaaaa	gcacaagtgt	ctggacaagg	cgatagttta	2160
catgaacata	ttgcaaat	agctggtagc	cctgctatta	aaaaaggat	tttacagact	2220
gtaaaagttg	ttgatgaatt	ggtcaaagta	atggggcggc	ataagccaga	aaatcgtt	2280
attgaaatgg	cacgtgaaaa	tcagacaact	caaaagggcc	agaaaaatc	gagagagcgt	2340
atgaaacgaa	tcgaagaagg	tatcaaagaa	ttaggaagtc	agattcttaa	agagcatcct	2400
gttgaata	ctcaattgca	aaatgaaaag	ctctatctct	attatctcca	aaatggaaga	2460
gacatgtatg	tggaccaaga	attagatatt	aatcgtttaa	gtgattatga	tgtcogatgcc	2520
attgttccac	aaagtttct	taaagaogat	tcaatagaca	ataaggctct	aacgcgttct	2580
gataaaaatc	gtggtaaatc	ggataacggt	ccaagtgaag	aagttagtcaa	aaagatgaaa	2640
aactattgga	gacaacttct	aaacgccaag	ttaatcactc	aacgtaagtt	tgataattta	2700
acgaaagctg	aacgtggagg	tttgagtga	cttgataaag	ctggttttat	caaacgcca	2760
ttggttga	ctcgccaaat	cactaagcat	gtggcacaaa	ttttggatag	tcgcatgaat	2820
actaaatcag	atgaaaatga	taaacttatt	cgagaggtta	aagtgattac	cttaaatct	2880
aaattagttt	ctgacttccg	aaaagatttc	caattctata	aagtacgtga	gattaacaat	2940
taccatcatg	cccatgatgc	gtatctaaat	gccgtcgttg	gaactgcttt	gattaagaaa	3000
tatccaaaac	ttgaatcggg	gtttgtctat	ggtgattata	aagtttatga	tgttcgtaaa	3060
atgattgcta	agtctgagca	agaaatagcc	aaagcaaccg	caaaatattt	cttttactct	3120
aatatcatga	acttcttcaa	aacagaaatt	acacttgcaa	atggagagat	tcgcaaacgc	3180
cctctaactg	aaactaatgg	ggaaactgga	gaaattgtct	gggataaagg	gagatattt	3240

-continued

```

gccacagtgc gcaaagtatt gtccatgccc caagtcaata ttgtcaagaa aacagaagta 3300
cagacaggcg gattctccaa ggagtcaatt ttacaaaaaa gaaattcggg caagcttatt 3360
gctcgtaaaa aagactggga tccaaaaaaa tatggtgggt ttgatagtcc aacggtagct 3420
tattcagtcc tagtgggtgc taagtgga aaagggaaat cgaagaagtt aaaatccgtt 3480
aaagagttac tagggatcac aattatggaa agaagttcct ttgaaaaaaa tccgattgac 3540
tttttagaag ctaaaggata taaggaagtt aaaaagact taatcattaa actacctaaa 3600
tatagtcttt ttgagttaga aaacggtcgt aaacggatgc tggctagtgc cggagaatta 3660
caaaaaggaa atgagctggc tctgccaaac aaatatgtga atttttata tttagctagt 3720
cattatgaaa agttgaagg tagtccagaa gataacgaac aaaaaaatt gtttgggag 3780
cagcataagc attattaga tgagattatt gagcaaatca gtgaattttc taagcgtgtt 3840
attttagcag atgccaat t agataaagtt cttagtgc ataaacaa tagagacaaa 3900
ccaatacgtg aacaagcaga aaatattatt catttattta cgttgacgaa tcttgagct 3960
cccgtgctt ttaaataatt tgatacaaca attgatcgta aacgatatac gtctacaaaa 4020
gaagttttag atgccactct tatccatcaa tccatcactg gtctttatga aacacgcatt 4080
gatttgagtc agctaggagg tgacgggtga ggaggttctg gaggtggagg ttctgctgag 4140
tatgtgcgag ccctcttga ctttaatggg aatgatgaag aggatcttcc cttaagaaa 4200
ggagacatcc tgagaatccg ggataagcct gaggagcagt ggtggaatgc agaggacagc 4260
gaaggaaaaga gggggatgat tctgtccct tacgtggaga agtattccgg agactataag 4320
gaccacgacg gagactacaa ggatcatgat attgattaca aagacgatga cgataagtct 4380
aggctcagat ccggagacta taaggaccac gacggagact acaaggatca tgatattgat 4440
tacaagaagc atgacgataa gtctaggatg accgacgctg agtacgtgag aatccatgag 4500
aagttggaca tctacacgtt taagaacag tttttcaaca acaaaaaatc cgtgtcgc at 4560
agatgctacg ttctcttga attaaaacga cggggtgaac gtagagcgtg tttttggggc 4620
tatgtctgta ataaaccaca gagcgggaca gaacgtggca ttcacgccga aatctttage 4680
attagaaaag tcgaagaata cctgcgcgac aaccccgac aattcagat aaattggtac 4740
tcatcctgga gtccttgc agattgcgct gaaaagatct tagaatgta taaccaggag 4800
ctgcggggga acggccacac tttgaaaatc tgggcttgc aactctatta cgagaaaaat 4860
gcgaggaatc aaattgggct gtggaacctc agagataacg gggttgggtt gaatgtaatg 4920
gtaagtgaac actaccaatg ttgcaggaaa atattcatcc aatcgtcgca caatcaattg 4980
aatgagaata gatggcttga gaagactttg aagcgagctg aaaaacgacg gagcgagttg 5040
tccattatga ttcaggtaaa aatactccac accactaaga gtccgtctgt tacttga 5097

```

<210> SEQ ID NO 39

<211> LENGTH: 105

<212> TYPE: DNA

<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 39

```

acgttaaact taccaccgca agggataaat atctaaccac gtgcgtgttg actatcttac 60
ctctggcggg gataatgggt gcagggccca ttttaggagg caaaa 105

```

<210> SEQ ID NO 40

-continued

```

<211> LENGTH: 247
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gRNA

<400> SEQUENCE: 40
ggttttagcaa gatggcagcg cctaaatgta gaatgataaa aggattaaga gattaatttc      60
cctaaaaaatg ataaaaacaag cgttttgaaa gcgcttgttt ttttggtttg cagtcagagt      120
agaatagaag tatcaaaaaa agcaccgact cggtgccact ttttcaagtt gataacggac      180
tagccttatt ttaacttgct atttctagct ctaaaactga gaccatcccg ggtctctact      240
gcagaat                                           247

<210> SEQ ID NO 41
<211> LENGTH: 64
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 41
tatcacgcc agtgggtatt atgtcaacac cgccagagat aatttatcac cgcagatggt      60
tate                                           64

<210> SEQ ID NO 42
<211> LENGTH: 10867
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Plasmid

<400> SEQUENCE: 42
gtcggaaactg actaaagtag tgagttatac acagggctgg gatctattct ttttatcttt      60
ttttattctt tctttattct ataaattata accacttgaa tataaacaaa aaaaacacac      120
aaaggtctag cggaatttac agagggctca gcagaattta caagttttcc agcaaaggtc      180
tagcagaatt tacagatacc cacaactcaa aggaaaagga ctagtaatta tcattgacta      240
gcccatctca attggtatag tgattaaaat cacctagacc aattgagatg tatgtctgaa      300
ttagttgttt tcaaaagaaa tgaactagcg attagtcgct atgacttaac ggagcatgaa      360
accaagctaa ttttatgctg tgtggcacta ctcaacccca cgattgaaaa ccctacaatg      420
aaagaacgga cggtatcggt cacttataac caatacgttc agatgatgaa catcagtagg      480
gaaaatgctt atgggtgatt agctaaagca accagagagc tgatgacgag aactgtggaa      540
atcaggaatc ctttggttaa aggccttgag attttccagt ggacaaaacta tgccaagttc      600
tcaagcgaaa aattagaatt agtttttagt gaagagatat tgccttatct tttccagtta      660
aaaaaattca taaaataata tctggaacat gttaagtctt ttgaaaacaa atactctatg      720
aggatttatg agtgggtatt aaaagaacta acacaaaaga aaactcacia ggcaaatata      780
gagattagcc ttgatgaatt taagttcatg ttaatgcttg aaaataacta ccatgagttt      840
aaaaggctta accaatgggt tttgaaacca ataagtaaag atttaaacac ttacagcaat      900
atgaaattgg tggttgataa gcgaggccgc ccgactgata cgttgatttt ccaagttgaa      960
ctagatagac aaatggatct cgtaaccgaa cttgagaaca accagataaa aatgaatggt      1020
gacaaaatc caacaacat tacatcagat tctacacctac ataacggact aagaaaaaca      1080

```

-continued

ctacacgatg	ctttaactgc	aaaaattcag	ctcaccagtt	ttgaggcaaa	atTTTTgagt	1140
gacatgcaaa	gtaagcatga	tctcaatggt	tcgTtctcat	ggctcaccga	aaaacaacga	1200
accacactag	agaacatact	ggctaaatac	ggaaggatct	gaggttctta	tggtctttgt	1260
atctatcagt	gaagcatcaa	gactaacaaa	caaaagtaga	acaactgttc	accgTtacat	1320
atcaaaggga	aaactgtcca	tatgcacaga	gataatctca	tgacccaaaac	cggtagctag	1380
aggggccgca	ttaggcacc	caggctttac	actttatgct	tccggctcgt	ataatgtgtg	1440
gattttgagt	taggatccgg	cgagattttc	aggagctaag	gaagctaaaa	tgagaaaaa	1500
aatcactgga	tataccaccg	ttgatatac	ccaatggcat	cgtaaagaac	atTTTTgaggc	1560
atTtcagtca	gTtgcTcaat	gtacctataa	ccagaccgTt	cagctggata	ttacggcctt	1620
tttaaagacc	gtaaagaaaa	ataagcacia	gttttatccg	gcctttattc	acattcttgc	1680
ccgcctgatg	aatgctcatc	cggaattccg	tatggcaatg	aaagacggTg	agctggTgat	1740
atgggatagt	gttaccocct	gttacaccgt	tttccatgag	caaaactgaaa	cgTtttcatc	1800
gctctggagt	gaataccacg	acgatttccg	gcagTttcta	cacatatatt	cgcaagatgt	1860
ggcgtgttac	ggTgaaaacc	tgccctattt	ccctaaaggg	tttattgaga	atatgttttt	1920
cgtctcagcc	aatccctggg	tgagTttcac	cagTtttgat	ttaaactggg	ccaatatgga	1980
caacttcttc	gcccccgTtt	tcaccatggg	caaattattat	acgcaaggcg	acaaggtgct	2040
gatgccgctg	gcgattcagg	ttcatcatgc	cgtctgtgat	ggcttccatg	tcggcagaat	2100
gcttaaatgaa	ttacaacagt	actgcgatga	gtggcagggc	ggggcgtaaa	cgcgTggatc	2160
cggcttacta	aaagccagat	aacagtatgc	gtatttgcgc	gctgattttt	gcggtctaga	2220
ggTttagcaa	gatggcagcg	cctaaatgta	gaatgataaa	aggattaaga	gattaatTtc	2280
cctaaaaaatg	ataaaaaaag	cgTtttgaaa	gcgctgtgtt	ttttggTttg	cagtcagagt	2340
agaatagaag	tatcaaaaaa	agcaccgact	cggtgccact	ttttcaagtt	gataacggac	2400
tagccttatt	ttaacttgct	atttctagct	ctaaaactga	gaccatcccc	ggtctctact	2460
gcagaattat	caccgccagT	ggtatttatg	tcaacaccgc	cagagataat	ttatcaccgc	2520
agatggTtat	cgatgaagat	tcttgctcaa	ttgttatcag	ctatgcgccg	accagaacac	2580
cttgccgatc	agccaaaact	ctcttcaggc	cactgactag	cgataaactt	ccccacaacg	2640
gaacaactct	cattgcatgg	gatcattggg	tactgtgggt	ttagTggTtg	taaaaacacc	2700
tgaccgctat	ccctgatcag	tttcttgaag	gtaaactcat	cacccccaaag	tctggctatg	2760
cagaaatcac	ctggctcaac	agcctgctca	gggtcaacga	gaattaacat	tccgtcagga	2820
aagctTggct	tggagcctgt	tggTgcggTc	atggaattac	cttcaacctc	aagccagaat	2880
gcagaatcac	tggctTTTTt	ggtTgtgctt	acccatctct	ccgcatcacc	ttTggTaaag	2940
gttctaagct	taggtgagaa	catccctgcc	tgaacatgag	aaaaaacagg	gtactcatac	3000
tcacttctaa	gtgacggctg	cataactaac	gcttcataca	tctcgtagat	ttctctggcg	3060
attgaagggc	taaattcttc	aacgctaaact	ttgagaattt	ttgtaagcaa	tgcggcgtta	3120
taagcattta	atgcattgat	gccattaaat	aaagcaccaa	cgctgactg	ccccatcccc	3180
atcttgtctg	cgacagattc	ctgggataag	ccaagtccat	ttttcttttt	ttcataaatt	3240
gctttaaggc	gacgtgcgTc	ctcaagctgc	tcttTgtTta	atggTttctt	ttttgtgctc	3300
atacgTtaaa	tctatcaccg	caagggataa	atatctaaca	ccgtgcgtgt	tgactatTtt	3360

-continued

acctctggcg	gtgataatgg	ttgcagggcc	catttttagga	ggcaaaaatg	gataagaaat	3420
actcaatagg	cttagctatc	ggcacaaaata	gcgtcgggatg	ggcgggtgatc	actgatgaat	3480
ataaggttcc	gtctaaaaag	ttcaagggttc	tgggaaatac	agaccgccac	agtatcaaaa	3540
aaaatcttat	aggggctctt	ttatttgaca	gtggagagac	agcgggaagcg	actcgtctca	3600
aacggacagc	tcgtagaagg	tatacacgtc	ggaagaatcg	tatttgttat	ctacaggaga	3660
ttttttcaaa	tgagatggcg	aaagttagatg	atagtttctt	tcatcgactt	gaagagtctt	3720
ttttgggtga	agaagacaag	aagcatgaac	gtcatcctat	ttttgaaat	atagtagatg	3780
aagttgctta	tcatgagaaa	tatccaacta	tctatcatct	gcgaaaaaaa	ttggtagatt	3840
ctactgataa	agcggatttg	cgcttaatct	atttggcctt	agcgcataatg	attaagtttc	3900
gtggtcattt	tttgattgag	ggagatttaa	atcctgataa	tagtgatgtg	gacaaaactat	3960
ttatccagtt	ggtacaaaacc	tacaatcaat	tatttgaaga	aaaccctatt	aacgcaagtg	4020
gagtagatgc	taaagcgatt	ctttctgcac	gattgagtaa	atcaagacga	ttagaaaaatc	4080
tcattgctca	gctccccggg	gagaagaaaa	atggcttatt	tgggaatctc	attgctttgt	4140
cattgggttt	gaccctaat	tttaaatcaa	attttgattt	ggcagaagat	gctaaattac	4200
agctttcaaa	agatacttac	gatgatgatt	tagataatth	attggcgcaa	attggagatc	4260
aatatgctga	ttgttttttg	gcagctaaga	atztatcaga	tgctatttta	ctttcagata	4320
tcctaagagt	aaactactgaa	ataactaagg	ctcccctatc	agcttcaatg	attaaacgct	4380
acgatgaaca	tcatcaagac	ttgactcttt	taaaagcttt	agttcgacaa	caacttcag	4440
aaaagtataa	agaaatcttt	tttgatcaat	caaaaaacgg	atatgcaggt	tatattgatg	4500
ggggagctag	ccaagaagaa	ttttataaat	ttatcaaacc	aattttagaa	aaaatggatg	4560
gtactgagga	attattgggtg	aaactaaatc	gtgaagattt	gctgcgcaag	caacggacct	4620
ttgacaacgg	ctctattccc	catcaaattc	acttgggtga	gctgcatgct	attttgagaa	4680
gacaagaaga	cttttatoca	tttttaaaag	acaatcgtga	gaagattgaa	aaaatcttga	4740
cttttcgaat	tccttattat	gttgggtccat	tggcgcgtgg	caatagtcgt	tttgcatgga	4800
tgactcggaa	gtctgaagaa	acaattacc	catggaatth	tgaagaagtt	gtcgataaag	4860
gtgcttcagc	tcaatcattt	attgaacgca	tgacaaaact	tgataaaaat	cttccaaatg	4920
aaaaagtact	accaaaaacat	agtttgcttt	atgagtattt	tacggtttat	aacgaattga	4980
caaaggtcaa	atatgttact	gaaggaatgc	gaaaaccagc	atctctttca	ggtgaacaga	5040
agaaagccat	tgttgattta	ctcttcaaaa	caaatcgaaa	agtaaccggt	aagcaattaa	5100
aagaagatta	tttcaaaaaa	atagaatggt	ttgatagtgt	tgaatthtca	ggagttgaag	5160
atagatttaa	tgcttcatta	ggtacctacc	atgatttgct	aaaaattatt	aaagataaag	5220
atthtttggg	taatgaagaa	aatgaagata	tcttagagga	tattgthtta	acattgacct	5280
tatttgaaga	tagggagatg	attgaggaaa	gacttaaaac	atatgctcac	ctctttgatg	5340
ataaggtgat	gaaacagctt	aaacgtcgcc	gttatactgg	ttggggacgt	ttgtctcgaa	5400
aattgattaa	tggatttagg	gataagcaat	ctggcaaaac	aatattagat	tttttgaat	5460
cagatggtht	tgccaatcgc	aatthtatgc	agctgatcca	tgatgatagt	ttgacattta	5520
aagaagacat	tcaaaaagca	caagtgtctg	gacaaggcga	tagthtcat	gaacatattg	5580
caatthtagc	tggtagccct	gctattaaaa	aaggtattht	acagactgta	aaagttgtht	5640

-continued

atgaattggt caaagtaatg gggcggcata agccagaaaa taccgttatt gaaatggcac 5700
gtgaaatca gacaactcaa aagggccaga aaaattcgcg agagcgtatg aaacgaatcg 5760
aagaaggat caaagaatta ggaagtcaga ttcttaaga gcacctgtt gaaaatactc 5820
aattgcaaaa tgaaaagctc tatctctatt atctccaaaa tggagagac atgtatgtgg 5880
accaagaatt agatattaat cgtttaagtg attatgatgt cgatgccatt gttccacaaa 5940
gtttccttaa agacgattca atagacaata aggtcttaac gcgttctgat aaaaatcgtg 6000
gtaaatcgga taacgttcca agtgaagaag tagtcaaaaa gatgaaaaac tattggagac 6060
aacttctaaa cgccaagtta atcactcaac gtaagtttga taatttaacg aaagctgaac 6120
gtggagggtt gagtgaactt gataaagctg gttttatcaa acgccaattg gttgaaactc 6180
gccaatcac taagcatgtg gcacaaattt tggatagtcg catgaatact aaatacgtg 6240
aaaatgataa acttattoga gaggttaaag tgattacctt aaaatctaaa ttagtttctg 6300
acttccgaaa agatttccaa ttctataaag tacgtgagat taacaattac catcatgccc 6360
atgatgcgta tctaaatgcc gtcgttgaa ctgctttgat taagaaatat ccaaaacttg 6420
aatcggagtt tgtctatggt gattataaag tttatgatgt tcgtaaaatg attgctaagt 6480
ctgagcaaga aataggcaaa gcaaccgcaa aatatttctt ttactctaat atcatgaact 6540
tcttcaaac agaaattaca cttgcaaatg gagagattcg caaacgccct ctaatcgaaa 6600
ctaattggga aactggagaa attgtctggg ataaagggcg agattttgcc acagtgcgca 6660
aagtattgtc catgccccaa gtcaatattg tcaagaaaac agaagtacag acaggcggat 6720
tctccaagga gtcaatttta ccaaaaagaa attcggacaa gcttattgct cgtaaaaaag 6780
actgggatcc aaaaaatat ggtggttttg atagtccaac ggtagcttat tcagtcctag 6840
tgggtgctaa ggtggaaaaa gggaaatcga agaagttaa atccgttaa gagttactag 6900
ggatcacaat tatggaaaga agttcctttg aaaaaatcc gattgacttt ttagaagcta 6960
aaggatataa ggaagttaa aaagacttaa tcattaaact acctaaatat agtctttttg 7020
agttagaaaa cggctgtaaa cggatgctgg ctagtgccg agaaattaca aaaggaaatg 7080
agctggctct gccaaagcaa tatgtgaatt ttttatatt agctagtcac tatgaaaagt 7140
tgaagggtag tccagaagat aacgaacaaa acaattgtt tgtggagcag cataagcatt 7200
atthagatga gattattgag caaatcagtg aattttctaa gcgtgttatt ttagcagatg 7260
ccaatttaga taaagttctt agtgcataa acaaacatag agacaaacca atacgtgaac 7320
aagcagaaaa tattattcat ttatttacgt tgacgaatct tggagctccc gctgctttta 7380
aatattttga tacaacaatt gatcgtaac gatatacgtc taaaaaaga gttttagatg 7440
ccactcttat ccatcaatcc atcactggtc tttatgaaac acgcattgat ttgagtcagc 7500
taggaggtga cggtgaggga ggttctggag gtggaggttc tgctgagtat gtgcgagccc 7560
tctttgactt taatgggaat gatgaagagg atcttcctt taagaaagga gacatctga 7620
gaatccggga taagcctgag gagcagtggt ggaatgcaga ggacagcga gaaagaggg 7680
ggatgattcc tgtcccttac gtggagaagt attccggaga ctataaggac cacgacggag 7740
actacaagga tcatgatatt gattacaaag acgatgacga taagtctagg ctcgagtccg 7800
gagactataa ggaccacgac ggagactaca aggatcatga tattgattac aaagacgatg 7860
acgataagtc taggatgacc gacgctgagt acgtgagaat ccatgagaag ttggacatct 7920

-continued

acacgtttaa	gaaacagttt	ttcaacaaca	aaaaatccgt	gtcgcataga	tgctacgttc	7980
tctttgaatt	aaaacgacgg	ggtgaacgta	gagcgtgttt	ttggggctat	gctgtgaata	8040
aaccacagag	cgggacagaa	cgtggcattc	acgcgcaaat	ctttagcatt	agaaaagtcg	8100
aagaatacct	gcgcgacaac	cccggacaat	tcacgataaa	ttggtactca	tcctggagtc	8160
cttgtgcaga	ttgcgctgaa	aagatcttag	aatggtataa	ccaggagctg	cgggggaacg	8220
gccacacttt	gaaaaatctgg	gcttgcaaac	tctattacga	gaaaaatgcg	aggaatcaaa	8280
ttgggctgtg	gaacctcaga	gataacgggg	ttgggttgaa	tgtaatggta	agtgaacact	8340
accaatgttg	caggaaaata	ttcatccaat	cgtcgcacaa	tcaattgaat	gagaatagat	8400
ggcttgagaa	gactttgaag	cgagctgaaa	aacgacggag	cgagttgtcc	attatgattc	8460
aggtaaaaat	actccacacc	actaagagtc	ctgctgttac	ttgacaggca	tcaataaaaa	8520
cgaaggctc	agtcgaaaga	ctgggccttt	cgttttatct	gttgtttgcg	gccgggtacc	8580
gagctcgaat	tactggcgcg	tcgttttaca	acgtcgtgac	tgggaaaacc	ctggcggtac	8640
ccaacttaat	cgcttgcag	cacatcccc	tttcgccagc	tggcgtaata	gcgaagaggc	8700
ccgcaccgat	cgcccttccc	aacagttgcg	cagcctgaat	ggcgaatggc	gattcacaaa	8760
aaataggtac	acgaaaaaca	agttaaggga	tgcaagttat	gcatccetta	acttacttat	8820
taaataat	atagctattg	aaaagagata	agaattgttc	aaagctaata	ttgtttaaat	8880
cgtcaattcc	tgcatgtttt	aaggaattgt	taaattgatt	ttttgtaaat	atcttcttgt	8940
attctttgtt	aaccatttc	ataacgaaat	aattatactt	ctgtttatct	ttgtgtgata	9000
ttcttgattt	ttttctat	aatctgataa	gtgagctatt	cactttaggt	ttaggatgaa	9060
aatattctct	tggaaaccata	cttaatatag	aaatatcaac	ttctgccatt	aaaaataatg	9120
ccaatgagcg	ttttgtat	aataatcttt	tagcaaaccc	gtattccacg	attaaataaa	9180
tctcatcagc	tatactatca	aaaacaattt	tgcgatttat	atccgtactt	atgttataag	9240
gtatattacc	aaatatttta	taggattggt	ttttaggaaa	tttaaacctg	aatataatcct	9300
tgtttaaaac	ttggaaatta	tcgtgatcaa	caagtttatt	ttctgtagtt	ttgcataatt	9360
tatggtctat	ttcaatggca	gttacgaaat	tacacctctg	tactaattca	agggtaaaaat	9420
gcccttttcc	tgagccgatt	tcaagatat	tatcatgttc	atttaatctt	atatttgtca	9480
ttattttctc	tatattatgt	tttgaagtaa	taaagttttg	actgtgtttt	atatttttct	9540
cgttcattat	aacctctttt	attttttctt	ccttataaaa	ttagtataat	tatagcacga	9600
gctctgataa	atatgaacat	gatgagtgat	cgttaaattt	atactgcaat	ctgatgcat	9660
tattgaataa	aagatatgag	agatttatct	agtttctttt	tttacaagaa	aaaagaaagt	9720
tcttaaagg	tttatacttt	tggtcgtaga	gcacacgggt	taacgaacta	attacgaagt	9780
aaataagctc	agtgtgttag	actttaatgt	tttttaagg	cattagtgca	tttaagcgtc	9840
agagcatggc	tttatgcoga	gaaaactatt	ggttggaatg	gcgtgtgtgt	tagccaaagc	9900
tttggcgagt	tggttggggg	tttcatggga	ttaatcccat	gaaagtacca	actcaacaac	9960
acactaacgc	ctgttggttc	caaccaatag	gaaattggaa	taagcaatta	gtataatgag	10020
agtataatgt	tggtataacg	ttagtataat	gatgcttttt	ttcattatat	tttttatgta	10080
ctttaaacct	gcacgcttat	gtgaattaga	aaaagcttaa	tcgcatttca	tagattgacc	10140
tcccaataac	tacgtgggtg	tattgggagg	tcaatctatt	tcatttgcct	cttgcctaaa	10200

-continued

```

gttcccaaat tcgagtaaga ggtatTTTTg tttttggtcg tgcctctca ttagtagttc 10260
aggggtttaac attaatactc cagtttttct ttttataata tttccttctt ctaagatttt 10320
aagtgttgtt attactgttt gtagacttgt tctgttagct tttgctattt ctcttgttgt 10380
agctatcatt gtattgttac ttaagtggac attatctagg atatagttaa cgattttaag 10440
tttttttccg ccaatcatat ctaacatact tattaattgc actatatatg cctttacgaa 10500
gttaccagac gtttgtttac ggtataactt gtctacctct atgacttctc cactttcttc 10560
gtctatgagc ctctgagagc ctttatagac tgttccatat ctttctttca tctttttctc 10620
actccttatt ttaactatt ctaactatat cataactggt ctaaaaaaaaa aagaacattt 10680
gttaaaagaa attagaacaa aatgagttaa aaattagaac aaacaaattc cttataaacc 10740
ttatcatctc aacctatatt aagattttac ctagtgaat cttcttttct atataaagcg 10800
tcggagcata tcaggggggtt atctaacgta aatgctacc ttcggctcgc ttcgctcgcg 10860
cattgac 10867

```

```

<210> SEQ ID NO 43
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: CAN1 ORF

```

```

<400> SEQUENCE: 43

```

```

ggggttaccg gccagttgg attccgttat tggagaaacc caggtgcctg ggggccaggt 60
ataata 66

```

```

<210> SEQ ID NO 44
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: CAN1 ORF

```

```

<400> SEQUENCE: 44

```

```

ccccaatggc cgggtcaacc taaggcaata acctctttgg gtccacggac cccaggtcca 60
tattat 66

```

```

<210> SEQ ID NO 45
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: mutant sequence

```

```

<400> SEQUENCE: 45

```

```

ggggttaccg gccagttgg attcccttat tggagaaacc caggtgcctg ggggccaggt 60
ataata 66

```

```

<210> SEQ ID NO 46
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target sequence

```

```

<400> SEQUENCE: 46

```

-continued

 aaatggcgag gatacgttct ctatggagga tggcataggt 40

<210> SEQ ID NO 47
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: target sequence

<400> SEQUENCE: 47

tttaccgctc ctatgcaaga gatacctcct accgtatcca 40

<210> SEQ ID NO 48
 <211> LENGTH: 47
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: CAN1 ORF

<400> SEQUENCE: 48

gattccgtta ttggagaaac ccaggtgcct ggggtccagg tataata 47

<210> SEQ ID NO 49
 <211> LENGTH: 47
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: CAN1 ORF

<400> SEQUENCE: 49

ctaaggcaat aacctctttg ggtccacgga ccccagggtcc atattat 47

<210> SEQ ID NO 50
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ade1

<400> SEQUENCE: 50

taacgataat gtcaattacg aagactgaac tggacggtat 40

<210> SEQ ID NO 51
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ade1

<400> SEQUENCE: 51

 Met Ser Ile Thr Lys Thr Glu Leu Asp Gly Ile
 1 5 10

<210> SEQ ID NO 52
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ade1

<400> SEQUENCE: 52

taacgataat gtgaattacg aagactgaac tggacggtat 40

-continued

<210> SEQ ID NO 53
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ade1

 <400> SEQUENCE: 53

 taacgataat gttaattacg aagactgaac tggacggtat 40

<210> SEQ ID NO 54
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Can1

 <400> SEQUENCE: 54

 gggggttaccg gccccagttgg attccgttat tggagaaacc 40

<210> SEQ ID NO 55
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Can1

 <400> SEQUENCE: 55

 Gly Val Thr Gly Pro Val Gly Phe Arg Tyr Trp Arg Asn
 1 5 10

<210> SEQ ID NO 56
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Can1

 <400> SEQUENCE: 56

 gggggttaccg gccccagttgg attcccttat tggagaaacc 40

<210> SEQ ID NO 57
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Can1

 <400> SEQUENCE: 57

 gggggttaccg gccccagttgg attccattat tggagaaacc 40

<210> SEQ ID NO 58
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Can1

 <400> SEQUENCE: 58

 gggggttaccg gccccagttga attcccttat tggagaaacc 40

<210> SEQ ID NO 59
 <211> LENGTH: 40
 <212> TYPE: DNA

-continued

<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Can1

<400> SEQUENCE: 59

ggggttaccg gcccgattgg attcctttat tggagaaacc 40

<210> SEQ ID NO 60
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: WT-Adel

<400> SEQUENCE: 60

acgataatgt caattacgaa g 21

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK

<400> SEQUENCE: 61

tcaatgggct aactacgttc 20

<210> SEQ ID NO 62
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK

<400> SEQUENCE: 62

ttaatgggct aactacgttc 20

<210> SEQ ID NO 63
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

<400> SEQUENCE: 63

gctgtctcag tttatggacc 20

<210> SEQ ID NO 64
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

<400> SEQUENCE: 64

cgacagagtc aaatacctgg 20

<210> SEQ ID NO 65
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

-continued

<400> SEQUENCE: 65
gctgtctcag tttatggacc 20

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

<400> SEQUENCE: 66
gctgtctcag tttatgaacc 20

<210> SEQ ID NO 67
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

<400> SEQUENCE: 67
gctgtctcag tttataaacc 20

<210> SEQ ID NO 68
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB

<400> SEQUENCE: 68
cgacagagtc aaatacttgg 20

<210> SEQ ID NO 69
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 69
tcgcttgaac atccagcgaa acaggccccc cccatcgagc agaaaaacggg ggtggatggc 60

<210> SEQ ID NO 70
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 70
tcgcttgaac atccagcgaa acaggttccc cccatcgagc agaaaaacggg ggtggatggc 60

<210> SEQ ID NO 71
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 71
tcgcttgaac atccagcgaa acaggtcccc cccatcgagc agaaaaacggg ggtggatggc 60

-continued

<210> SEQ ID NO 72
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 72

tcgcttgaac atccagcgaacacaggcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 73
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 73

tcgcttgaac atccagcgaacacaggcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 74
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 74

tcgcttgaac atccagcgaacacaggcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 75
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 75

tcgcttgaac atccagcgaacacaggtcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 76
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 76

tcgcttgaac atccagcgaacacaggcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 77
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 77

tcgcttgaac atccagcgaacacaggcccccccatcgagcagaaaacggtggtggatggc 60

<210> SEQ ID NO 78
<211> LENGTH: 60
<212> TYPE: DNA

-continued

<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 78

tcgcttgaac atccagcgaa acaggcctcc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 79
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 79

tcgcttgaac atccagcgaa acaggccttc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 80
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 80

tcgcttgaac atccagcgaa acaggccttc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 81
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 81

tcgcttgaac atccagcgaa acaggccttc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 82
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 82

tcgcttgaac atccagcgaa acaggccttc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 83
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

<400> SEQUENCE: 83

tcgcttgaac atccagcgaa acaggccccc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 84
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: target

-continued

<400> SEQUENCE: 84

tcgcttgaac atccagcgaa acaggccttt cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 85

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: target

<400> SEQUENCE: 85

tcgcttgaac atccagcgaa acaggccctt cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 86

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: target

<400> SEQUENCE: 86

tcgcttgaac atccagcgaa acaggcccct cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 87

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: target

<400> SEQUENCE: 87

tcgcttgaac atccagcgaa acaggcccca cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 88

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: target

<400> SEQUENCE: 88

tcgcttgaac atccagcgaa acaggcccct tccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 89

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: target

<400> SEQUENCE: 89

tcgcttgaac atccagcgaa acaggccccc cccatcgagc agaaaacggt ggtggatggc 60

<210> SEQ ID NO 90

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: rpoB1r

<400> SEQUENCE: 90

gctgtctcag tttatggacc 20

-continued

```

<210> SEQ ID NO 91
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: rpoB1r

<400> SEQUENCE: 91

gctgtctcag tttatgaacc                20

<210> SEQ ID NO 92
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK 8

<400> SEQUENCE: 92

actcacacca ttcaggcgcc                20

<210> SEQ ID NO 93
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK 8

<400> SEQUENCE: 93

acttacacca ttcaggcgcc                20

<210> SEQ ID NO 94
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK 12

<400> SEQUENCE: 94

tcaatgggct aactacgttc g              21

<210> SEQ ID NO 95
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: galK 12

<400> SEQUENCE: 95

ttaatgggct aactacgttc g              21

```

What is claimed is:

1. A method of modifying a targeted site of a double stranded DNA, comprising:

contacting said double stranded DNA with at least one complex which comprises (i) a nucleic acid base converting enzyme linked to (ii) a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in the targeted site of the double stranded DNA,

thereby to convert one or more nucleotides in the targeted site to one or more different nucleotides or to delete one or more nucleotides in the targeted site or to insert one or more nucleotides into said targeted site, without

introducing a double strand break (DSB) in said double stranded DNA in the targeted site, wherein the nucleic acid sequence-recognizing module is a CRISPR-Cas system, and wherein the CRISPR-Cas system comprises a nickase protein.

2. The method of claim **1** which comprises contacting the double stranded DNA with two or more complexes that each comprise a nucleic sequence-recognizing module that specifically binds to a different target nucleotide sequence.

3. The method of claim **2**, wherein the different target nucleotide sequences are present in different genes.

4. The method of claim **1**, wherein the nucleic acid base converting enzyme is a deaminase.

5. The method of claim 4, wherein the deaminase is a cytidine deaminase.

6. The method of claim 1, wherein the step of contacting comprises introducing a nucleic acid encoding the at least one complex into a cell which comprises the double stranded DNA.

7. The method of claim 6, wherein the cell is a prokaryotic cell, an eukaryotic cell, a microbial cell, a plant cell, an insect cell, an animal cell, a vertebrate cell, or a mammalian cell.

8. A method of modifying a targeted site in double stranded genomic DNA in each of two or more targeted alleles on homologous chromosomes in a polyploid cell, the method comprising:

contacting said double stranded genomic DNA of the polyploid cell with at least one complex which comprises (i) a nucleic acid base converting enzyme linked to (ii) a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in the targeted site in the double stranded genomic DNA in each of said two or more targeted alleles on homologous chromosomes in the polyploid cell,

thereby to convert one or more nucleotides in said targeted site in the double stranded genomic DNA in each of said two or more targeted alleles on homologous chromosomes to one or more different nucleotides, or to delete one or more nucleotides in said targeted site in the double stranded genomic DNA in each of said two or more targeted alleles on homologous chromosomes, or to insert one or more nucleotides into said targeted site in the double stranded genomic DNA in each of said two or more targeted alleles on homologous chromosomes,

without introducing a double strand break (DSB) in said double stranded genomic DNA, wherein the nucleic acid sequence-recognizing module is a CRISPR-Cas system, and wherein the CRISPR-Cas system comprises a nickase protein.

9. The method of claim 6, wherein the step of introducing the nucleic acid encoding the at least one complex into the cell comprises introducing an expression vector comprising the nucleic acid encoding the at least one complex into the

cell, wherein the nucleic acid is under regulation of an inducible regulatory region, the method further comprising a step of inducing expression of the nucleic acid for an expression period to stabilize the conversion of one or more nucleotides in the targeted site to one or more different nucleotides, or the deletion of one or more nucleotides, or the insertion of one or more nucleotides into said targeted site in the double stranded DNA

10. The method of claim 9, wherein the target nucleotide sequence in the targeted site in the double stranded DNA is present in a gene essential for survival of the cell.

11. A nucleic acid-modifying enzyme complex, comprising:

a nucleic acid base converting enzyme, linked to (ii) a nucleic acid sequence-recognizing module that specifically binds to a target nucleotide sequence in a targeted site of a double stranded DNA,

wherein the nucleic acid sequence-recognizing module is a CRISPR-Cas system comprising either a Cas protein that is incapable of introducing a double strand break (DSB) in double stranded DNA or a Cas protein in which cleavage activity for only one strand of double stranded DNA has been inactivated,

and wherein the complex is capable of converting one or more nucleotides in the targeted site to one or more other nucleotides, or is capable of deleting one or more nucleotides, or is capable of inserting one or more nucleotides into said targeted site, without introducing a double strand break (DSB) in double stranded DNA in the targeted site.

12. A nucleic acid encoding the nucleic acid-modifying enzyme complex of claim 11.

13. The method of claim 1, wherein the nickase protein is a Cas9 D10A mutant nickase protein (nCas9(D10A)).

14. The method of claim 1, wherein the nickase protein is a Cas9 H840A mutant nickase protein (nCas9(H840A)).

15. The nucleic acid-modifying enzyme complex of claim 11, wherein only one of two DNA cleavage abilities of the Cas protein is inactivated.

16. A nucleic acid encoding the nucleic acid-modifying enzyme complex of claim 15.

* * * * *