

Prepared 6-30-05

	NDC 63135-312-04
	<i>Topical Anesthetic Gel</i>
	Drug Facts
	Active Ingredient Lidocaine 4% w/w..... Topical Anesthetic
	Purpose
	Uses For the temporary relief of pain and itching due to minor: ■ skin irritations ■ burns ■ cuts ■ insect bites.
	Warnings <i>For external use only</i>
	Flammable Keep away from sparks, heat and fire.
	When using this product ■ do not use in or near the eye ■ do not use in large quantities, particularly over raw surfaces or blistered areas ■ do not exceed recommended dosage unless directed by a doctor.
	Stop use and ask a doctor if:
	■ allergic reaction occurs ■ condition worsens or does not improve within 7 days ■ symptoms clear up and return within a few days ■ you notice any unusual effects.
	Keep out of the reach of children
	If swallowed, get medical help or contact a Poison Control Center right away.
	Directions ■ Adults and children 2 years and older: Apply externally to the affected area up to 3 to 4 times a day ■ Children under 2 years of age: consult a doctor.
	Inactive Ingredients Aloe Barbadosis Leaf Extract, Benzyl Alcohol, Carbomer, Disodium EDTA, Glycerin, Glyceryl Laurate, SD Alcohol 40, Simmondsia Chinensis (Jojoba) Seed Oil, Water.
	Other information Apply a thick layer and allow 30-60 min. for numbness to develop. Optional: cover with occlusive film.
	Questions Call toll free 800-677-9299 or visit www.topicaïne.com
	Manuf. for:  ESBA Laboratories Inc. Jupiter, FL 33458
	Rev. 506
	MADE IN USA
	NET WEIGHT 4 OZ (113 grams)



Pantone 3125C (83% C+ 23% Y)

Pantone 293 CVC (100 C+56 % M)

Gradient in parts of Topicaïne Name