NDC 63135-312-30

Topical Anesthetic Gel



Lidocaine 4%

Net Wt. 1 Oz (30 grams)

Topical Anesthetic Gel



Lidocaine 4%

www.topicaine.com

Drug Facts

Active Ingredient Lidocaine 4% w/w...

Topical Anesthetic

Uses For the temporary relief of pain and itching due to : irritations ■minor burns ■ minor cuts ■ insect bites

Warnings

For external use only

Manuf, for Jupiter FL 33458

Rev 511 MADE IN USA Topical Anesthetic Gel



Lidocaine 4% Net Wt. 1 Oz (30 grams) www.topicaine.com 800-677-9299

Drug Facts (continued)

Flammable Keep away from sparks, heat and fire. Cap tube tightly when not in use.

When using this product ■ do not use in or near the eye ■ do not use in large quantities, particularly over raw surfaces or blistered areas ■do not exceed recommended dosage unless directed by a doctor

Stop use and ask a doctor if: ■allergic reaction occurs ■condition worsens or does not improve within 7 days ■symptoms clear up and return within a few days ■ you notice any unusual effects such as blurred vision, dizziness or drowsiness, difficulty breathing, trembling, chest pain, or irregular heartbeat.

UPC Symbol: 363135312302 (UCC-12)

www.topicaine.com 800-677-9299

TOPICAINE ®

is a registered trademark of ESBA LABORATORIES Inc.

Drug Facts (continued)

Keep out of the reach of children If swallowed, get medical help or Contact a Poison Control Center right away

Directions Adults and children 2 years and older. Apply externally to the affected area up to 3 to 4 times a day ■ Children under 2 years of age: consult a doctor

Inactive Ingredients Aloe Barbadensis Leaf Extract, Benzyl Alcohol, Carbomer, Disodium EDTA, Glycerin, Glyceryl Laurate, SD Alcohol 40, Simmondsia Chinensis (Jojoba) Seed Oil, Water.

Purpose

minor skin