

GROPRO

AWAKENING

POLLINATION

3-3-5

GUARANTEED ANALYSIS

Total Nitrogen (N)	3.00%
0.30% Urea Nitrogen	
2.70% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	3.00%
Soluble Potash (K ₂ O)	5.00%
Boron (B)	0.10%
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc (Zn)	
*Maximum Chlorine (Cl) 0.54%	

Derived from: Fish protein hydrolysate, Soy protein hydrolysate, Potassium phosphate, Potassium chloride, Boric acid, Zinc sulfate.

ALSO CONTAINS
NONPLANTFOOD
INGREDIENT: 3.00%
Seaweed extract
(Purpose: Microbe food)

KEEP OUT OF REACH OF CHILDREN WARNING

See Next Panel for Precautionary Statements



GHS07
May be harmful if swallowed.
Causes mild skin and eye irritation.

NET CONTENTS:

9.2 lbs. per gallon (1102.4 grams/liter) @ 68 F

■ **22.5 GAL**
(9.46353 L)

■ **50 GAL**
(189.271 L)

■ **264 GAL**
(999.349 L)

MANUFACTURED BY: GROPRO

12600 West Frontage Road,
Brunsville MN 55337
Phone: 1-833-476-7761
www.groproag.com

PRECAUTIONARY STATEMENTS:

- Do not get in eyes, on skin or on clothing.
- Wash hands, forearms, and other exposed areas thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Do not breathe vapors, mist, or spray.

WARNING: Contains Boron. Do not use on plants sensitive to Boron. Use of Boron on plants other than those recommended may result in serious injury to the plants.

Personal Protective Equipment (PPE): Wear protective eyewear (goggles or face shield), chemical-resistant gloves, long-sleeved shirt and long pants, and shoes plus socks when using this product. Take off any contaminated clothing and wash before reuse.

Keep product locked up and out of the reach of children. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or smoking tobacco. Remove and wash contaminated clothing before reuse. Do not take internally. Avoid contact with or inhalation of spray application mist if present. Do not apply this product in such a manner as to directly expose workers or other persons.

If product is being mixed with pesticides and/or spray adjuvants, follow all precautionary statements on the accompanying product(s) labeling. Not for human or animal consumption.

Information about the components of this lot of fertilizer may be obtained by writing to GROPRO INC, and giving the lot number which is found on the container. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Contents Made In South Korea
FLORIDA FERTILIZER LICENSE NUMBER F002877

FIRST AID

IF SWALLOWED: • Call a POISON CENTER or doctor immediately. • Rinse mouth. Do NOT induce vomiting. • Do not give anything by mouth to an unconscious person. • Immediately call a POISON CENTER or doctor.

IF IN EYES: • Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. • Call a poison control center or doctor for treatment advice.

IF INHALED: • Move person to fresh air and keep at rest in a position comfortable for breathing if they feel unwell. • If not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a POISON CENTER or doctor for treatment advice.

IF ON SKIN OR HAIR: • Take off immediately all contaminated clothing. Rinse skin with water or shower. • Get medical attention if irritation develops or persists. • Wash contaminated clothing before reuse.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only. Keep container tightly closed. Do not allow water to be introduced into the contents of this container. Do not store near heat or open flame. Do not store with oxidizing agents or ammonium nitrate. Keep locked up and out of the reach of children

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) during mixing and loading and add rinse water to spray tank. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at www.acrecycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

For help in chemical emergencies involving spill, leak, fire or exposure, call toll free 1- 800-424-9300.

DIRECTIONS FOR USE

SHAKE CONTAINER WELL, add AWAKENING® to partially filled spray tank. Wash any residual AWAKENING® with remaining water. All other spray materials should be added prior to AWAKENING®. AWAKENING® may be applied diluted for conventional ground foliar applications (50-500 gallons of water per acre) or at high concentrations for ground low volume or aerial applications (5 to 30 gallons of water per acre). In the case of very alkaline spray water (ph 7.5 to 8.0), it is strongly recommended to use an acidifying agent or buffering agent to lower the pH of the mixture down to 6.5 - 7.0.

CONDITIONS OF SALE—LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

IMPORTANT: READ BEFORE read the entire Directions for Use. Conditions of Warranties and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer or Warranties and Limitation of Liability.

Conditions: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness of other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of GROPRO Inc. All such risks shall be assumed by the user or buyer.

Disclaimer or Warranties: To the extent consistent with applicable law, GROPRO Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of GROPRO Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, GROPRO Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

Limitations of Liability: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at GROPRO Inc's election, the replacement of product.

GENERAL INFORMATION

Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by crops or other intended plants.

AWAKENING® is a liquid foliar formulation of essential nutrient and micronutrients.

AWAKENING® provides the plant with NPK, boron and zinc.

AWAKENING® efficiently corrects Nitrogen, Phosphorus, Potassium, zinc and boron deficiencies on most agronomic crops.

COMPATIBILITY

AWAKENING® is compatible with most fungicides and insecticides, provided they are used in accordance with the registered label directions. Some compatibility problems may occur. When there is a question of compatibility prior to charging the spray tank, use the "jar compatibility test" in the following manner. Add 30 fl. oz. (20 parts) water to a glass jar with at least one quart capacity, then add 1.5 fl. oz. (1/3 part) of the products to be tank mixed. Shake well and allow to stand for at least 5 minutes. Observe to determine if any flocculation, precipitation, or unusual settling out occurs. If flocculation, precipitation, or other incompatible properties, etc. occur, do not tank mix AWAKENING® with this product. Select another product with suitable compatibility properties.

CROP RECOMMENDATIONS

FRUIT CROPS

Apply AWAKENING® to most fruit crops at pre-bloom, full bloom and soon after fruit set at a rate of 2 to 3 quarts per acre.

APPLES & PEARS: Dosage per acre per application: 2.25-3 pints. Three applications: 1) Pre-bloom. 2) Early king bloom (for apples)/bloom (for pears). 3) Post bloom (petal fall - fruit set).

BLUEBERRY, CRANBERRY AND CANE BERRY: Dosage per acre per application: 1.25 -2 pints. Four applications: 1) At first bloom. 2) At full bloom. 3) At fruit set. 4) At petal fall.

CHERRIES: Dosage per acre per application: 3-3.5 pints. Three applications: 1) At first white flower. 2) At petal fall. 3) At shuck fall.

GRAPES: Raisin, Table and Wine: Three applications: 1) 12 inch shoot growth, 2 pints per acre. 2) At bloom stage, 2 pints per acre. 3) After fruit set, 2 pints per acre.

Concord, Niagara and other American cultivars: Two applications: 1) Early bloom, 3-3.25 pints per acre. 2) Full bloom, 3 pints per acre.

OLIVES: Dosage per acre per application: 2.5 pints. Three applications: 1) Bud differentiation. 2) Early to full bloom. 3) 1st growth stage of fruit.

STONE FRUIT (Peaches, Nectarines, Apricots, Plums): Dosage per acre per application: 2.5 pints-3 pints. Three applications: 1) Early bloom. 2) Petal fall. 3) 10-20 days later (dependent upon variety).

STRAWBERRIES: Dosage per acre per application: 1.5 pints. 1) Within 14 days of transplant. 2) At first bloom. 3) 14-21 days later. Subsequent applications: Every 14-21 days thereafter.

VEGETABLE CROPS

FRUIT-BEARING VEGETABLES: For fruit-bearing vegetable crops not specifically listed on the label, apply AWAKENING® at pre-bloom, bloom and every one to two weeks thereafter at a rate of 2.5 to 3.5 pints per acre.

POTATOES (Irish): Dosage per acre per application: 2.5 pints. Two applications: 1) 100% sprouting. 2) 15 days after first treatment.

SNAP BEANS, PEAS AND OTHER LEGUMES: Dosage per acre per application: 3.25 pints. Two applications: 1) At 1st bloom. 2) At first pod.

TOMATOES, PEPPERS, CUCURBITS AND OTHER FRUITING VEGETABLES: Dosage per acre per application: 2.5 -3 pints. Three applications: 1) at first flower. 2) 7-14 days later. 3) 7-14 days later.

OTHER CROPS

ALMONDS & PISTACHIOS: Dosage per acre per application: 3 - 3.5 pints. Two applications: 1) Pre-bloom to early bloom. 2) Full bloom to petal fall. Or, apply one application of 6 pints per acre at full bloom.

COTTON: Dosage per acre per application: 1.5 -2 pints. One application: 1) At early to mid-bloom.

ENGLISH WALNUTS & PECANS: Dosage per acre per application: 4 pints. Two applications: 1) At early pistillate bloom. 2) At full pistillate bloom.

PEANUTS: Dosage per acre per application: 2 pints. Two applications: 1) At pegging. 2) 10-14 days after.

SUGAR BEET: 2.5 - 3 pints per acre applied at 8- to 10-leaf stage followed by 2.25 pints per acre 14-21 days later.

SUNFLOWER: 2.25 pints per acre applied at 3 pairs of true leaves to flower bud appearance.