



Engineered Chemistries
SUPERIOR SOLUTIONS

1 (800) 426-0512



HOME

PRODUCTS ▾

CONTACT US

HISTORY OF WECEM

SDS

LOGIN

0

HOME > [DISINFECTANTS](#) > D-GERM SPRAY DISINFECTANT AEROSOL

PRODUCT CATEGORIES ▾



D-GERM SPRAY DISINFECTANT AEROSOL

SKU: D-Germ Spray Disinfectant

Categories: [Aerosols](#), [Disinfectants](#)

* Registered with the U.S. Environmental Protection Agency (EPA).

D-Germ Spray Disinfectant:

A Broad-Spectrum Disinfectant for Environmental Surfaces.

Eliminates Odors by Attacking them at their Source.
Disinfectant – Tuberculocidal – Virucidal

TUBERCULOCIDAL ACTIVITY: This product exhibits disinfectant efficacy against *Mycobacterium bovis* (BCG) in 3 minutes at 20°C when used as directed on hard, nonporous, inanimate surfaces in the presence of 5% blood serum.

BACTERICIDAL and *VIRUCIDAL ACTIVITY: This product exhibits disinfectant efficacy in 3 minutes against the following organisms when used as directed on hard, non-porous, inanimate surfaces in the presence of 5% blood serum: *Pseudomonas aeruginosa* (ATCC 15442), *Staphylococcus aureus* (ATCC 6538), *Salmonella enterica* (ATCC 10708), Methicillin-Resistant *Staphylococcus aureus* (MRSA, ATCC 33592), Influenza A2 Virus

(A/2/Japan/305/57, strain H2N2), Human Immunodeficiency Virus Type 1 (HIV-1) (strain H2N2), Human Immunodeficiency Virus Type 1 (HIV-1) (strain IIB), and Herpes Simplex Virus Type 2 (strain G, ATCC VR-734). ***KILLS HIV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces / objects with blood or body fluids and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1.

USED IN: HOSPITALS AND OTHER MEDICAL, INDUSTRIAL, INSTITUTIONAL COMMERCIAL AND RESIDENTIAL FACILITIES.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

HOLD CONTAINER UPRIGHT. SHAKE OCCASIONALLY DURING USE.

FOR TUBERCULOCIDAL USE: Remove heavy soil and spray from 6-8" until thoroughly wet. Allow surface to remain undisturbed for at least 3 minutes at 20o C before wiping.

FOR DISINFECTING and deodorizing: remove heavy soil and spray from 6-8" until thoroughly wet. Allow surface to remain undisturbed for at least 3 minutes before wiping. For use against HIV-1 see special instructions for cleaning.

THIS IS AN EPA REGISTERED PRODUCT, DELIVERY WILL BE MADE ONLY IN THE FOLLOWING STATES: AR, CA, LA, MS, TX



LABEL



SDS



Ingredient Disclosure

To request a quote or sample, please register with Wechem.

PAGES

- [HOME](#)
- [PRODUCTS](#)
- [HISTORY OF WECHEM](#)
- [CORPORATE NEWS](#)
- [JOB OPPORTUNITIES](#)
- [CONTACT US](#)

CONTACT US

5734 Susitna Drive
Harahan, LA 70123

Toll Free | 1 (800) 426 - 0512
Phone | (504) 733 - 1152
Fax | (504) 733 - 2218

SEARCH PRODUCTS

SEARCH

DISCLAIMER

- [TERMS AND CONDITIONS](#)
- [PRIVACY POLICY](#)
- [SITE MAP](#)

Copyright(©) 2020 - Wechem, INC. Managed by Yael Consulting



Engineered Chemistries
SUPERIOR SOLUTIONS

1 (800) 426-0512



HOME

PRODUCTS ▾

CONTACT US

HISTORY OF WECEM

SDS

LOGIN

 0

[HOME](#) > [DISINFECTANTS](#) > D-GERM FOAMING GERMICIDAL CLEANER AEROSOL

PRODUCT CATEGORIES ▾



D-GERM FOAMING GERMICIDAL CLEANER AEROSOL

SKU: D-Germ Foaming Germicidal

Categories: [Aerosols](#), [Disinfectants](#)

* Registered with the U.S. Environmental Protection Agency (EPA).

KILLS HIV-1, HBV, AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES

CONTROLS MOLD AND MILDEW

FUNGICIDAL AGAINST ATHLETE'S FOOT FUNGUS (Trichophyton Mentagrophytes)

TUBERCULOCIDAL DISINFECTANT ACTIVITY AGAINST: Mycobacterium tuberculosis (BCG) in 10 minutes at 20 degrees Centigrade when used as directed on previously cleaned hard nonporous inanimate surfaces.

BACTERICIDAL DISINFECTANT ACTIVITY AGAINST: Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, Escherichia coli 0157:H7, Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin intermediate resistant Staphylococcus aureus (VISA), Vancomycin resistant Enterococcus faecalis (VRE) and meets the requirements for hospital use

D-Germ Foaming Germicidal Cleaner is designed for cleaning, deodorizing, and disinfecting for hospitals, nursing homes, schools, hotels, homes, and food processing establishments. This product is effective where housekeeping is of prime importance in controlling cross-contamination from treated surfaces.

This product when used as directed is formulated to disinfect inanimate hard non-porous surfaces: walls, floors, sink tops, toilet bowls, toilet seats, chairs, tables, telephones, restrooms, shower stalls, bathtubs, hampers, and cabinets.

This product is not for use on medical device surfaces.

This product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls, and other areas which are prone to odors caused by microorganisms.

DISINFECTION, DEODORIZATION AND CLEANING

Shake can before using. Hold container six to eight inches from surface to be treated. Spray area until it is covered with white foam. Allow foam to penetrate and remain wet for 10 minutes. No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. This product will not leave grit or soap scum.

For heavily soiled areas, a pre-cleaning step is required before treatment.

When used in food processing establishments, cover or remove all food and packaging materials, remove all gross oils, saturate all surfaces with the spray product, scrub to loosen all soils, allow to soak for 10 minutes, thoroughly rinse all wetted and cleaned surfaces with potable water.

THIS IS AN EPA REGISTERED PRODUCT, DELIVERY WILL BE MADE ONLY IN THE FOLLOWING STATES: AR, LA, MS, TX



LABEL



SDS

To request a quote or sample, please register with Wechem.

PAGES

- [HOME](#)
- [PRODUCTS](#)
- [HISTORY OF WECHEM](#)
- [CORPORATE NEWS](#)
- [JOB OPPORTUNITIES](#)
- [CONTACT US](#)

CONTACT US

5734 Susitna Drive
Harahan, LA 70123

Toll Free | 1 (800) 426 - 0512
Phone | (504) 733 - 1152
Fax | (504) 733 - 2218

SEARCH PRODUCTS

SEARCH

DISCLAIMER

- [TERMS AND CONDITIONS](#)
- [PRIVACY POLICY](#)
- [SITE MAP](#)

Copyright(©) 2020 - Wechem, INC. Managed by Yael Consulting



Engineered Chemistries
SUPERIOR SOLUTIONS

1 (800) 426-0512



HOME

PRODUCTS ▾

CONTACT US

HISTORY OF WECHEM

SDS

LOGIN

 0

HOME > DISINFECTANTS > D GERM TB

PRODUCT CATEGORIES ▾



D GERM TB

SKU: D Germ TB

Category: [Disinfectants](#)

* Registered with the U.S. Environmental Protection Agency. * Kills Pandemic 2009 H1N1 Influenza A Virus.

D-GERM TB- Ready to Use Spray & Wipe Germicidal Cleaner is designed specifically as a general non-acid cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food service establishments (restaurants) (commercial kitchens), transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross-contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact

with food. In addition, this product deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

DIRECTIONS FOR USE It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces. Fill or refill the 32 ozs. D-Germ TB Ready to Use Spray & Wipe Germicidal Cleaner spray bottle with this product refill. Pour product into spray bottle over a sink or tub basin. **DISINFECTION, DEODORIZING, AND CLEANING-** Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from the surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum. To disinfect food service establishment food contact surfaces: Before using this product food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow this product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. do not use on eating utensils, glassware, and cookware. Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, the product kills the following organisms on hard, non-porous, inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA): Pseudomonas aeruginosa, (ATCC 15442), Staphylococcus aureus, (ATCC 6538), Salmonella (choleraesuis) enterica, (ATCC 10708), Escherichia coli, (ATCC 11229), Escherichia coli O157:H7,

(ATCC 43895), Streptococcus pyogenes (Necrotizing Fasciitis Group A) (V.A. Medical Center Isolate 04001), Listeria monocytogenes, (ATCC 35152), Yersinia enterocolitica, (ATCC 23715), Enterococcus faecium, (ATCC 6569), Corynebacterium ammoniagenes, (ATCC 6871), Salmonella (typhi) enterica, (ATCC 6539)

ANTIBIOTIC-RESISTANT (STRAINS OF) (ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY):

Methicillin-resistant Staphylococcus aureus (MRSA), (ATCC 33593), Vancomycin-resistant Enterococcus faecalis (VRE), (ATCC 51575), Vancomycin intermediate resistant Staphylococcus aureus (VISA), (CDC Isolate 99287), Methicillin-resistant Staphylococcus epidermidis (MRSE), (ATCC 51625), Community Associated Methicillin-resistant Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400, Community Associated Methicillin-resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY- This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium bovis) at 20 degrees Centigrade (20°C) when the treated surface is allowed to remain wet for 5 minutes. ***VIRUCIDAL**









ACTIVITY- This product kills on hard, non-porous, inanimate surfaces when allowed to remain wet for a 10 minute contact time against Canine Parvovirus (ATCC VR-2017), Hepatitis A Virus (HAV) (University of Ottawa), Poliovirus Type 1 (ATCC VR-1000) This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against Hepatitis B Virus (HBV) (Hepadnavirus Testing, Inc.), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV) (NVSL) when the treated surface is allowed to remain wet for 5 minutes; Human Coronavirus (ATCC VR-740), SARS associated Coronavirus (ZeptoMetrix), Avian Influenza A stain H3N2 (ATCC VR-2072) and Avian Influenza A strain H9N2 (Charles River Laboratories) (Avian Influenza strains H3N2 and H9N2) for 2 minutes; Paramyxovirus (Mumps) (ATCC VR-1438), Rhinovirus type 39 (ATCC VR-340) and Rotavirus (University of Ottawa) for 3 minutes; HIV-1 (associated with AIDS) (Advanced Biotechnologies, Inc.) for 1 minute (60 seconds); and Norovirus (Norwalk virus), Feline Calicivirus (University of Ottawa) and Rabies Virus (CDC) for 30 seconds. **KILLS PANDEMIC 2009 H1N1 INFLUENZA A VIRUS. KILLS HIV-1, HBV AND HCV ON**

PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface to remain wet for HBV AND HCV 5 minutes, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. **FUNGICIDAL ACTIVITY** - This product is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (ATCC 9533) (Athlete's Foot Fungus) (a cause of ringworm of the foot), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. **EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).** **MILDEWSTAT** - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

THIS IS AN EPA REGISTERED PRODUCT, DELIVERY WILL BE MADE ONLY TO THE FOLLOWING STATES: AK, AL, AR, AZ, CA, CO, CT, DE, GA, IA, ID, IL, KY, LA, MA, MD,

MI, MN, MO, MS, MT, NC, ND, NE, NH, NM, NV, NY, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VA, VT, WA, WI, WV AND WY.

 SDS	 Ingredient Disclosure	 QT LABEL
 GAL LABEL	 5 GAL LABEL	 35 GAL LABEL
 55 GAL LABEL	 TECH SHEET	

To request a quote or sample, please register with Wechem.

PAGES

- HOME
- PRODUCTS
- HISTORY OF WECEM
- CORPORATE NEWS
- JOB OPPORTUNITIES
- CONTACT US

CONTACT US

5734 Susitna Drive
Harahan, LA 70123

Toll Free | 1 (800) 426 - 0512
Phone | (504) 733 - 1152
Fax | (504) 733 - 2218

SEARCH PRODUCTS

SEARCH

DISCLAIMER

[TERMS AND CONDITIONS](#)

[PRIVACY POLICY](#)

[SITE MAP](#)

Copyright(©) 2020 - Wechem, INC. Managed by Yael Consulting

D-Germ TB Ready To Use Spray & Wipe Germicidal Cleaner is designed specifically as a general non-acid cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food service establishments (restaurants) (commercial kitchens), transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, this product deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING - Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food service establishment food contact surfaces: Before using this product food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow this product to penetrate and remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, the product kills the following organisms on hard, non-porous, inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA) - Pseudomonas aeruginosa, (ATCC 15442), Staphylococcus aureus, (ATCC 6538), Salmonella (choleraesuis) enterica, (ATCC 10708), Escherichia coli, (ATCC 11229), Escherichia coli 0157:H7, (ATCC 43895), Streptococcus pyogenes (Necrotizing Fasciitis Group A) (V.A. Medical Center Isolate 04001), Listeria monocytogenes, (ATCC 35152), Yersinia enterocolitica, (ATCC 23715), Enterococcus faecium, (ATCC 6569), Corynebacterium ammoniagenes, (ATCC 6871), Salmonella (typhi) enterica, (ATCC 6539)

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant Staphylococcus aureus (MRSA), (ATCC 33593), Vancomycin resistant Enterococcus faecalis (VRE), (ATCC 51575), Vancomycin intermediate resistant Staphylococcus aureus (VISA), (CDC Isolate 99287), Methicillin resistant Staphylococcus epidermidis (MRSE), (ATCC 51625), Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400, Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY - This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium bovis) at 20 degrees Centigrade (20°C) when the treated surface is allowed to remain wet for **5 minutes**.

***VIRUCIDAL ACTIVITY** - This product kills on hard, non-porous, inanimate surfaces when allowed to remain wet for a 10 minute contact time against: Canine Parvovirus (ATCC VR-2017), Hepatitis A Virus (HAV) (University of Ottawa), Poliovirus Type 1 (ATCC VR-100)

This product, when used on environmental, inanimate hard surfaces exhibits antiviral activity against: **Hepatitis B Virus (HBV)** (Hepadnavirus Testing, Inc.), **Hepatitis C Virus (HCV)** and **Bovine Viral Diarrhea Virus (BVDV)** (NVSL) when the treated surface is allowed to remain wet for **5 minutes**; **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZeptoMetric), **Avian Influenza A stain H3N2** (ATCC VR-2072) and **Avian Influenza A strain H9N2** (Charles River Laboratories) for **2 minutes**; **Paramyxovirus** (Mumps) (ATCC VR-1438), **Rhinovirus type 39** (ATCC VR-340) and **Rotavirus** (University of Ottawa) for **3 minutes**; **HIV-1** (associated with AIDS) (Advanced Biotechnologies, Inc.) for **1 minute (60 seconds)**; and **Norovirus** (Norwalk virus), **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.



Engineered Chemistries
SUPERIOR SOLUTIONS

D-GERM™

TB

Ready To Use Spray & Wipe Germicidal Cleaner

TUBERCULOCIDAL · DISINFECTANT
CLEANER · DEODORIZER
MILDEWSTAT (ON HARD INANIMATE SURFACES)
*VIRUCIDAL · FUNGICIDE (AGAINST PATHOGENIC FUNGI)
KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ACTIVE INGREDIENTS
n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides.....0.105%
n-Alkyl (68% C₁₂, 32% C₁₄)
dimethyl ethylbenzyl ammonium chlorides0.105%
INERT INGREDIENTS99.790%
TOTAL.....100.000%
NET CONTENTS: 1 U.S. QUART (32 FL. OZ.) BATCH #

See side panel for additional precautionary statements.
EPA Reg. No. 1839-83-34370
EPA Est. No. 72026-IL-01
U.S. PATENT No. 5,444,094

KILLS PANDEMIC 2009 H1N1 INFLUENZA A VIRUS. KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for HBV and HCV **5 minutes**, for HIV-1 **1 minute**. These contact times will not control other common types of viruses and bacteria.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

FUNGICIDAL ACTIVITY - This product is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (ATCC 9533) (Athlete's Foot Fungus) (a cause of ringworm of the foot), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

MILDEWSTAT - To control mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.
PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Non-refillable container. Do not reuse or refill container. Clean container promptly after emptying. Triple rinse as follows: Fill container 1/4 full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.

Sold by:
WECHEM, INC.
5734 Susitna Drive
Harahan, LA 70123
(504) 733-1152 • 1(800) 426-0512
FAX (504) 733-2218
www.wechem.com



D-GERM TB™

READY TO USE SPRAY & WIPE GERMICIDAL CLEANER



- Registered with the U.S. Environmental Protection Agency.
- Kills Pandemic 2009 H1N1 Influenza A Virus and COVID-19.

D-GERM TB has demonstrated effectiveness against viruses similar to COVID-19 on hard, non-porous surfaces. Therefore, D-GERM TB can be used against COVID-19 when used in accordance with the directions for use against Norovirus, Feline Calicivirus, Poliovirus, Rhinovirus or Reovirus. Refer to the CDC website for additional information.

D-GERM TB Ready to Use Spray & Wipe Germicidal Cleaner is designed specifically as a general non-acid cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food service establishments (restaurants) (commercial kitchens), transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, this product deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING - Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food service establishment food contact surfaces: Before using this product food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow this product to penetrate and remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, the product kills the following organisms on hard, non-porous, inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA):

Pseudomonas aeruginosa, (ATCC 15442), *Staphylococcus aureus*, (ATCC 6538), *Salmonella (choleraesuis) enterica*, (ATCC 10708), *Escherichia coli*, (ATCC 11229), *Escherichia coli* O157:H7, (ATCC 43895), *Streptococcus pyogenes* (Necrotizing

Fasciitis Group A) (V.A. Medical Center Isolate 04001), *Listeria monocytogenes*, (ATCC 35152), *Yersinia enterocolitica*, (ATCC 23715),

Enterococcus faecium, (ATCC 6569), *Corynebacterium ammoniagenes*, (ATCC 6871), *Salmonella (typhi) enterica*, (ATCC 6539)

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant *Staphylococcus aureus* (MRSA), (ATCC 33593), Vancomycin resistant *Enterococcus faecalis* (VRE), (ATCC 51575), Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), (CDC Isolate 99287), Methicillin resistant *Staphylococcus epidermidis* (MRSE), (ATCC 51625), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 123) Genotype USA400, Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY - This product exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20 degrees Centigrade (20°C) when the treated surface is allowed to remain wet for **5 minutes**.

***VIRUCIDAL ACTIVITY** - This product kills on hard, non-porous, inanimate surfaces when allowed to remain wet for a **10 minute** contact time against: Canine Parvovirus (ATCC VR-2017), Hepatitis A Virus (HAV) (University of Ottawa), Poliovirus Type 1 (ATCC VR-1000)

This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Hepatitis B Virus (HBV)** (Hepadnavirus Testing, Inc.), **Hepatitis C Virus (HCV)** and **Bovine Viral Diarrhea Virus (BVDV)** (NVSL) when the treated surface is allowed to remain wet for **5 minutes**; **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZeptoMatrix), **Avian Influenza A stain H3N2** (ATCC VR-2072) and **Avian Influenza A strain H9N2** (Charles River Laboratories) for **2 minutes**; **Paramyxovirus** (Mumps) (ATCC VR-1438), **Rhinovirus type 39** (ATCC VR-340) and **Rotavirus** (University of Ottawa) for **3 minutes**; **HIV-1 (associated with AIDS)** (Advanced Biotechnologies, Inc.) for **1 minute (60 seconds)**; and **Norovirus (Norwalk virus)**, **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.

KILLS PANDEMIC 2009 H1N1 INFLUENZA A VIRUS.

KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled

with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

WEICHEM, INC.
5734 Susitna Dr. Harahan, LA 70123
504.733.1152 1.800.426.0512
FAX 504.733.2218 www.wechem.com



**Engineered Chemistries
SUPERIOR SOLUTIONS**