

Made to U.S. Pharmacopeia (USP) quality, purity and potency standards. ate, Silicon Dioxide, Acacia, Maltodextrin, Hypro-Laboratory tested to dissolve within 60 minutes.

Supplement Facts

Each Tablet Contains % DV Each Tablet Contains % DV /itamin A 3500 I.U., Zinc 11 mg (29% as Beta Carotene) Selenium 55 mca. Copper 0.9 mg Vitamin D 400 I.I 100% Manganese 2.3 mg Vitamin E 30 I.U. 100% Chromium 35 mcg Vitamin K 25 mcg. Molybdenum 45 mcg Thiamin (Vit. B1) 1.5 mg. 100% Chloride 72 ma. Riboflavin (Vit. B2) 1.7 mg. 100% Potassium 80 mg 100% Niacin 20 mg Boron 150 mca Vitamin B6 2 mg. 100% Nickel 5 mcg Folic Acid 500 mcg. 125% Silicon 2 ma Vitamin B12 6 mcg. 100% Tin 10 mcg Biotin 30 mca. 10% Vanadium 10 mcg Pantothenic Acid 10 mg. Lutein± (Tagetes Calcium 200 mg., . 20% erecta) (flower) 250 mcg Iron 18 ma.. 100%

25% *Daily Value (DV) not established INGREDIENTS: Dicalcium Phosphate, Magnesium Oxide, Potassium Chloride, Calcium Carbonate, Cellulose, Ascorbic Acid, Ferrous Fumarate, Corn Starch, dl-Alpha Tocopheryl Acetate, Niacinamide, Polyvinyl Alcohol, Gelatin, Croscarmellose Sodium, d-Calcium Pantothenate, Crospovidone, Zinc Oxide, Magnesium Stearate, Titanium Dioxide, Polyethylene Glycol, Talc, Manganese Sulf

1009

Phosphorus 109 mg

Magnesium 100 mg

lodine 150 mca.

Lycopene (Lycopersicon

esculentum) (fruit) 300 mcg ..

mellose, Pyridoxine Hydrochloride, Glucose, Cupric Sulfate, Riboflavin, Thiamine Mononitrate, Vitamin A Acetate, Boric Acid, Sucrose, Folic Acid, Beta Carotene, Yellow 6 Lake, Chromium Pico-linate, Lycopene, Lutein, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Tricalcium Phosphate, Sodium Ascorbate, Tocopherols, Red 40 Lake, Phytonadione, Biotin, Sodium Metavanadate, Nickelous Sulfate, Stannous Chloride, Cholecalciferol, Cyanocobalamin, Ascorbyl Palmitate. Contains: Soy

Distributed by: BI-MART 220 South Seneca, Eugene, OR 97402

Oil- and Water-soluble Vitamins with Minerals Suggested use: Take one tablet daily with a ful

glass of water, preferably after a meal *Pertains to new forms of chromium and copper compared to original formula.

WARNING: Accidental overdose of ironcontaining products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

NO VARNISH THIS AREA

2-1/4 X 6-3/4 BM5308

CAUTION:

It is the vendors responsibility to print a scannable UPC code from either our artwork or your alternative UPC method.

