

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

INDICATIONS: For the treatment and control of gastrointestinal roundworms (including inhibited *Ostertagia ostertagi*), lungworms, grubs, horn flies, sucking and biting lice and sarcoptic mange mites in cattle.

See package insert for complete indications and use directions.

Recommended Dose: 1 mL per 22 lb of body weight.

RESIDUE WARNING: Cattle must not be treated within 48 days of slaughter for human consumption. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

This product should not be applied to self or others because it may be irritating to human skin and eyes and absorbed through the skin. To minimize accidental skin contact, the user should wear a long-sleeved shirt and rubber gloves. If accidental skin contact occurs, wash immediately with soap and water. If accidental eye exposure occurs, flush eyes immediately with water and seek medical attention.

PRECAUTIONS: Store at 20°-25°C (68°-77°F), excursions permitted between 15° and 30°C (between 59° and 86°F). Keep container tightly closed when not in use and protect from light.



Manufactured for:
Aspen Veterinary Resources,® Ltd.
Liberty, MO 64068 USA

A191FP 10/12

Manufactured by:
FIRST PRIORITY, INC.,
Elgin, IL 60123-1146



Ivermax[®]

POUR-ON for Cattle (ivermectin)

5 mg ivermectin per mL

PARASITICIDE

Kills: Roundworms (including Brown Stomach Worm), Lungworms, Grubs, Sucking Lice, Biting Lice, Mange Mites, Horn Flies

For Animal Use Only

Keep Out of Reach of Children

ANADA 200-340, Approved by FDA

NET CONTENTS:
250 mL
(8.5 fl oz)

Use only in well-ventilated areas or outdoors.

Close container tightly when not in use.

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced.

Do not use when rain is expected to wet cattle within six hours after treatment.

This product is for application to skin surface only. Do not give orally or parenterally.

Cloudiness in the formulation may occur when Ivermax[®] (ivermectin) Pour-On is stored at temperatures below 32°F. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

Antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of the skin with mange scabs or lesions, or with dermatosis or adherent materials, e.g., caked mud or manure.

Ivermectin has been associated with adverse reactions in sensitive dogs; therefore, Ivermax[®] (ivermectin) Pour-On is not recommended for use in species other than cattle.

Do not contaminate water by direct application or by the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

WARNING! FLAMMABLE! KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME AND OTHER SOURCES OF IGNITION.

WARNING! NOT FOR USE IN HUMANS.
Keep this and all drugs out of the reach of children.

Restricted Drug (California) - Use Only as Directed

To report adverse effects, obtain an MSDS, or for assistance call Aspen Veterinary Resources,® Ltd. at (816) 415-4324.

Lot No.

Exp. Date