

1802 0101101300

NDC 72493-103-03



Jelmyto™
(mitomycin) for pyelocalyceal solution
SINGLE-DOSE CARTON

Jelmyto™
(mitomycin) for pyelocalyceal solution
SINGLE-DOSE CARTON

Distributed by:
UroGen Pharma, Inc.
Princeton, NJ 08540



NDC 72493-103-03

JELMYTO™ is a Product of Belgium
Sterile Hydrogel is a Product of Israel
Usual Dosage: See Full Prescribing Information

NDC 72493-103-03
Jelmyto™
(mitomycin) for pyelocalyceal solution
SINGLE-DOSE CARTON
UroGen
Pharma

LP-P-004207/16c-1



NDC 72493-103-03

Jelmyto™
(mitomycin) for pyelocalyceal solution

SINGLE-DOSE CARTON
Warning: For Pyelocalyceal Use Only

Attention Pharmacist: Reconstitution
is required prior to dispensing.
See the full prescribing information in the pharmacy
before proceeding.

Rx Only

Contents of Carton:

- 2 Vials of JELMYTO™ (mitomycin) for pyelocalyceal solution, 40 mg per Vial
- 1 Vial Sterile Hydrogel, 20 mL per Vial
- Full Prescribing Information
- Instructions for Pharmacy
- Instructions for Administration

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F).
Avoid excessive heat over 40°C (104°F). Protect from light.
Reconstituted JELMYTO should be infused as soon as possible after reconstitution. If immediate institution
is not possible, reconstituted JELMYTO may be stored at 2°C to 8°C (36°F to 46°F) for up to 8 hours.
Protect reconstituted JELMYTO from light.
When ready to reutilize, chill JELMYTO to 3°C to 5°C (27°F to 41°F) for at least 30 minutes.
But no longer than one hour, to reconvert it to a liquid form.

