

YESCARTA® is a treatment for your non-Hodgkin lymphoma. It is used when you have failed at least two other kinds of treatment. YESCARTA® is different than other cancer medicines because it is made from your own white blood cells. [Read More...](#)




[PRESCRIBING INFORMATION & MEDICATION GUIDE](#) [IMPORTANT SAFETY INFORMATION](#) [FOR HCPs](#)

[ABOUT CAR T](#) [YESCARTA® THERAPY](#) [TREATMENT PROCESS](#) [TREATMENT CENTERS](#) [PATIENT SUPPORT](#)

WHAT IS CAR T?

CAR T cells are created by collecting your own T cells and sending them to a manufacturing facility. At the facility, the T cells are modified by adding a chimeric antigen receptor (CAR) gene. Once the CAR gene is added, your new CAR T cells are returned to the treatment facility to be infused back into your body. In your body, these CAR T cells work to recognize and attack your cancer.



Discuss CAR T therapy with your doctor to see if it might be right for you.

[LEARN HOW YESCARTA®
USES CAR T TECHNOLOGY](#)

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about YESCARTA®?

YESCARTA® may cause side effects that are life-threatening and can lead to death. Call or see your healthcare provider or get emergency help right away if you get any of the following:

- Fever (100.4°F/38°C or higher)
- Difficulty breathing
- Chills or shaking chills
- Confusion
- Dizziness or lightheadedness
- Severe nausea, vomiting, or diarrhea
- Fast or irregular heartbeat
- Severe fatigue or weakness

It is important to tell your healthcare provider that you received YESCARTA® and to show them your YESCARTA® Patient Wallet Card. Your healthcare provider may give you other medicines to treat your side effects.

Before getting YESCARTA®, tell your healthcare provider about all your medical problems, including if you have or have had:

- Neurologic problems (such as seizures, stroke, or memory loss)
- Lung or breathing problems
- Heart problems
- Liver problems
- Kidney problems
- A recent or active infection

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive YESCARTA®?

- Since YESCARTA® is made from your own white blood cells, your blood will be collected by a process called "leukapheresis" (loo-kah-fur-ee-sis), which will concentrate your white blood cells.
- Your blood cells will be sent to a manufacturing center to make your YESCARTA®.
- Before you get YESCARTA®, you will get 3 days of chemotherapy to prepare your body.
- When your YESCARTA® is ready, your healthcare provider will give it to you through a catheter placed into your vein (intravenous infusion). The infusion usually takes less than 30 minutes.
- You will be monitored where you received your treatment daily for at least 7 days after the infusion.
- You should plan to stay close to the location where you received your treatment for at least 4 weeks after getting YESCARTA®. Your healthcare provider will help you with any side effects that may occur.
- You may be hospitalized for side effects and your healthcare provider will discharge you if your side effects are under control, and it is safe for you to leave the hospital.
- Your healthcare provider will want to do blood tests to follow your progress. It is important that you do have your blood tested. If you miss an appointment, call your healthcare provider as soon as possible to reschedule.

What should I avoid after receiving YESCARTA®?

- Do not drive, operate heavy machinery, or do other dangerous things for 8 weeks after you get YESCARTA® because the treatment can cause sleepiness, confusion, weakness, temporary memory and coordination problems.
- Do not donate blood, organs, tissues, and cells for transplantation.

What are the possible or reasonably likely side effects of YESCARTA®?

The most common side effects of YESCARTA® include:

- Fever (100.4°F/38°C or higher)
- Low white blood cells (can occur with a fever)
- Low red blood cells
- Low blood pressure (dizziness or lightheadedness, headache, feeling tired, short of breath)
- Fast heartbeat
- Confusion
- Difficulty speaking or slurred speech
- Nausea
- Diarrhea

These are not all the possible side effects of YESCARTA®. Call your healthcare provider about any side effects that concern you. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see [full Prescribing Information](#), including **IMPORTANT WARNING** and Medication Guide.

INDICATION

YESCARTA® is a treatment for your non-Hodgkin lymphoma. It is used when you have failed at least two other kinds of treatment. YESCARTA® is different than other cancer medicines because it is made from your own white blood cells, which have been modified to recognize and attack your lymphoma cells.

Limitation of Use: YESCARTA® is not indicated for the treatment of patients with primary central nervous system lymphoma.

[Terms and Conditions](#) [Privacy Policy](#)

The content on this site is intended for US residents only.
YESCARTA, the YESCARTA Logo, KITE KONNECT, KITE,
and the KITE Logo are trademarks of KITE Pharma, Inc.
GILEAD is a trademark of Gilead Sciences, Inc.
All other trademarks referenced herein are the property of their respective owners.
© 2018 Kite Pharma | PRC-00390 03/2018

