



GET YOUR CARD
Have a card? [Activate](#) **>**
[Terms & Conditions apply.](#)

WHY PRISTIQ

For a decade, brand-name PRISTIQ has been prescribed to 2.5 million patients. If you have depression, ask your doctor about moving forward with brand-name PRISTIQ—only from Pfizer.



PRISTIQ IS CLINICALLY PROVEN

In clinical studies, PRISTIQ 50 mg also helped improve patients' ability to function according to a scale[†] used to measure how depression disrupts the following areas:

- Work
- Social life/leisure activities
- Family life/home responsibilities

Clinical studies of PRISTIQ 50 mg have also shown:

- **No significant difference in average weight gain** versus placebo[†]
- **A low occurrence of sexual side effects** in men and women taking PRISTIQ,

Scroll for Important Safety Information and Indication IMPORTANT SAFETY INFORMATION

Suicidality and Antidepressant Drugs


Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy or when the dose is changed should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior such as becoming agitated, irritable, hostile, aggressive, impulsive, or restless. Should these occur, report them to a doctor right away. PRISTIQ is not approved for use in children under 18.

Do not take PRISTIQ if you are allergic to desvenlafaxine, venlafaxine, or any of the ingredients in PRISTIQ. Do not take PRISTIQ if you currently take, or have taken within the last

†Sheehan Disability Scale (a validated measure of functional impairment) total score for PRISTIQ vs placebo.
‡Side effects may increase at higher doses. Data based on short-term studies and a 6-month phase of a long-term study of patients who responded to PRISTIQ. Individual results may vary.

PRISTIQ IS THE BRAND YOU RECOGNIZE

- The PRISTIQ pill has a distinctive shape and marking to let you know you're getting the brand-name medication—pill to pill, month to month
- Brand-name PRISTIQ is from a company you know—only from Pfizer
- You have the assurance that Pfizer stands behind its manufacturing standards



PRISTIQ OFFERS SAVINGS TO PATIENTS

If eligible, patients may pay as little as \$4 for each 30-day fill of brand-name PRISTIQ—and could save up to \$1,080 a year.*

[GET YOUR CARD](#)

Have a card? [Activate](#) ▶
[Terms & Conditions apply.](#)

[LEARN MORE ABOUT PRISTIQ](#)

Scroll for Important Safety Information and Indication **IMPORTANT SAFETY INFORMATION**

Suicidality and Antidepressant Drugs

Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy or when the dose is changed should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior such as becoming agitated, irritable, hostile, aggressive, impulsive, or restless. Should these occur, report them to a doctor right away. PRISTIQ is not approved for use in children under 18.

Do not take PRISTIQ if you are allergic to desvenlafaxine, venlafaxine, or any of the ingredients in PRISTIQ. Do not take PRISTIQ if you currently take, or have taken within the last 14 days, any medicine known as an MAOI (including intravenous methylene blue or the antibiotic linezolid). Allow 7 days after stopping PRISTIQ before starting an MAOI. Taking an MAOI with PRISTIQ can cause serious or even life-threatening side effects.

Before taking PRISTIQ, tell your healthcare professional about all prescription and over-the-counter

Browser address bar: <https://www.pristiq.com/about-pristiq>

Browser tabs: Query, About PRISTIQ® (desvenlaf... x)

Browser extensions: Share Browser, WebEx

Taskbar: PfizerWorld Home, IP Manager, EFR, Trademarks SharePoint Gr..., IP SharePoint Home Folder, IP Portal, IP Renewals, ACT Communications Tra..., BT On Demand, Pfizer YRR - Your Reportin..., Trademarks USPTO

HAVE A QUESTION ABOUT THE PRISTIQ SAVINGS CARD?
SPEAK TO A LIVE REPRESENTATIVE AT 1-855-440-6852
24/7 EXCEPT FOR ALL MAJOR U.S. HOLIDAYS

IMPORTANT SAFETY INFORMATION

Suicidality and Antidepressant Drugs

Antidepressants increased the risk of suicidal thinking and behavior in children, teens, and young adults. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy or when the dose is changed should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior such as becoming agitated, irritable, hostile, aggressive, impulsive, or restless. Should these occur, report them to a doctor right away. PRISTIQ is not approved for use in children under 18.

Do not take PRISTIQ if you are allergic to desvenlafaxine, venlafaxine, or any of the ingredients in PRISTIQ. Do not take PRISTIQ if you currently take, or have taken within the last 14 days, any medicine known as an MAOI (including intravenous methylene blue or the antibiotic linezolid). Allow 7 days after stopping PRISTIQ before starting an MAOI. Taking an MAOI with PRISTIQ can cause serious or even life-threatening side effects.


Before taking PRISTIQ, tell your healthcare professional about all prescription and over-the-counter medications and supplements you take or plan to take including those to treat migraines and psychiatric disorders (including other antidepressants or amphetamines); to avoid serotonin syndrome, a potentially life-threatening condition; aspirin, NSAID pain relievers, or blood thinners because they may increase the risk of bleeding.

PRISTIQ may cause or worsen some conditions, so tell your healthcare professional about all the medical conditions you have or had including:

- High blood pressure, which should be controlled before starting PRISTIQ and monitored regularly
- Heart problems, high cholesterol or triglyceride levels, a history of stroke, kidney or liver problems, or low sodium levels in your blood
- Bleeding problems
- Depression, suicidal thoughts or behavior
- Mania, bipolar disorder, or seizures or convulsions
- Are pregnant or plan to become pregnant. Talk to your healthcare provider about the risk to your unborn baby if you take PRISTIQ during pregnancy.
- Are breastfeeding or plan to breastfeed. PRISTIQ can pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with PRISTIQ.

Some people are at risk for visual problems such as eye pain, changes in vision, or swelling or redness around the eye. You may want to undergo an eye examination to see if you are at risk and get preventative treatment if you are.

Discontinuation symptoms may occur when stopping or reducing PRISTIQ, so talk to your healthcare professional before stopping or changing your dose.



Until you see how PRISTIQ affects you, be careful driving a car or operating machinery. Avoid drinking alcohol while taking PRISTIQ.

In clinical studies, most common side effects with PRISTIQ 50 mg were nausea, dizziness, sweating, constipation, and decreased appetite.

INDICATION

PRISTIQ is a prescription medication approved for the treatment of major depressive disorder in adults.

Please see [Full Prescribing Information, including BOXED WARNING](#), and [Medication Guide](#).

PRISTIQ SAVINGS CARD TERMS AND CONDITIONS

By participating in the PRISTIQ Savings Offer Program, you acknowledge that you currently meet the eligibility criteria and will comply with the terms and conditions described below:

- This Savings Offer is not valid for prescriptions that are eligible to be reimbursed, in whole or in part, by Medicaid, Medicare, Tricare, or other federal or state healthcare programs (including any state prescription drug assistance programs) and the Government Health Insurance Plan available in Puerto Rico (formerly known as “La Reforma de Salud”)
- You must deduct the savings received under this program from any reimbursement request submitted to your insurance plan, either directly by you or on your behalf
- Eligible patients may pay a minimum of \$4 per monthly prescription fill. By using the Savings Offer, eligible patients will receive a savings of up to \$90 per fill off their co-pay or out-of-pocket costs. The Savings Offer is good for a maximum savings of \$1,080 per year (\$90 per month x 12 months). The Savings Offer limits your prescription cost to \$4, subject to a \$90 maximum monthly benefit. Thus, if your co-pay or out-of-pocket cost is more than \$94, you will save \$90 off of your co-pay or total out-of-pocket costs. [Example: If your co-pay or out-of-pocket costs are \$100, you will pay \$10 (\$100 – \$90 = \$10).] If your co-pay or out-of-pocket costs are no more than \$94, you pay \$4. For a mail-order 3-month prescription, your total maximum savings may be \$270 (\$90 x 3)
- The Savings Offer is not valid when the entire cost of your prescription drug is eligible to be reimbursed by your private insurance plans or other health or pharmacy benefit programs
- The Savings Offer is not valid for Massachusetts residents whose prescriptions are covered, in whole or in part, by third-party insurance
- This Savings Offer is not valid where prohibited by law
- The Savings Offer cannot be combined with any other rebate/coupon, free trial, or similar offer for the specified prescription
- The Savings Offer may not be redeemed more than once per month per patient.
- **The Savings Offer will be accepted only at participating pharmacies**
- **The Savings Offer is not health insurance**

party insurance

- This Savings Offer is not valid where prohibited by law
- The Savings Offer cannot be combined with any other rebate/coupon, free trial, or similar offer for the specified prescription
- The Savings Offer may not be redeemed more than once per month per patient.
- **The Savings Offer will be accepted only at participating pharmacies**
- **The Savings Offer is not health insurance**
- This Savings Offer is good only in the U.S. and Puerto Rico
- The Savings Offer is limited to 1 per person during this offering period and is not transferable
- Pfizer reserves the right to rescind, revoke, or amend the program without notice
- No membership fees. The savings offer and program expire on 12/31/2020

For reimbursement when using a non-participating pharmacy/mail order: Pay for your PRISTIQ prescription and mail copy of original pharmacy receipt (cash register receipt NOT valid) with product name, date, and amount circled to: PRISTIQ Savings Offer, 2250 Perimeter Park Drive, Suite 300, Morrisville, NC 27560. Be sure to include a copy of the front of your PRISTIQ Savings Card, your name, and mailing address. Please expect up to 4 to 6 weeks for reimbursement.

Patients should always ask their doctors for medical advice about adverse events.

You may report an adverse event related to Pfizer products by calling 1-800-438-1985 (U.S. only). If you prefer, you may contact the U.S. Food and Drug Administration (FDA) directly. The FDA has established a reporting service known as MedWatch where healthcare professionals and consumers can report serious problems they suspect may be associated with the drugs and medical devices they prescribe, dispense, or use. Visit [MedWatch](#) or call 1-800-FDA-1088.

The health information contained herein is provided for educational purposes only and is not intended to replace discussions with a healthcare provider. All decisions regarding patient care must be made with a healthcare provider, considering the unique characteristics of the patient.

The product information provided in this site is intended for residents of the United States. The products discussed herein may have different product labeling in different countries.

In need of prescription assistance?
Pfizer RxPathways® connects eligible patients to assistance programs that offer insurance support, co-pay assistance, and medicines for free or at a savings.
www.pfizer.com/rxpathways
1-800-438-1985

[TERMS OF USE](#)

[PRIVACY POLICY](#)

[SITE MAP](#)

[CONTACT PFIZER](#)

[ABOUT PFIZER](#)

