



Your position: Home / Pet Supplies



### 正典球苗SCOCVAC Coccidiosis Trivalent Vaccine for Chickens, Live

Launch Date: September 01, 2020

Price:  
USD \$300.00

select:

Quantity:  
 (136 available)

ADD TO CART

BUY NOW

♥ Add to Favorite Items



Description 1 Reviews



Message Us



Animal Vaccine

### Coccidiosis Trivalent Vaccine for Chickens, Live

(*E. tenella* PTMZ Strain, *E. maxima* PMHY Strain and *E. acervulina* PAHY Strain)

**【Product Name】**Coccidiosis Trivalent Vaccine for Chickens, Live (*E. tenella* PTMZ Strain, *E. maxima* PMHY Strain and *E. acervulina* PAHY Strain)

**【Main Ingredients and Contents】**The vaccine contains 3 kinds of coccidium oocysts, including *E. tenella* (PTMZ Strain), *E. maxima* (PMHY Strain) and *E. acervulina* (PAHY Strain), each dose contains  $400 \pm 10\%$  sporulated oocysts.

**【Properties】**White or kind of solution, a little sedimentation will be shown after it has been kept statically for a while.

**【Indication and Usage】**It is used for preventing coccidiosis in broiler chicken. 14 days after vaccinated, the chicken start to generate immunity, and immunity may continue at the end of the raising period.

**【Usage and dosage】**

Message Us

1. Vaccination procedure: Put the vaccine in the drinking water, which is used for immunizing chickens with age of 3~7 days.
2. Method and dosage of vaccination: Vaccinated through drinking water. Every chicken will take 1 dose of vaccine. Every bottle has 1000 (or 2000) doses of vaccine, which should be added with 6L (or 12L) water. Then put into 1 bottle 50 g/ bottle (or 1 bottle 100 g/bottle) coccidiosis vaccine suspending agent, which will be made into suspension. The final product can be provided for 1000 (or 2000) doses of vaccine to take freely, every chicken will take 6 ml coccidiosis vaccine suspension averagely, 4-6 hours later it will be drank off.

**【Side Effects】**12~14 days after vaccinated vaccine, some chickens may excrete bloody excrement, no need to use drugs. If the bloody excrement is very severe or chickens die for coccidiosis, then feed them with Sulfaquinoxaline or sulfadimidine with recommended dosage for 1~2 days, then this problem can be controlled.

**【Precautions】**

- 1.The Product is prohibited to be stored in the frozen or near heat source place. It should be shaken fully before used.
- 2.Requirements for the application of drugs in feed: it is prohibited to add any anti-coccidium drugs in feed.
- 3.Requirements for the management of expanding the number and padding.
  - ① It is suggested that do not expand the number day by day. On the 7th day after vaccinated coccidiosis vaccine, the brood area would be expanded to that required by the chickens on the 17th day after vaccinated at once, so as to enable the chicken group to obtain same opportunities of superinfection.
  - ② Do not change the padding within the 8th~16th day after vaccinating coccidiosis vaccine.
  - ③ The humidity of padding should be better at 25%~30% (when catching the padding by hands, the palms feel slight wet).
- 4.Prevention and control of immune suppression disease: Many immune suppression diseases, such as infectious bursal disease, marek's disease, poisoning for mycotoxin, etc., will severely influence the establishment of anti-coccidium immunity, and exacerbate the reaction of vaccine. Try to avoid these diseases to interfere the effect of vaccine immunity.
5. Reduction the influence of stress factors: On the 12th~14th day after vaccinating coccidiosis vaccine, during which time the vaccine might react strongly. Try to avoid breaking the mouth, injecting other vaccine and moving chicken groups.

**【Treatment measures for waste packages】**Residual vaccine and empty bottles should not be discarded randomly, and it should be discarded after boiled.

**【Specification】** (1) 1000 doses/bottle ;(2) 2000 doses/bottle

**【Package】**10 bottles /box

**【Storage and validity】**Stored under 2~8°C, no frozen, and the validity is 6

[Message Us](#)

months.

【Approval Number】Biological Vet-drug 190462195

【Manufacturer】Foshan Standard Bio-tech Co., Ltd.

【Country of Origin】China

Use Under the Guidance of Veterinarian Only



Animal Vaccine

### Coccidiosis Quadrivalent vaccine for Chickens, live

(*E. tenella* PTMZ Strain, *E. necatrix* PNHZ Strain, *E. maxima* PMHY Strain and *E. acervulina* PAHY Strain)

【Product Name】Coccidiosis Quadrivalent vaccine for Chickens, live (*E. tenella* PTMZ Strain, *E. necatrix* PNHZ Strain, *E. maxima* PMHY Strain and *E. acervulina* PAHY Strain)

【Main Ingredients and Contents】The vaccine contains 4 kinds of coccidium

Message Us

oocysts, including *E. tenella* (PTMZ Strain), *E. necatrix* (PNHZ Strain), *E. maxima* (PMHY Strain) and *E. acervulina* (PAHYStrain), each dose contains  $1100 \pm 10\%$  sporulated oocysts.

**【Properties】** White or kind of solution, a little sedimentation will be shown after it has been kept statically for a while.

**【Indication and Usage】** It is used for preventing coccidiosis in chicken. 14 days after vaccinated, the chickens start to generate immunity, and immunity may continue at the end of the raising period.

**【Administration and dosage】**

1. Vaccination procedure: Put the vaccine in the drinking water, which is used for immunizing chickens with age of 3~7 days.
2. Method and dosage of vaccination: Vaccinated through drinking water. Every chicken will take 1 pcs. Every bottle has 1000 (or 2000) doses of vaccine, which should be added with 6L (or 12L) water. Then put into 1 bottle 50g/ bottle (or 1bottle 100 g/bottle) coccidiosis vaccine suspending agent, which will be made into suspension. The final product will be provided for 1000 (or 2000) doses of vaccine to take freely, every chicken will take 6 ml coccidiosis vaccine suspension averagely, 4-6 hours later it will be drank off.

**【Side Effects】** 12~14 days after vaccinated vaccine, some chickens may excrete bloody excrement, no need to use drugs. If the bloody excrement is very severe or chickens die for cocciosis, then feed them with Sulfaquinoxaline or sulfadimidine with recommended dosage for 1~2 days, then this problem can be controlled.

**【Precautions】**

1. The Product is prohibited to be stored in the frozen or near heat source place.
2. It is only used for vaccinating healthy chicken, and it should be shaken fully before used.
3. Requirements for the application of drugs in feed: it is prohibited to add any anti-coccidium drugs in feed.
4. Requirements for the management of expanding the number and padding
  - ① It is suggested that do not expand the number day by day. On the 7th day after vaccinated coccidiosis vaccine, the brood area would be expanded to that required by the chickens on the 17th day after vaccinated at once, so as to enable the chicken group to obtain same opportunities of superinfection.
  - ② Do not change the padding within the 8th~16th day after vaccinating coccidiosis vaccine.
  - ③ The humidity of padding should be better at 25%~30% (when catching the padding by hands, the palms feel slight wet).
5. Prevention and control of immune suppression disease: Many immune suppression diseases, such as infectious bursal disease, marek's disease, poisoning for mycotoxin, etc., will severely influence the establishment of anti-coccidium immunity, and exacerbate the reaction of vaccine. Try to avoid these diseases to interfere the effect of vaccine immunity.
6. Reduction the influence of stress factors: On the 12th~14th day after

[Message Us](#)

vaccinating coccidiosis vaccine, during which time the vaccine might react strongly. Try to avoid breaking the mouth, injecting other vaccine and moving chicken groups.

**【Treatment measures for waste packages】** Residual vaccine and empty bottles should not be discarded randomly, and it should be discarded after boiled.

**【Specification】** (1) 1000 doses/bottle

**【Package】** 10 bottles/box

**【Storage and Validity】** Stored under 2~8°C, no frozen, and the validity is 7 months.

**【Approval Number】** Biological Vet-drug 190462139

**【Manufacturer】** Foshan Standard Bio-tech Co., Ltd.

**【Country of Origin】** China

Use Under the Guidance of Veterinarian Only

MEMBER

[Order Tracking](#)

[WishList](#)

[Change Account](#)

[Create an Account](#)

OVERVIEW

[PAYMENT METHODS](#)

[SHIPPING & RETURNS](#)

[Location & Working Hours](#)

GET IN TOUCH

SUBSCRIBE

[Message Us](#)