## Official 2018 Scorecard

Sandpiper scorecard will NOT be accepted

Plo	ayers Names		
_			PSYCHOTOUR®
1 2 3 4 5 6 7 8 9	PAR SCORE 5 511 [] 4 447 [] 4 407 [] 3 226 [] 5 495 [] 3 173 [] 4 368 [] 4 370 [] 4 401 []	TRUE FALSI ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )	Removal of colon polyps can prevent colon cancer.  There are more than 1 million colon cancer survivors in the U S.  Lifetime risk of men developing colon cancer is about 1 in 22.  Surgery is the most common treatment for colon cancer.  Colon cancer is expected to cause about 51,020 deaths this year.  For most people screening should start at the age of 45.  Colon cancer can be present for years before symptoms develop.  Lifetime risk of women developing colon cancer is about 1 in 24.  Estimated # of new colon cancer cases in 2019 in US is 4,321.
10 11 12 13 14 15 16 17	4 345 [] 3 202 [] 4 325 [] 5 455 [] 4 425 [] 5 582 [] 4 373 [] 4 374 []	TRUE FALS ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )	Regular exercise may decrease the risk of developing colon cancer. Colon cancer is among the least preventable and treatable cancers. Risk factors for colon cancer include heredity. Most colorectal cancers develop from polyps. 100% of new colon cancer cases occur in people 50 or over. Colon cancer screening for people at average risk should start at 55 1 in 5 colon cancer patients are 20 to 54 years old. Risk factors for colon cancer include family history.

## **SMART SCRAMBLE**

TOTAL [\_\_\_]