

COMPOUND INTEREST?

Because custom-compounded products have not been rigorously tested, your doctor may recommend an FDA-approved therapy that is also bioidentical.



Confident coverage morning, noon, and night<sup>†</sup>  
Once-daily dosing of estradiol

- Home
- About Divigel®
- Using Divigel®
- Resources
- Offers
- News
- Menopause
- Estrogen Therapy
- Healthcare Professionals


Safety Information

### The Benefits of a Gel

Divigel® uses *transdermal delivery* to administer relief directly through your skin—no patches, no pills.

The gel medication is absorbed through your skin and goes directly into your bloodstream, avoiding circulation through the liver, so it can be used at lower doses than hormone pills<sup>2,8,9</sup>

A convenient and easy-to-use treatment approach that fits into your daily routine



### Convenient Gel Formulation

Divigel® is an appealing gel formulation that's convenient and discreet; it's very similar to the look and feel of hand sanitizer.

The clear gel is invisible and odorless when dry, so no one knows you're using medication

It dries quickly, without a sticky residue; and once you have washed your hands, there is little risk of transfer from your skin to another person's skin<sup>2</sup>

It's skin-friendly: Less than 1% of women treated with Divigel® experienced application-site reactions<sup>2</sup>

Compared with all other gel or lotion estrogen therapy products, Divigel® requires a smaller amount of gel applied to a smaller area of skin<sup>2-5</sup>

### Discreet and Easy to Use

Small packets are portable and easy to use at home and away.

Premeasured, individual-use foil packets to ensure precise, accurate dosing

Each foil packet contains a single day's dose for convenient, consistent delivery of Divigel®

Store up to 14 foil packets in a handy Divigel® travel compact, available free when you [click here](#)



[HOME](#) | [FULL PRESCRIBING INFORMATION](#) | [PATIENT INFORMATION](#) | [CONTACT](#) | [CORPORATE](#) | [REFERENCES](#) | [PRIVACY POLICY](#) | [SITE MAP](#)

#### Important Safety Information for Patients

##### What is the most important information I should know about Divigel® (an estrogen hormone)?

- Using estrogen-alone increases your chance of getting cancer of the uterus (womb). Report any unusual vaginal bleeding right away while you are using Divigel®. Vaginal bleeding after menopause may be a warning sign of cancer of the uterus (womb). Your healthcare provider should check any unusual bleeding to find out the cause.
- Do not use estrogen-alone to prevent heart disease, heart attacks, strokes or dementia (decline of brain function)
- Using estrogen-alone may increase your chances of getting strokes or blood clots
- Using estrogen-alone may increase your chance of getting dementia, based on a study of women 65 years of age or older
- Do not use estrogens with progestins to prevent heart disease, heart attacks, strokes or dementia
- Using estrogens with progestins may increase your chances of getting heart attacks, strokes, breast cancer, or blood clots
- Using estrogens with progestins may increase your chance of getting dementia, based on a study of women 65 years of age or older
- You and your healthcare provider should talk regularly about whether you still need treatment with Divigel®

Divigel® should not be used if you have unusual vaginal bleeding, currently have or have had certain cancers, including cancer of the breast or uterus, had a stroke or heart attack; currently have or have had blood clots, currently have or have had liver problems, have been diagnosed with a bleeding disorder, are allergic to Divigel® or any of its ingredients, or think you may be pregnant.

Tell your healthcare provider about all of your medical problems and the medicines you take, if you are going to have surgery or will be on bed rest, and if you are breastfeeding.

Call your healthcare provider right away if you get any of the following symptoms: new breast lumps, unusual vaginal bleeding, changes in vision or speech, sudden new severe headaches, or severe pains in your chest or legs with or without shortness of breath, weakness and fatigue.

Common side effects that may occur with Divigel® include headache; breast pain; irregular vaginal bleeding or spotting; stomach or abdominal cramps, bloating; nausea and vomiting; hair loss; fluid retention and vaginal yeast infection.

Serious but less common side effects include heart attack, stroke, blood clots, dementia, breast cancer, cancer of the uterus, ovarian cancer, high blood pressure, high blood sugar, gallbladder disease, liver problems, and enlargement of benign uterus tumors ("fibroids").

Alcohol-based gels are flammable. Avoid fire, flame or smoking until the gel has dried.

Please see Patient Information for Divigel® and talk to your healthcare provider. For more information, call 1-888-650-3789 or visit [www.divigel.com](http://www.divigel.com).

**You are encouraged to report negative side effects to Upsher-Smith Laboratories, Inc. at 1-855-899-9180, or to the FDA by visiting [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or calling 1-800-FDA-1088.**

©2012 Upsher-Smith Laboratories, Inc., Maple Grove, MN 55369