.8" Hinge

TOP FRONT (S1)

Aspi-Cor 81mg®

Adult Low Dose Aspirin 81 mg
Pain Reliever (NSAID)

ENTERIC

SAFETY COATED

120 Tablets

Actual Size

HEALTH PRODUCTS CORPORATION
Yonkers, N.Y. 10703
Product #6-45291

0 39686 01492 4

DO NOT EAT OR DRINK WHILE USING THIS PRODUCT
DO NOT EAT OR DRINK WHILE USING THIS PRODUCT

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*	Pain reliever

*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doc tor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
- hives
- facial swelling
- shock
- asthma (wheezing)

PACKAGE EXP. NOTE
MFG. DATE

PEEL HERE FOR MORE DRUG FACTS

7/16" unfinished area

.8" Hinge

TOP BACK (S2)

1.875
DIE SIZE
ACROSS

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause a severe stomach bleeding. The chance is higher if you:
- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if
- you have a history of stomach problems, such as heartburn
- stomach bleeding warning applies to you
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for
- gout
- diabetes
- arthritis

Stop use and ask a doctor if
- you get ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

4.6875

.8" Hinge

TOP BASE (S3)

SAI 139-4 40 1.8

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless clearly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients

black iron oxide, colloidal silicon dioxide, corn starch, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, polydextrose, polyethylene glycol, propylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

844-222-0990

Existing Tool: TBA
Size: 1.875 x 4.6875 with .0625"cr
Date: 03-16-15 new size