



2021 Cataract and Refractive  
**Product Catalog**



**Alcon**



Customer Service is available to assist you from 7:00 AM to 7:00 PM Central Time, Monday through Friday.

**BY TELEPHONE**

Call TOLL FREE 1-800-TO-ALCON (1-800-862-5266)

**BY FAX**

We offer TOLL FREE fax service. Fax your purchase order directly to:  
Alcon Customer Service Department  
1-800-241-0677

**ONLINE ORDERING**

Alcon offers the convenience and ease of online ordering and account access (invoices, order tracking, IOL inventory reporting, etc.). Please ask your customer service representative for additional details about MyAlconStore.com

**EDI**

Alcon offers and encourages the ease of automated ordering to customers with compatible systems configurations. Please ask your customer service representative for additional details on EDI.

**AUTOMATIC ORDER SCHEDULING SERVICE**

Regular automated orders can be scheduled based on interim dates most convenient for you. Ask your customer service representative about the benefits of our standing order program.

**PAYMENT TERMS**

Payment terms are net 30 days.





CATARACT EQUIPMENT

IRRIGATION & ASPIRATION

OVDs

CATARACT DISPOSABLES

SOLUTIONS

CUSTOM PAK®

SPECIALTY

PROCEDURAL EYE DROPS

IOLs

GLAUCOMA

REFRACTIVE SUITE

TETRACAINE  
Hydrochloride Ophthalmic  
Solution 0.5% STERI-UNITS®

ISOPTO® ATROPINE  
(atropine sulfate ophthalmic  
solution) 1%

ALCAINE®  
(proparacaine hydrochloride  
ophthalmic solution, USP) 0.5% 15 mL

FLUORESCITE®  
(fluorescein injection, USP) 10%

MYDRIACYL®  
(tropicamide ophthalmic  
solution, USP) 1%

CYCLOMYDRIL®  
(cyclopentolate hydrochloride and phenylephrine  
hydrochloride ophthalmic solution)

CYCLOGYL®  
(cyclopentolate hydrochloride  
ophthalmic solution, USP)

**ALCAINE®**

(proparacaine hydrochloride  
ophthalmic solution, USP)  
0.5% 15 mL



Available direct from Alcon

15 mL - 0.5%

- **ALCAINE®** Solution 0.5% is a topical local anesthetic indicated for corneal anesthesia of short duration (eg, during tonometry, gonioscopy, and removal of corneal foreign bodies), and for short corneal and conjunctival procedures
- ALCAINE® Solution 0.5% should not be used to treat patients with known hypersensitivity to any ingredient in the preparation
- 15-mL bottle available by the unit directly from Alcon; to order, please call (800) 862-5266 or visit MyAlconStore.com
- 15-mL bottle available by the unit through your authorized wholesaler or distributor

**EACH mL OF ALCAINE® SOLUTION 0.5% CONTAINS:**

**Active:** proparacaine hydrochloride 5 mg 0.5%

**Preservative:** benzalkonium chloride (0.01%)

**Inactive:** glycerin and purified water; pH may be adjusted with hydrochloric acid and / or sodium hydroxide

Bottle must be stored in unit carton to protect contents from light.

Store bottles under refrigeration at 2°C to 8°C (36°F-46°F).

**INDICATION AND DOSING**

ALCAINE® Solution is indicated for procedures in which a topical ophthalmic anesthetic is indicated: corneal anesthesia of short duration, e.g., tonometry, gonioscopy, removal of corneal foreign bodies, and for short corneal and conjunctival procedures.

**DOSAGE AND ADMINISTRATION**

**Usual Dosage:** Removal of foreign bodies and sutures, and for tonometry: 1 to 2 drops (in single instillations) in each eye before operating.

**Short Corneal and Conjunctival Procedures:** 1 drop in each eye every 5 to 10 minutes for 5 to 7 doses.

**IMPORTANT SAFETY INFORMATION**

**CONTRAINDICATIONS**

ALCAINE® Solution should be considered contraindicated in patients with known hypersensitivity to any of the ingredients of this preparation.

**WARNINGS AND PRECAUTIONS**

**NOT FOR INJECTION—FOR TOPICAL OPHTHALMIC USE ONLY.** Prolonged use of a topical ocular anesthetic is not recommended. It may produce permanent corneal opacification with accompanying visual loss.

**ADVERSE REACTIONS**

Occasional temporary stinging, burning and conjunctival redness may occur with the use of proparacaine. A rare, severe, immediate-type, apparently hyperallergic corneal reaction characterized by acute, intense and diffuse epithelial keratitis, a gray, ground glass appearance, sloughing of large areas of necrotic epithelium, corneal filaments and, sometimes, iritis with descemetitis has been reported. Allergic contact dermatitis from proparacaine with drying and fissuring of the fingertips has also been reported.

For additional information on ALCAINE® Solution, please [click here](#) for the full Prescribing Information.