

Drug Facts (continued)

- for adults and children 12 years of age and over
- children under 12 years of age: do not use; it is not known if the drug works or is safe in children under 12

Before first use (new inhaler)
activate new inhaler by shaking then spraying into air 4 separate times

each time you dose

- remove red cap (if attached)
- shake then spray into the air 1 time
- exhale completely, place inhaler in mouth
- inhale deeply while pressing down on top of inhaler, then continue the deep breath
- hold breath as long as possible, exhale
- wait 1 minute. If symptoms not relieved, take a second inhalation by repeating steps above.

after use

- wait at least 4 hours between doses
- do not use more than 8 inhalations in 24 hours
- wash inhaler after each day of use. Run water through the mouthpiece for 30 seconds.

Other information

- store at room temperature, between 15-25°C (59-77°F)
- contains no sulfites
- keep this label and enclosed materials. They contain important additional information.

Inactive ingredients

dehydrated alcohol (1%), hydrofluoroalkane (HFA 134a), polysorbate 80, thymol

Questions or comments?

call 1-8 Primatene or 1-877-462-8363
Mon. - Fri. 7 am - 5 pm PST



Visit www.primatene.com
See Insert and Side Panels
For Special Directions On:

- Activating Your New Inhaler
- Dosing with Your New Inhaler
- Using Spray Indicator

Primatene[®] MIST

Epinephrine Inhalation Aerosol
0.125 mg per spray
Bronchodilator

For Oral
Inhalation Only

For **TEMPORARY** relief
of **MILD** symptoms
of **INTERMITTENT** asthma



Suspension:
Shake then Spray
into the air before
each inhalation.

NEW FORMULATION:
See Important Usage
Information on Insert
and on Side Panels

160 Metered Sprays
Net Weight: 11.7g



Manufactured by:
Armstrong Pharmaceuticals, Inc.
Canton, MA 02021
© 2018 Made in USA



U5530R

Open Here

Drug Facts

Active ingredient <i>(in each spray)</i>	Purpose
Epinephrine 0.125 mg.....	Bronchodilator

Uses
for temporary relief of mild symptoms of intermittent asthma

- wheezing
- tightness of chest
- shortness of breath

Warnings
For oral inhalation only
Asthma alert: Because asthma may be life threatening, see a doctor if you

- are not better in 20 minutes
- get worse
- need more than 8 inhalations in 24 hours
- have more than 2 asthma attacks in a week

These may be signs that your asthma is getting worse.

Do not use

- unless a doctor said you have asthma
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs taken for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or a pharmacist before taking this product.

Ask a doctor before use if you have

- ever been hospitalized for asthma
- heart disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- thyroid disease
- seizures
- narrow angle glaucoma

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions
- taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain)

When using this product

- your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death.
- your risk of heart attack or stroke increases if you:
 - have a history of high blood pressure or heart disease
 - take this product more frequently or take more than the recommended dose.
- avoid foods or beverages that contain caffeine
- avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect
- avoid spraying in eyes
- contents under pressure. Do not puncture or incinerate.
- do not store near open flame or heat above 120°F (49°C). May cause bursting.

Stop use and ask a doctor if

- your asthma is getting worse (see asthma alert)
- you have difficulty sleeping
- you have a rapid heart beat
- you have tremors, nervousness, or seizure

If pregnant or breast-feeding, ask a health care professional before use
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- read the Consumer Information Insert for detailed directions on how to use this product
- do not use more than directed



U5530R